

ANNALES DE L'I. H. P., SECTION A

BRIAN DOLAN

Spinors on CP^2 in $d \geq 9$ supergravity

Annales de l'I. H. P., section A, tome 42, n° 4 (1985), p. 375-382

<http://www.numdam.org/item?id=AIHPA_1985__42_4_375_0>

© Gauthier-Villars, 1985, tous droits réservés.

L'accès aux archives de la revue « Annales de l'I. H. P., section A » implique l'accord avec les conditions générales d'utilisation (<http://www.numdam.org/legal.php>). Toute utilisation commerciale ou impression systématique est constitutive d'une infraction pénale. Toute copie ou impression de ce fichier doit contenir la présente mention de copyright.

NUMDAM

Article numérisé dans le cadre du programme
Numérisation de documents anciens mathématiques

<http://www.numdam.org/>

Spinors on CP^2 in $d \geq 9$ Supergravity

by

Brian DOLAN

Department of Natural Philosophy, University of Glasgow,
Glasgow G12 8QQ

ABSTRACT. — When the internal space of a $d \geq 9$ supergravity theory is $CP^2 \times M$, for some M , it is shown how to put spinors on CP^2 using spin^c structures, without the introduction of any extra fields. When M contains S^1 , the spinors couple to a $U(1)$ field with the number of right-handed spinors (v_+) greater than the number of left-handed spinors (v_-). When M contains S^2 , the spinors are $SU(2)$ doublets with $v_+ - v_- = -1$.

RÉSUMÉ. — Lorsque l'espace interne d'une théorie de supergravité à $d \geq 9$ est $CP^2 \times M$ pour un certain M , on montre comment on peut définir des spineurs sur CP^2 en utilisant des spin^c -structures sans introduire de champs supplémentaires. Lorsque M contient S^1 , les spineurs sont couplés à un champ $U(1)$, le nombre de spineurs droits (v_+) étant plus grand que le nombre de spineurs gauches (v_-). Lorsque M contient S^2 , les spineurs sont des doublets de $SU(2)$ avec $v_+ - v_- = -1$.

There are now many solutions known of the bosonic field equations of $d = 11$ and $d = 10$ supergravity theories. The basic idea is to use the extra fields, whose presence is dictated by supergravity arguments, to split the d dimensional metric up into the direct sum of a 4-dimensional metric on a non-compact spacetime and a $(d-4)$ dimensional metric on a compact, internal, manifold. The isometry group of the internal manifold is identified with the gauge group of nature.

In particular, some of these solutions involve an internal manifold whose topology is that of $CP^2 \times M$, where M is a compact space [1]-[5]. One

of the reasons that CP^2 attracts attention is that it admits an Einstein metric whose group of isometrics has the algebra of $SU(3)$, so that the total gauge algebra of such solutions is that of $SU(3) \times G_M$ where G_M is the group of isometries of the metric on M . However, these solutions suffer from the difficulty that CP^2 does not admit a spinor structure [5]-[8] and so such a solution will not admit spinors. One way round this problem is to put a $spin^c$ structure on CP^2 [6] [7] [8] i. e. put a $U(1)$ field on CP^2 and then add spinor fields which couple to the $U(1)$ field in such a way that they are defined consistently. This is the approach of Ref. [1], where the $U(1)$ fields is simply added to the theory, but this is somewhat contrary to the spirit of supergravity theories.

In this paper, a method is presented of consistently putting spinors on CP^2 without the introduction of any extra bosonic fields, other than those already present in the supergravity theory. The spinors can couple to an $SU(2)$ field as doublets on CP^2 or a $U(1)$ field as singlets on CP^2 , depending on the topology of M , though the total particle spectrum will depend on the spinor content of the other dimensions. This indicates a possible connection between the particles spectrum and the topology, as first suggested by Hawking and Pope [6].

In many solutions the equations of motion for the metric on the internal space, obtained by dimensional reduction from $d = 10$ or 11 , are simply Einstein's equations on the internal manifold ⁽¹⁾.

$$\mathbb{R}_{AB} \times e^{ABC} = \lambda \times e^C \quad (1)$$

where e^A , $A = 1, \dots, d-4$ are a basis of orthonormal one-forms. $e^{A \dots C} = e^A \wedge e^B \wedge \dots \wedge e^C$, \times is the Hodge dual in $d-4$ dimensions, \mathbb{R}_{AB} are the curvature 2-forms and λ is a constant coming from the curvature scalar of 4-dimensional space-time and any other fields present e. g. the 4-form field of $N = 1, d = 11$ supergravity. More generally, if the internal space is an outer product of two or more Einstein spaces, λ could be different for each space, e. g. if the internal space contains an S^1 , as in the dimensional reduction of $N = 1, d = 11$ down to $N = 2, d = 10$ [3].

In what follows we shall examine the case where the internal space is a product space $CP^2 \times M$, where M is a compact manifold with a Riemannian metric. The Fubini-Study metric on CP^2 is given by [7]

$$g_{CP^2} = \sum_{a=1}^4 e^a \otimes e^a$$

⁽¹⁾ For the particular case of the $N = 2, d = 10$ chiral supergravity theory, no such solution has been constructed. However, it seems reasonable to suppose that some solutions of this theory may involve Einstein metrics, of the form (1).

where e^a are given by

$$\begin{aligned} e^1 &= \frac{r\sigma^1}{(1+r^2)^{\frac{1}{2}}} & e^2 &= \frac{r\sigma^2}{(1+r^2)^{\frac{1}{2}}} \\ e^3 &= \frac{r\sigma^3}{(1+r^2)} & e^2 &= \frac{dr}{(1+r^2)} \end{aligned} \tag{2}$$

where $\sigma^i (i=1,2,3)$ are left invariant one forms on S^3 and obey $d\sigma^i = \varepsilon^{ijk}\sigma^j \wedge \sigma^k$. Explicitly

$$\begin{aligned} \sigma^1 &= \frac{1}{2} (\sin \psi d\theta - \sin \theta \cos \psi d\phi) \\ \sigma^2 &= -\frac{1}{2} (\cos \psi d\theta + \sin \theta \sin \psi d\phi) \\ \sigma^3 &= \frac{1}{2} (\cos \theta d\phi + d\psi) \end{aligned} \tag{3}$$

where $0 \leq \theta < \pi, 0 \leq \phi < 2\pi, 0 \leq \psi < 4\pi, 0 \leq r < \infty$. Of course, CP^2 is not covered by this single co-ordinate patch, one must choose different co-ordinates to cover the points $r = \infty$.

The orientation is given by $*1 = e^{1243}$ ($*$ is the Hodge operator on CP^2).

The unique, torsion free, metric compatible connection obtained from (2) is

$$\begin{aligned} \omega_{14} &= \frac{1}{r} e^1 & \omega_{23} &= \frac{1}{r} e^1 \\ \omega_{24} &= \frac{1}{r} e^2 & \omega_{31} &= \frac{1}{r} e^2 \\ \omega_{34} &= \frac{1-r^2}{r} e^3 & \omega_{12} &= \frac{1+2r^2}{r} e^3 \end{aligned} \tag{4}$$

The curvature 2-forms, $R_{ab} = d\omega_{ab} + \omega_{ac} \wedge \omega^c_b$, are

$$\begin{aligned} R_{14} &= e^{14} + e^{23} & R_{23} &= e^{14} + e^{23} \\ R_{24} &= e^{24} + e^{31} & R_{31} &= e^{24} + e^{31} \\ R_{34} &= 4e^{34} - 2e^{12} & R_{12} &= -2e^{34} + 4e^{12} \end{aligned}$$

and satisfy

$$\begin{aligned} R_{ab} \wedge *e^{abc} &= 12*e^c \\ \Rightarrow R_{ab} \wedge *e^{ab} &= 24*1 \end{aligned}$$

One can show that chiral spin $\frac{1}{2}$ fields cannot be put on CP^2 by using the Atiyah-Singer index theorem [7] to calculate the difference between

the number of right handed (v_+) and the number of left handed (v_-) massless spinors with the above curvature,

$$\begin{aligned} v_+ - v_- &= -\frac{1}{24} \frac{1}{8\pi^2} \int_{\text{CP}^2} R_{ab} \wedge R^{ab} \\ &= -\frac{1}{8} \end{aligned}$$

Since this is not an integer, one cannot put chiral spinors on CP^2 . However, if the spinors interact with a gauge field by minimal coupling then the Dirac operator is modified and the index theorem, for complex bundles, yields:

$$v_+ - v_- = -\frac{P}{8} + \frac{1}{8\pi^2} \int_{\text{CP}^2} \text{Tr}(G \wedge G) \quad (5)$$

where G is the field strength 2-form for the gauge field, which may be non-abelian, hence the trace. P is the dimension of the representation space on which the group acts. Thus, by suitable choice of gauge-group and G one can hope to adjust the R.H.S. of equation (5) so as to make it an integer, and indeed this is what a spin^c structure does.

Suitable gauge fields, G , can be constructed from the Levi-Civita connection (ref. [9]). The connection is an $\text{SO}(4)$ Lie Algebra valued one form and so can be decomposed into two $\text{SU}(2)$ Lie Algebra valued one forms

$$\begin{aligned} B_{\pm}^1 &= \frac{1}{2}(\omega^{41} \pm \omega^{23}) \\ B_{\pm}^2 &= \frac{1}{2}(\omega^{42} \pm \omega^{31}) \\ B_{\pm}^3 &= \frac{1}{2}(\omega^{43} \pm \omega^{12}) \end{aligned} \quad (6)$$

The resulting field strengths $G_{\pm}^i = dB_{\pm}^i + \varepsilon^{ijk} B_{\pm}^j \wedge B_{\pm}^k$ are (anti) self-dual. When this procedure is applied to the standard metric and connection on S^4 , one obtains the BPST instanton [10].

Take the positive sign in equations (6). Then using (4)

$$B^1 = B^2 = 0, \quad B^3 = \frac{3}{2} re^3 \equiv B \quad (7)$$

and the field is that of $\text{U}(1)$ rather than $\text{SU}(2)$. The field strength is

$$G = 3(e^{43} + e^{12}) = \frac{3}{4} G \quad (8)$$

It is self-dual and is, in fact the Kahler 2-form on CP^2 . To see how this field can be introduced into equations (1), we consider the case where the manifold, M , is the direct product of S^1 with an Einstein space. Then equa-

tions (1) split into three cases (a, b, c = 1, . . . , 4 label CP²; e⁵ is the volume element of S¹; μ = 1, . . . , d-9 label the rest of M)

$$\mathbb{R}_{AB} \wedge * e^{ABc} = \lambda_1 * e^c \tag{9}$$

$$\mathbb{R}_{AB} \wedge * e^{AB5} = \lambda_2 * e^5 \tag{10}$$

$$\mathbb{R}_{AB} \wedge * e^{AB\mu} = \lambda_3 * e^\mu \tag{11}$$

Now consider S¹ over CP² in the light of a standard Kaluza-Klein theory and use the Killing symmetry of the metric on S¹ to construct a U(1) field over CP².

Let e⁵ = dx⁵, 0 ≤ x⁵ < 2π, and consider a new, five dimensional metric

$$g_{(5)} = \sum_{a=1}^4 e^a \otimes e^a + (e^5 + B) \otimes (e^5 + B)$$

where B is any one form on CP² (not necessarily that of equation (7)), with no x⁵ dependence. This is the standard Kaluza-Klein ansatz, except that it is being applied purely to the internal space.

Of course the metric g₍₅₎ will not be a metric for CP² × S¹, in general, but for some other space with a different global topology.

The reduction of equations (9) and (10), with e⁵ replaced by (e⁵ + B) yields

$$d_4^* G = 0 \tag{12}$$

$$G \wedge * G = \frac{2}{3} (2\tilde{\lambda}_1 - \tilde{\lambda}_2) * 1 \tag{13}$$

$$R_{ab} \wedge * e^{abc} = \tilde{\lambda}_1 * e^c + \frac{1}{2} [G \wedge i^c * G - i^c G \wedge * G] \tag{14}$$

where $\tilde{\lambda}_i = \lambda_i - \mathcal{R}$ with \mathcal{R} the curvature scalar for the remaining dimensions, G = dB and i^c is the interior derivative, sending p-forms to (p - 1) forms and satisfying i^c(e^a) = δ^{ca}, i^c(e^{ab}) = δ^{ca}e^b - δ^{cb}e^a, i^c₄(e^a) = *₄e^{ac}.

If we now use the one-form derived from the connection on CP², equation (7), then (12) is automatically satisfied since G is self-dual. (13) is satisfied if $\tilde{\lambda}_2 = 2\tilde{\lambda}_1 - 27$ and (14) reduces to (9) with $\tilde{\lambda}_1 = 12$. (11) will be satisfied by suitably adjusting the curvature scalar of the remaining Einstein space, \mathcal{R} .

Substituting equation (8) into (5) with P = 1, the dimension of S¹ (the volume of CP² is π²/2)

$$v_+ - v_- = -\frac{1}{8} + \frac{9}{8} = 1$$

Thus this construction enables us to put spinors on CP² in a consistent fashion, with one more right-handed than left-handed spinor. In general the R.H.S. could be made into any positive integer, simply by rescaling B. This result is no more than a combination of the standard, 5 dimensional Kaluza-Klein ansatz, applied to the internal manifold, and the spin^c struc-

tures of ref. [6]. Its attractive feature is that it avoids the ad hoc introduction of an extra U(1) field.

Let us take the negative sign in equations (6). Then

$$B^1 = -\frac{1}{r} e^1, \quad B^2 = -\frac{1}{r} e^2, \quad B^3 = -\frac{(2+r^2)}{2r} e^3 \quad (15)$$

giving

$$G^1 = e^{41} - e^{23}, \quad G^2 = e^{42} - e^{31}, \quad G^3 = e^{43} - e^{12} \quad (16)$$

G is anti-self dual.

This field can be introduced in a consistent way if M is the direct product of S² with an Einstein space. Again, equations (1) split into three cases (a, b, c = 1, . . . , 4 label CP², m, n = 5, 6 label S², μ labels the rest of M)

$$\mathbb{R}_{AB} \wedge \ast e^{ABc} = \lambda_1 \ast e^c \quad (17)$$

$$\mathbb{R}_{AB} \wedge \ast e^{ABm} = \lambda_2 \ast e^m \quad (18)$$

$$\mathbb{R}_{AB} \wedge \ast e^{AB\mu} = \lambda_3 \ast e^\mu \quad (19)$$

Now use the Killing symmetries of the standard metric on S² to construct an SU(2) gauge field over CP² in the manner of a non-abelian Kaluza-Klein theory [11]-[14].

Let e⁵ = dα, e⁶ = sin α dβ 0 ≤ α < π, 0 ≤ β < 2π be a basis of orthonormal one forms on S², and consider a new, six dimensional metric

$$g_{(6)} = \sum_{a=1}^4 e^a \otimes e^a + \sum_{m=5}^6 \left[\left[e^m - 2 \sum_{j=1}^3 e^m(\tilde{K}_j) B^j \right] \otimes \left[e^m - 2 \sum_{k=1}^3 e^m(\tilde{K}_k) B^k \right] \right]$$

Here \tilde{K}_j are the three Killing vectors of the metric on S² and the numbers e^m(\tilde{K}_j) are the components of their metric dual one-forms in the e^m basis. B^j are SU(2) Lie Algebra valued one forms on CP², not necessarily those of equation (15), which are independent of α and β.

The Killing one forms K_j, metric dual to the Killing vectors, can be chosen as

$$K_1 = \sin \beta e^5 + \cos \beta \cos \alpha e^6$$

$$K_2 = \cos \beta e^5 - \sin \beta \cos \alpha e^6$$

$$K_3 = \sin \alpha e^6$$

and satisfy

$$dK_i = -\varepsilon_{ijk} K_j \wedge K_k$$

It is convenient also to define the quantities

$$h_{ij} = \sum_{m=5}^6 e^m(\tilde{K}_i) e^m(\tilde{K}_j)$$

which will be used for the contraction of group indices.

Note that

$$\delta^{jk}h_{jk} = 2 \quad \text{and} \quad \delta^{jk}K_j \wedge i^m \frac{*}{2}K_k = - \frac{*}{2}e^m \tag{20}$$

where $\frac{*}{2}$ is the Hodge dual on S² ($\frac{*}{2}1 = e^{56}$).

The reduction of equations (17) and (18), with e^m replaced by

$$e^m - 2 \sum_{j=1}^3 e^m(\tilde{K}_j)B^j$$

yields

$$R_{ab} \wedge \frac{*}{4}e^{abc} = (\lambda_1 - R_2)\frac{*}{4}e^c + 2[G^j \wedge i^c \frac{*}{4}G^k - i^c G^j \wedge \frac{*}{4}G^k]h_{jk} \tag{21}$$

$$(G^j \wedge \frac{*}{4}G^k)[3h_{jk}\frac{*}{2}e^m + K_j \wedge i^m(\frac{*}{2}K_k) + K_k \wedge i^m(\frac{*}{2}K_j)] = \frac{1}{2}(R_4 - \lambda_2)\frac{*}{4}1 \wedge \frac{*}{2}e^m \tag{22}$$

$$D(\frac{*}{4}G^j) = 0 \tag{23}$$

R₂ = 2 is the curvature scalar for S², R₄ = 24 is the curvature scalar for CP², and D($\frac{*}{4}G^j$) = d($\frac{*}{4}G^j$) + 2ε^{ijkl}B^k ∧ $\frac{*}{4}G^l$. If we now substitute (15) and (16), then (23) is automatically satisfied since Gⁱ is anti-self dual. Since G^j ∧ $\frac{*}{4}G^k$ = 2 $\frac{*}{4}$ 1δ^{jk} then (22) is satisfied (using 20) with λ₂ = 8 and (21) reduces to (17) which is satisfied provided λ₁ = 14. Finally (19) can be satisfied by adjusting λ₃ (for d=11, m has only one value as the remaining manifold must be S¹). This requires that λ₁λ₂ and λ₃ all be positive.

Now, putting (16) into (5) with G = τ_jG^j where τ_j are the generators of SU(2), normalised so that Tr(τ_jτ_k) = 2δ_{jk}, and P = 2, the dimension of S², one obtains

$$v_+ - v_- = -\frac{1}{4} - \frac{3}{4} = -1$$

Thus we now have one more left-handed spinor than right-handed ones. These spinors are SU(2) doublets, since P = 2.

The term $-\frac{3}{4}$ deserves some comment. It is, in fact, the topological charge of the SU(2) « instanton » field over CP² and being non-integral indicates a discontinuity in the fields. The construction (6) only gives us an ansatz for the Algebra. If we had used SO(3) as the group rather than SU(2), then the topological charge would have been -3 instead of $-\frac{3}{4}$. One factor of two arises because SU(2) is the double covering of SO(3) and another because the relevant quantity is the first Pontrjagin number for real bundles, as opposed to the second Chern number for complex bundles and, when

a complex bundle is regarded as real by forgetting the complex structure, these differ by a factor of two ⁽²⁾.

Fractionally charged instanton configurations have been studied extensively in the literature, for a review see ref. [15].

CONCLUSION

It has been shown that it is possible to consistently define spinors on CP^2 within the context of supergravity theories, without the introduction of any extra fields. This is achieved by minimally coupling the spinors to gauge fields which arise as extra components of the metric. If the internal manifold has the topology of $CP^2 \times M$ before the introduction of the extra components, then the spectrum of the spinors depends on the topology of M , being $SU(2)$ doublets if M contains an S^2 and $U(1)$ singlets if M contain an S^1 . If M is $S^2 \times S^1$, then a combination of both singlets and doublets could be achieved, there being more left-handed than right-handed doublets, and more right-handed than left-handed singlets. The overall particle spectrum of d -dimensional spinors will, however, depend on the spinor fields on the remaining $d-4$ dimensions apart from CP^2 , in particular the chirality content of the total particle spectrum could well be different from that of the spinors on CP^2 only.

REFERENCES

- [1] S. WATAMURA, *Phys. Lett.*, t. **129 B**, 1983, p. 188.
- [2] S. WATAMURA, *Spontaneous Compactification and CP^N* , University of Tokyo Preprint UT-Komaba, 83-15.
- [3] I. C. G. CAMPBELL and P. C. WEST, $N = 2, D = 10$ *Non-Chiral supergravity and its spontaneous Compactification*, King's College Preprint.
- [4] B. DOLAN, *Phys. Lett.*, t. **140 B**, 1984, p. 304.
- [5] L. CASTELLANI, L. J. ROMANS and N. P. WARNER, *A Classification of Compactifying Solutions for $d = 11$ Supergravity*, Cal. Tech. Preprint, 68-1055.
- [6] S. W. HAWKING and C. N. POPE, *Phys. Lett.*, t. **73 B**, 1978, p. 42.
- [7] T. EGUCHI, P. B. GILKEY and A. J. HANSON, *Phys. Rep.*, t. **66**, n° 6, 1980.
- [8] A. TRAUTMAN, *Intern. J. of Theor. Phys.*, t. **16**, 1977, p. 561.
- [9] M. F. ATIYAH, N. J. HITCHIN and I. M. SINGER, *Proc. Roy. Soc. Lond.*, t. **A 363**, 1978, p. 425-461.
- [10] A. A. BELAVIN, A. M. POLYAKOV, A. S. SCHWARZ and Yu. S. TYUPKIN, *Phys. Lett.*, t. **59 B**, 1975, p. 85.
- [11] Y. M. CHO and P. G. O. FREUND, *Phys. Rev.*, t. **D 12**, 1975, p. 1711.
- [12] Y. M. CHO and P. S. JANG, *Phys. Rev.*, t. **D 12**, 1975, p. 3789.
- [13] I. M. BENN and R. W. TUCKER, *A Riemannian Description of Non-Abelian Gauge Interaction*. Lancaster University Preprint, 1983.
- [14] S. M. BARR, *Phys. Lett.*, t. **129 B**, 1983, p. 303.
- [15] A. CHAKRABATI, *Classical Solutions of Yang-Mills Fields*, École Polytechnique Preprint.

(Manuscrit reçu le 20 août 1984)

⁽²⁾ I am grateful to M. F. Atiyah for clarification on this point.