



Cite this: *Chem. Commun.*, 2015, 51, 10107

Received 30th April 2015,
Accepted 14th May 2015

DOI: 10.1039/c5cc03625a

www.rsc.org/chemcomm

An oxothiosquaramide was shown to bind to chloride through hydrogen bonding interactions in DMSO and found to exhibit pH switchable chloride transport across phospholipid bilayers via an antiport transport mechanism.


The transport of anions across cellular membranes is an important biological function often mediated by specialised proteins embedded within cellular lipid bilayers.^{1,2} Intense effort is currently focussed on the development of small synthetic molecules that can mimic the role of such proteins. These have potential to treat diseases characterised by faulty anion transport^{3,4} or to induce cellular processes such as apoptosis.^{5,6} We have developed a number of simple hydrogen bonding receptors for anion transport containing urea, thiourea and squaramide groups.^{7–9} Transporters containing squaramides have proven particularly effective and mediate higher fluxes of anions than the analogous ureas or thioureas due to significantly higher anion affinities.^{9,10} More recently, we have shown that thiosquaramides display pH-dependent anion transport behaviour mimicking the gating/switching behaviour seen in biological systems.¹¹ Thiosquaramides are significantly more acidic than their oxosquaramide analogues, and at pH > 7 are present as deprotonated species, unable to bind to or transport anions. Lowering the pH reprotonates the thiosquaramide and switches on anion transport, providing a

pH switchable anion transport by an oxothiosquaramide†

Robert B. P. Elmes,‡§^a Nathalie Busschaert,‡¶^b Dawid D. Czech,^b Philip A. Gale*^b and Katrina A. Jolliffe*^a

potential method for targeting acidic environments in biological systems, without the need for external stimulation.

Given the significant differences in anion binding strength and p*K*_a between oxo- and thiosquaramides,^{11–13} we were intrigued by the possibility that a mixed oxothiosquaramide derivative might exhibit intermediate behaviour between the two extremes, with ‘switch-ON’ anion transport behaviour occurring at a different pH to that observed for **2**, which could be more relevant to switching in biological systems. We therefore report here the synthesis, chloride binding and transport properties of the mixed oxothiosquaramide **3**.



We previously found that conversion of oxosquaramide **1** to the thiosquaramide analogue **2** was most readily achieved using an excess of the crystalline P₄S₁₀-pyridine thionating agent¹⁴ in MeCN.¹¹ We reasoned that treatment of **1** with 1 equivalent of this reagent should yield a statistical mixture of compounds **1–3**,¹⁵ and found that this mixture was readily separable by flash chromatography to give **3** in 26% yield (see ESI†).

Mixed oxothiosquaramides have not previously been investigated as anion binding motifs. In order to evaluate the chloride binding ability of **3**, ¹H NMR spectroscopic titrations at 298 K in 0.5% H₂O in DMSO-*d*₆ were performed by addition of tetrabutylammonium chloride (TBACl). The data (Δδ) was fitted to a 1 : 1 binding model using Hyperquad[©] (see ESI† for fitted data).¹⁶ The association constants (*K*_a) of **1–3** in 0.5% H₂O in DMSO-*d*₆ with Cl[–] are summarised in Table 1. Both NH signals of **3** underwent significant downfield shifts (1.31 and 1.75 ppm) upon titration with chloride, indicating that both protons are involved in

^a School of Chemistry (F11), The University of Sydney, 2006 NSW, Australia.

E-mail: kate.jolliffe@sydney.edu.au; Fax: +61 2 9351 3329; Tel: +61 2 9351 2297

^b Chemistry, University of Southampton, Southampton, SO17 1BJ, UK.

E-mail: philip.gale@soton.ac.uk; Tel: +44 (0)23 8059 3332

† Electronic supplementary information (ESI) available: Synthesis and characterisation of the receptors, experimental details and figures relating to stability, association constant determination, stack plots and fitplots of the ¹H NMR titrations, pH-spectrophotometric titrations, various vesicle assays methods and Hill plots. The data underlying this communication is available online at <http://dx.doi.org/10.5258/SOTON/376697>. See DOI: 10.1039/c5cc03625a

‡ These authors contributed equally to this work.

§ Current address: Department of Chemistry, Maynooth University, National University of Ireland, Maynooth, Co. Kildare, Ireland.

¶ Current address: Department of Chemistry, Chemistry Research Laboratory, University of Oxford, 12 Mansfield Road, Oxford, OX1 3TA, UK.



Table 1 Summary of the chloride association constants K_a (M^{-1}) of receptors **1–3** in 0.5% H_2O in $DMSO-d_6$ at 298 K^a and pK_a values of receptors **1–3** (6×10^{-5} M) measured by spectrophotometric titrations in acetonitrile/water (9/1 v/v; in presence of 0.1 M TBAPF₆)

Receptor	K_a (M^{-1})	pK_a
1	458 ^b	9.8 ^b
2	60 ^b	5.3 ^b
3	470	6.6

^a Data fitted to a 1:1 model with errors <15%. ^b Value taken from ref. 11.

hydrogen bonding. Compound **3** exhibited similar affinity for Cl^- to that of its squaramide counterpart **1**, suggesting that incorporation of a single sulfur atom does not significantly alter the anion binding capacity of these compounds. This contrasts with the behaviour of thiosquaramide **2**, which binds only weakly to Cl^- in this solvent.

To compare the acidity of the NH protons in compounds **1–3**, pK_a values were determined by pH-spectrophotometric titrations in a mixture of acetonitrile/water (9/1 v/v; in the presence of 0.1 M TBAPF₆) as described previously.¹¹ Under these conditions, **3** was observed to undergo a single deprotonation event, characterised by a hypochromism at ~ 372 nm and a concomitant hyperchromism at ~ 396 nm (see Fig. 1). The titration profiles comparing **1–3** are shown in Fig. 1. The obtained pK_a values corresponding to a single deprotonation event are summarised in Table 1. As expected, the mixed oxothiosquaramide **3** has a pK_a intermediate between those of **1** and **2**. However, the pK_a is much closer to that of the dithionated derivative **2** than




Fig. 1 (a) Absorption spectra taken over the course of a pH-spectrophotometric titration of **3** (6×10^{-5} M) in an acetonitrile–water mixture (9/1 v/v; in presence of 0.1 M TBAPF₆). (b) Comparison plots of normalised absorbance for **1** (\blacktriangle), **2** (\blacksquare) and **3** (\blacklozenge) vs. pH. pK_a values were determined by fitting four parameter sigmoidal curves through the data points using Sigma Plot (Systat Software Inc., Chicago, IL, USA) with the point of inflexion corresponding to the pK_a .

to that of **1**. Importantly, the pK_a value of 6.6 suggests that **3** also exists in its deprotonated form at neutral pH and may therefore be useful as a pH-switchable anion transporter.

In order to evaluate the chloride transport properties of compound **3**, previously reported vesicle-based procedures¹⁷ were employed under conditions allowing direct comparison to the reported chloride transport properties of **1** and **2**.^{9,11} Unilamellar 1-palmitoyl-2-oleoyl-*sn*-glycero-3-phosphocholine (POPC) liposomes loaded with a NaCl solution (489 mM) buffered to pH 7.2 were prepared and suspended in an isotonic NaNO₃ solution (489 mM) buffered to pH 7.2. A small volume of a DMSO solution of **3** was added to these liposomes and the resulting chloride efflux was monitored using an ion selective electrode (ISE). An analogous series of experiments were also performed at pH 4.0. The results for oxothiosquaramide **3** (Fig. 2) indicate that, like the thiosquaramide analogue **2**, **3** possesses significantly enhanced anion transport ability at pH 4.0, compared to that observed at pH 7.2 (the results for the oxo-/thiosquaramides can be found in the ESI†). This is consistent with the pK_a value determined for **3** (6.6), which implies that at pH 7.2 a significant proportion of this receptor will be deprotonated and thus negatively charged. Interestingly, the transport behaviour of **3** at low loading (0.1 mol%) showed a more pronounced pH switching effect compared to higher loadings (1 mol%). This suggests that the ‘trigger’ to switch on anion transport may occur at a higher pH for **3** compared to that required for receptor **2**.

Hill analysis¹⁸ of the anion transport of **3** was performed at both pH 7.2 and pH 4.0, and revealed that **3** is a poor anion transporter at pH 7.2 but is an efficient anion transporter at pH 4.0 (Table 2). The mixed oxothiosquaramide derivative **3**




Fig. 2 Chloride efflux from POPC vesicles at pH 7.2 (empty symbols) and pH 4.0 (filled symbols) mediated by oxothiosquaramide **3** (1 mol% or 0.1 mol% with respect to lipid). POPC vesicles were loaded with a 489 mM NaCl solution buffered to pH 7.2 with 5 mM phosphate salts or to pH 4.0 with 5 mM citrate salts, and were suspended in a 489 mM NaNO₃ solution buffered to pH 7.2 with 5 mM phosphate salts or to pH 4.0 with 5 mM citrate salts. At the end of the experiment (300 s), detergent was added to lyse the vesicles and calibrate the ISE to 100% chloride efflux. Each point represents the average of a minimum of 6 independent trials. DMSO was used as a control.



Table 2 Summary of the anion transport behaviour of receptor 1–3

	EC50 at pH 7.2 ^a (mol%)	EC50 at pH 4.0 ^a (mol%)	Retention time ^b (min)	Apparent pK _a ^c
1	0.06 ^d	0.08 ^d	26.0	11.2 (±0.2)
2	0.68 ^d	<0.0125 ^d	26.6	6.0 (±0.2)
3	0.22	0.03	26.7	6.9 (±0.1)

^a Concentration of transporter (mol% with respect to lipid) needed to achieve 50% chloride efflux in 270 s from POPC vesicles filled with NaCl and buffered to pH 7.2 or pH 4.0. ^b Retention time of the compounds on a reversed-phase HPLC column. ^c pK_a value obtained by performing anion transport studies at various pH (see main text for details). ^d Value taken from ref. 11.

was found to exhibit chloride transport efficiencies intermediate between those of 1 and 2. At pH 7.2, transport activity was observed to follow the trend 1 > 3 > 2 which is consistent with the percentage of deprotonation as indicated by the trend in pK_a values. A similar dependence of anion transport on the percentage of protonated receptor at physiological pH has also been observed for prodigiosenes.¹⁹ At pH 4.0, transport activity follows the general trend 2 > 3 > 1 and is consistent with a combination of lipophilicity (as determined by HPLC), which is an important factor in transport ability, and the expected anion binding ability of the receptors in their fully protonated forms.^{20–23}

The chloride transport mediated by compound 3 may be either *via* the formation of membrane-spanning channels or *via* a mobile carrier mechanism. Oxo and thiosquaramides have previously been shown to function primarily *via* a mobile carrier mechanism^{9,11} and a series of vesicle-based experiments point towards 3 operating *via* a similar mechanism (see ESI†). Calcein leakage assays²⁴ revealed that the large highly anionic dye calcein does not leak from POPC vesicles at pH 7.2 or pH 4.0 in the presence of 3 (see ESI†), excluding the formation of pores or channels with large internal diameter and vesicle disruption induced by the transporters.

The charge imbalance created by transmembrane anion transport can be compensated by means of an antiport process or a symport process. The chloride efflux shown in Fig. 2 is therefore most likely the result of either a pure 'switch-on' Cl[−]/NO₃[−] antiport mechanism or the occurrence of a H⁺/Cl[−] symport process at lower pH (or a combination of both). To investigate which mechanism is dominant for receptor 3, POPC liposomes containing a NaCl solution and suspended in a Na₂SO₄ solution were prepared at pH 7.2 and pH 4.0. The double negatively charged sulfate anion is considerably more hydrophilic than nitrate ($\Delta G_{\text{hydr}}(\text{SO}_4^{2-}) = -1080 \text{ kJ mol}^{-1}$; $\Delta G_{\text{hydr}}(\text{NO}_3^-) = -300 \text{ kJ mol}^{-1}$)²⁵ and is therefore more challenging to transport across a lipid bilayer. It was found that under these conditions the efficiency of chloride transport was significantly reduced with a minimal amount of chloride being transported out of the liposomes at both pH 7.2 and pH 4.0 (see ESI†). While these results strongly suggest that an antiport mechanism is the dominant process for the chloride transport facilitated by 3, in principle HCl symport may account for some of the anion transport efficiency. A series of vesicle-based experiments were thus conducted using the pH-dependent fluorescence properties of 8-hydroxy-1,3,6-pyrenetrisulfonate (HPTS) in order to clarify this effect

(see ESI†).²⁴ Initially, POPC liposomes were prepared containing NaCl (489 mM) and HPTS (1 mM) buffered to pH 7.2 or pH 6.0, and suspended in a NaNO₃ (489 mM) solution buffered to pH 7.2 or pH 6.0. The results revealed that 3 is capable of effecting large changes in intravesicular pH even in the absence of a pH gradient (although to a lesser extent). The ability of 1 and 2 to change intravesicular pH was previously observed, however in the absence of a pH gradient the intravesicular pH did not change significantly upon the addition of these transporters (see ESI†).¹¹ Taken together, these results indicate that the chloride effluxes shown in Fig. 2 are mainly the result of Cl[−]/NO₃[−] antiport processes but that H⁺/Cl[−] symport (or OH[−]/Cl[−] antiport) may also be an underlying mechanism. In conclusion, it appears that the chloride transport ability of receptor 3 is mainly the result of a Cl[−]/NO₃[−] antiport mechanism that can be switched on by lowering the pH, while H⁺/Cl[−] symport may also be contributing to the overall transport ability of 3.

To probe whether the pH-switch of the thiosquaramides mirrors the pK_a values obtained through spectrophotometric titrations, the Cl[−]/NO₃[−] experiments depicted in Fig. 2 were repeated at a range of pHs for receptors 1–3 (see ESI† for details on the different pHs and buffers used). The exact position of the pH switch should correspond to the pK_a value of the receptor under the conditions of the anion transport experiments ('apparent pK_a').¹¹ In order to calculate the apparent pK_a of 3, we plotted the normalised initial rate of chloride efflux *versus* pH and fitted the data to a sigmoidal function (Fig. 3). The values obtained using this method are summarised in Table 2 and largely agree with the pK_a values determined above. These results clearly show that the chloride transport ability of 3 can be switched on at pH < 7, with the major switch occurring at pH ≈ 7.0. Fig. 3 also confirms that the chloride transport by the analogous (thio)squaramides 1 and 2 is also pH-dependent but in different




Fig. 3 pH dependence of the chloride transport ability of oxo/thio/mixed squaramides 1–3 (1 mol%). Chloride transport experiments were conducted as described in Fig. 2 but at various different pHs. The obtained data was fitted to an asymptotic function and the initial rate of chloride efflux (k_{ini}) was calculated (see ESI† for details) and plotted as a function of pH after normalising by the maximum rate achievable at pH 4.0. Apparent pK_a values could subsequently be estimated by fitting the obtained plot to the following sigmoidal function: $y = \text{max} + ((\text{min} - \text{max}) / (1 + 10^{(\text{pK}_a - x)}))$.



pH windows. Clearly, squaramide **1** displays potent transmembrane anion transport activity across a large pH range (pH 4–10), while thiosquaramide **2** is inactive above pH \approx 6 and oxothiosquaramide **3** shows activity over an intermediate pH range.

In summary we have reported the first use of a mixed oxothiosquaramide as an efficient chloride receptor and pH-switchable chloride transporter. Receptor **3** has similar anion binding properties to oxosquaramide **1**, forming strong complexes with chloride anions, as indicated by proton NMR titrations in DMSO-*d*₆. However, its p*K*_a is closer to that of thiosquaramide **2**. Vesicle studies indicated that the *trans*-membrane anion transport ability of **3** is switched OFF at pH > 7 and switched ON at pH < 7. Importantly, the effect of the addition of one sulfur to the squaramide structure is large enough to cause the desired pH switch for transport while addition of a second sulfur moves the switch to lower pH. This paper provides further evidence for the controllable and switchable anion transport nature of synthetic receptors containing the squaramide moiety. We are currently studying these and related systems for the development of future biologically active anion transporters.

We thank the EPSRC (EP/J009687/1) and the ARC (DP140100227) for funding. PAG thanks the Royal Society and the Wolfson Foundation for a Royal Society Wolfson Research Merit Award. PAG and KAJ thank the European Cooperation in Science and Technology (COST) action CM1005 “Supramolecular Chemistry in Water” for support.

Notes and references

- J. T. Davis, O. Okunola and R. Quesada, *Chem. Soc. Rev.*, 2010, **39**, 3843–3862.
- F. M. Ashcroft, *Ion Channels and Disease*, Academic Press, San Diego, 2000.
- P. A. Gale, R. Pérez-Tomás and R. Quesada, *Acc. Chem. Res.*, 2013, **46**, 2801–2813.
- N. Busschaert and P. A. Gale, *Angew. Chem., Int. Ed.*, 2013, **52**, 1374–1382.
- S.-K. Ko, S. K. Kim, A. Share, V. M. Lynch, J. Park, W. Namkung, W. Van Rossom, N. Busschaert, P. A. Gale, J. L. Sessler and I. Shin, *Nat. Chem.*, 2014, **6**, 885–892.
- J. T. Davis, in *Topics in Heterocyclic Chemistry*, ed. P. A. Gale and W. Dehaen, Springer, Berlin, Heidelberg, 2010, vol. 24, pp. 145–176.
- L. E. Karagiannidis, C. J. E. Haynes, K. J. Holder, I. L. Kirby, S. J. Moore, N. J. Wells and P. A. Gale, *Chem. Commun.*, 2014, **50**, 12050–12053.
- C. J. E. Haynes, N. Busschaert, I. L. Kirby, J. Herniman, M. E. Light, N. J. Wells, I. Marques, V. Felix and P. A. Gale, *Org. Biomol. Chem.*, 2014, **12**, 62–72.
- N. Busschaert, I. L. Kirby, S. Young, S. J. Coles, P. N. Horton, M. E. Light and P. A. Gale, *Angew. Chem., Int. Ed.*, 2012, **51**, 4426–4430.
- S. J. Edwards, H. Valkenier, N. Busschaert, P. A. Gale and A. P. Davis, *Angew. Chem., Int. Ed.*, 2015, **54**, 4592–4596.
- N. Busschaert, R. B. P. Elmes, D. D. Czech, X. Wu, I. L. Kirby, E. M. Peck, K. D. Hendzel, S. K. Shaw, B. Chan, B. D. Smith, K. A. Jolliffe and P. A. Gale, *Chem. Sci.*, 2014, **5**, 3617–3626.
- X. Ni, X. Li, Z. Wang and J.-P. Cheng, *Org. Lett.*, 2014, **16**, 1786–1789.
- R. B. P. Elmes, P. Turner and K. A. Jolliffe, *Org. Lett.*, 2013, **15**, 5638–5641.
- J. Bergman, B. Pettersson, V. Hasimbegovic and P. H. Svensson, *J. Org. Chem.*, 2011, **76**, 1546–1553.
- M. B. Onaran, A. B. Comeau and C. T. Seto, *J. Org. Chem.*, 2005, **70**, 10792–10802.
- P. Gans, A. Sabatini and A. Vacca, *Talanta*, 1996, **43**, 1739–1753.
- B. D. Smith and T. N. Lambert, *Chem. Commun.*, 2003, 2261–2268.
- A. V. Hill, *Biochem. J.*, 1913, **7**, 471–480.
- E. Marchal, S. Rastogi, A. Thompson and J. T. Davis, *Org. Biomol. Chem.*, 2014, **12**, 7515–7522.
- N. Busschaert, S. J. Bradberry, M. Wenzel, C. J. E. Haynes, J. R. Hiscock, I. L. Kirby, L. E. Karagiannidis, S. J. Moore, N. J. Wells, J. Herniman, G. J. Langley, P. N. Horton, M. E. Light, I. Marques, P. J. Costa, V. Félix, J. G. Frey and P. A. Gale, *Chem. Sci.*, 2013, **4**, 3036–3045.
- H. Valkenier, C. J. E. Haynes, J. Herniman, P. A. Gale and A. P. Davis, *Chem. Sci.*, 2014, **5**, 1128–1134.
- N. Busschaert, M. Wenzel, M. E. Light, P. Iglesias-Hernández, R. Pérez-Tomás and P. A. Gale, *J. Am. Chem. Soc.*, 2011, **133**, 14136–14148.
- V. Saggiomo, S. Otto, I. Marques, V. Félix, T. Torroba and R. Quesada, *Chem. Commun.*, 2012, **48**, 5274–5276.
- V. Sidorov, F. W. Kotch, G. Abdrakhmanova, R. Mizani, J. C. Fettinger and J. T. Davis, *J. Am. Chem. Soc.*, 2002, **124**, 2267–2278.
- Y. Marcus, *J. Chem. Soc., Faraday Trans.*, 1991, **87**, 2995–2999.

