

Ca²⁺-Regulatory Muscle Proteins in the Alcohol-Fed Rat

Kay Ohlendieck, Shona Harmon, Michael Koll, Alistair G. Paice, and Victor R. Preedy

Alcoholic myopathy is characterized by muscle weakness and difficulties in gait and locomotion. It is one of the most prevalent skeletal muscle disorders in the Western hemisphere, affecting between 40% and 60% of all chronic alcohol misusers. However, the pathogenic mechanisms are unknown, although recent studies have suggested that membrane defects occur as a consequence of chronic alcohol exposure. It was our hypothesis that alcohol ingestion perturbs membrane-located proteins associated with intracellular signalling and contractility, in particular those relating to calcium homeostasis. To test this, we fed male Wistar rats nutritionally complete liquid diets containing ethanol as 35% of total dietary energy. Controls were pair-fed identical amounts of the same diet in which ethanol was replaced by isocaloric glucose. At the end of 6 weeks, rats were killed and skeletal muscles dissected. These were used to determine important ion-regulatory skeletal muscle proteins including sarcalumenin (SAR), sarcoplasmic-endoplasmic reticulum Ca²⁺-adenosine triphosphatase (ATPase) (SERCA1), the junctional face protein of 90 kd (90-JFP), α_1 - and α_2 -dihydropyridine receptor (α_1 -DHPR and α_2 -DHPR), and calsequestrin (CSQ) by immunoblotting. The relative abundance of microsomal proteins was determined by immunoblotting using the enhanced chemiluminescence (ECL) technique. The data showed that alcohol-feeding significantly reduced gastrocnemius and hind limb muscle weights ($P < .05$ in both instances). Concomitant changes included increases in the relative amounts of SERCA1 ($P < .05$) and Ca²⁺-ATPase activity ($P < .025$). However, there were no statistically significant changes in either SAR, 90-JFP, α_1 -DHPR or α_2 -DHPR ($P > .2$ in all instances). Reductions in CSQ were of marginal significance ($P = .0950$). We conclude that upregulation of SERCA1 protein and Ca²⁺-ATPase activity may be an adaptive mechanism and/or a contributory process in the pathology of alcohol-induced muscle disease.

© 2003 Elsevier Inc. All rights reserved.

CHRONIC ALCOHOL DEPENDENTS may not only suffer from cirrhosis-induced impairment of liver function, cardiac complications, and various alcohol-based psychotic disorders, but up to two thirds have severe skeletal muscle dysfunction, ie, alcohol-induced muscle disease.^{1,2} Chronic alcoholic muscle disease is characterized by defects in contractility, muscle weakness, and loss of muscle mass. However, the mechanisms for this pathology are unknown.^{1,2} Nevertheless, 3 lines of evidence from our group suggest that disturbances in the membrane domain may occur as a consequence of alcohol exposure. The first pertains to adduct formation, which occurs especially within the sarcolemmal region of muscle.^{3,4} Secondly, reductions in ribosomal protein in alcohol-exposed muscle occurs preferentially in the subsarcolemmal region in contrast with the more moderate changes in the intermyofibrillar ribosomes.⁵ Finally, as a consequence of oxidative stress, alcohol increases the concentration of the membrane located hydroperoxides, 7 α -hydroperoxycholest-5-en-3 β -ol (7 α -OOH) and 7 β -hydroperoxycholest-5-en-3 β -ol (7 β -OOH).⁶ Furthermore, studies from other groups have also shown disturbances

in the membrane domain of muscle, such as alcohol-induced decreases in Ca²⁺ transients in cultured human myotubes.⁷

In skeletal muscle fibers, excitation-contraction coupling and muscle relaxation is regulated by changes in cytosolic Ca²⁺ levels.⁸ Ca²⁺ cycling through the sarcoplasmic reticulum is mediated by the junctional ryanodine receptor Ca²⁺ release channel complex and the Ca²⁺-adenosine triphosphatase (ATPase) units.⁹ Following voltage sensing by the α_1 -subunit of the dihydropyridine receptor,¹⁰ direct physical triad receptor coupling triggers Ca²⁺ efflux through the ryanodine receptor.¹¹ The energy-dependent reuptake of Ca²⁺ ions is achieved by Ca²⁺ pumps of the longitudinal tubules and the terminal cisternae.¹² Within the lumen of the sarcoplasmic reticulum, the high-capacity Ca²⁺-binding protein calsequestrin (CSQ)¹³ acts as an ion reservoir and endogenous regulator of Ca²⁺ efflux.¹⁴ Other luminal Ca²⁺-binding proteins are represented by sarcalumenin (SAR)¹⁵ and calreticulin (CAL).¹⁶

Since ethanol causes biochemical abnormalities within the membrane domain of muscle,^{3,7} it was our hypothesis that chronic alcohol ingestion perturbs key components of calcium homeostasis. This is a reasonable supposition, since abnormal Ca²⁺ handling is involved in various skeletal muscle disorders.¹⁷ Such damage may contribute to potential pathways of alcohol-induced tissue damage including functional deficiencies such as weakness (for example see, Odermatt et al¹⁸). To determine the relative expression of key Ca²⁺-regulatory muscle proteins, we have performed immunoblotting of microsomal proteins derived from control and chronic alcohol-exposed skeletal muscle fibers. To unequivocally identify the various muscle proteins, established antibodies were used for highly specific immunodecoration of the α_1 - and α_2 -subunit of the transverse-tubular DHPR, the Ca²⁺-binding proteins CSQ of the terminal cisternae and SAR of the longitudinal tubules, as well as the Ca²⁺-ATPase. For the internal standardization of immunoblots, the expression of the abundant dystrophin-associated surface marker β -dystroglycan was employed.¹⁹

From the Department of Biology, National University of Ireland, Maynooth, Co. Kildare; Department of Pharmacology, Conway Institute of Biomolecular and Biomedical Research, University College Dublin, Belfield, Dublin, Ireland; and the Department of Nutrition and Dietetics, King's College London, London, UK.

Submitted May 16, 2002; accepted January 7, 2003.

Supported by project grants from Enterprise Ireland and the European Commission (FMRX-CT960032; RTN2-2001-00337).

Address reprint requests to Victor R. Preedy, PhD, DSc, FRCPATH, Department of Nutrition and Dietetics, King's College London, 150 Stamford St, London SE1 9NN, UK.

© 2003 Elsevier Inc. All rights reserved.

0026-0495/03/5209-0040\$30.00/0

doi:10.1016/S0026-0495(03)00063-5

Table 1. Composition and Energy Values of Control and Alcohol-Containing Liquid Diets

	Content (g)	Total kcal	Total kJ
Alcohol diet			
Water	126	0.0	0.0
Fresubin	600	586	2,448
Added glucose	0.0	0.0	0.0
Casein	15.0	60	251
Orovite (sucrose)	2.46	9	39
Alcohol (61.8 ml)	49.08	348	1,457
Total	1,063	1,003	4,194
Control diet			
Water	126	0.0	0.0
Fresubin	600	586	2,448
Added glucose	93	349	1,458
Casein	15.0	60	251
Orovite (sucrose)	2.46	9	39
Alcohol	0	0	0
Total	1,106	1,004	4,196

NOTE. Calorific values were calculated as follows: ethanol = 7.1 kcal/g; carbohydrate = 3.8 kcal/g; protein = 4.0 kcal/g; fat = 9.0 kcal/g; and 4.18 kJ was assumed to be = 1.0 kcal.

MATERIALS AND METHODS

Materials

Protease inhibitors and acrylamide stock solutions were obtained from Boehringer-Mannheim (Lewis, UK). Chemiluminescence substrates were purchased from Pierce and Warriner (Chester, UK). Im-

mobilon NC nitrocellulose membranes were from Millipore (Bedford, MA). Commercially available primary antibodies were obtained from Affinity Bioreagents, Golden, CO: monoclonal antibody (mAb) VIIIID1₂ to CSQ, mAb 20A to the α_2 -subunit of the DHPR, mAb XIIC4 to SAR, mAb VF1c to the 90-kd junctional face protein, mAb IHH11 to the fast SERCA1 isoform of the sarcoplasmic endoplasmic reticulum Ca²⁺-ATPase, and pAb PA3-900 to calreticulin. Peroxidase-conjugated secondary antibodies were purchased from Chemicon (Temecula, CA). Monoclonal antibody IIID5 to the α_1 -subunit of DHPR was a generous gift from Dr Kevin P. Campbell (Howard Hughes Medical Institute, University of Iowa, Iowa City, IA). All other chemicals were of analytical grade and purchased from Sigma Chemical Co (Dorset, UK).

Animal Treatments

Male Wistar rats were obtained from accredited commercial suppliers at about 60 g body weight then maintained for approximately 1 week in a Home Office-approved animal house until they weighed approximately 0.1 kg. At this stage, rats were ranked and divided into 2 groups of equal mean body weight, and designated to either controls or ethanol-treated groups. The treated rats were subjected to a Lieber-DeCarli alcohol-feeding regimen in which animals were fed a nutritionally complete liquid diet containing 35% of total dietary energy as ethanol.^{20,21} Controls were pair-fed identical volumes of the same diet in which ethanol was replaced by isocaloric glucose^{20,21} (Tables 1 and 2). After approximately 6 weeks, animals were killed by decapitation. Skeletal muscles were dissected and limb muscle mass determined. Dissected muscle was frozen in liquid nitrogen (-196°C) and stored at -70°C until use. For analysis, either the gastrocnemius or hind limb musculature was used. A group of rats were fed on a standard solid

Table 2. Comparison of Chow Diet and Liquid Diets Used for Chronic Alcohol Feeding

	Chow Diet (per kg)	Liquid Control (glucose) Diet (per kg)	Liquid Alcohol Diet (per kg)
Nutrients			
Energy	12.5 MJ (2,988 kcal)	3,816 kJ (912 kcal)	3,812 kJ (911 kcal)
Fat/oil	208 g	18 g	18 g
Protein	174 g	34 g	34 g
Carbohydrates	495 g	142 g	107 g
of which glucose	n/a	83 g	0
of which alcohol	n/a	0	49 g
Water	100 g	779 g	779 g
Vitamins and selected minerals			
Vitamin A	5,614 μg	243 μg	243 μg
Vitamin D	0.129 μg	136 μg	136 μg
Vitamin E	103 μg	4 μg	4 μg
Vitamin K	185 μg	26.9 μg	26.9 μg
Vitamin B ₁	15.8 mg	1.16 mg	1.16 mg
Vitamin B ₂	13.6 mg	1.46 mg	1.46 mg
Niacin	78.7 mg	12.9 mg	12.9 mg
Vitamin B ₆	18.5 mg	1.54 mg	1.54 mg
Vitamin B ₁₂	83.5 μg	1.07 μg	1.07 μg
Pantothenic acid	26.1 mg	1.88 mg	1.88 mg
Biotin	0.359 mg	53.7 mg	53.7 mg
Folic acid	4.4 mg	54 μg	54 μg
Ascorbic acid	8.0 mg	51.1 mg	51.1 mg
Calcium	8.0 g	322 mg	322 mg
Iron	110 mg	540 μg	540 μg
Iodine	0.60 mg	57.1 mg	57.1 mg

NOTE. Data to show differences between solid chow and liquid diets for selective nutrients. Chow CRM pellets diets (SDS, Witham, UK) were used. The liquid diet was made up using Fresubin (Fresenius Kabi Ltd, Mano Park, UK), glucose and casein (both Merck BDH, Lutterworth, UK), Orovite 7 (Boots Chemist), 99.9 % alcohol (Hospital Pharmacy), and water.

chow diet (ie, pelleted) ad libitum for the duration of the study for comparative purposes. These were from the same batch of rats selected for liquid feeding with either glucose or ethanol.

Liquid Diets

Fresh liquid diets used for the 6-week chronic ethanol feeding experiment were prepared on a daily basis according to the recipe described in Table 1. A food blender was used to thoroughly mix the ingredients. To prevent the possibility of ethanol precipitating the protein in the alcohol diet, absolute ethanol was the last ingredient to be added carefully, and contents were then thoroughly stirred during the addition. The composition of the diet is given in Table 2. Control and alcohol-containing diets were isolipidic, isonitrogenous, and isoenergetic (Tables 1 and 2).

The compositions of the solid chow and liquid diets are entirely different with respect to macro- and micronutrients such as vitamins, minerals, and other essential and nonessential dietary elements (Table 2). Thus, any comparisons between the rats fed ad libitum on the solid chow and animals fed on the 2 liquid diets will be potentially difficult to interpret and may be due to any one or several of over 100 dietary components (see also Table 2). For this reason, detailed and comparative biochemical analysis of the muscle from solid chow and liquid fed rats was not performed (for further details on the feeding protocols, see Preedy et al²¹ and Lieber and DeCarli²²). Rather, representative blots are presented to illustrate the fact that the liquid diet-feeding regimen had no overt effect on the gross features of immunoblotting (electrophoretic separation, migration, etc).

Subcellular Fractionation

For subcellular fractionation, muscle specimens were homogenized in 5 vol of 20 mmol/L Tris-maleate, pH 7.0, 3 mmol/L EGTA, 10% (wt/vol) sucrose using an Ultra-Turrax T25 (IKA-Labor Technik, Staufen, Germany) and subsequently crude microsomal membranes prepared by standard differential centrifugation.¹⁹ All preparative steps were performed at 0 to 4°C in a cold room and all buffers were supplemented with a protease inhibitor cocktail consisting of 1 mmol/L EDTA, 0.2 mmol/L pepabloc, 1.4 μmol/L pepstatin A, 0.15 μmol/L aprotinin, 0.3 μmol/L E-64, 1 μmol/L leupeptin, and 0.5 μmol/L soybean trypsin inhibitor to minimize protein degradation.^{23,24} Protein concentration of resuspended microsomal pellets was determined by the method of Bradford²⁵ using bovine serum as standards. To determine potential differences in the Ca²⁺-ATPase activity between control and ethanol-treated microsomal preparations, a direct colourimetric assay was employed.²⁶

Gel Electrophoresis

For electrophoretic separation, protein samples were mixed with an equal volume of double strength sample buffer containing 3% (wt/vol) sodium dodecyl sulfate (SDS), 8 mol/L urea, 2 mol/L thiourea Bradford, 0.05% (wt/vol) bromophenol blue, and 50 mmol/L Tris-HCl, pH 6.8, 75 mmol/L dithiothreitol (DTT),²⁷ and 20 μg protein run per lane using 7% (wt/vol) separation gels at a constant voltage for 280 Vh in a Mini-Protean II gel system from Bio-Rad Laboratories (Hemel Hempstead, UK).

Immunoblot Analysis

To determine alcohol-induced effects on the isoform expression level of key Ca²⁺-regulatory proteins, muscle proteins separated by gel electrophoresis were transferred to Immobilon NC membranes according to previously described methods²⁸ using a Bio-Rad Mini-Protean II blotting system (Bio-Rad Laboratories). Blocking, washing, and incubation with primary and peroxidase-conjugated secondary antibodies were performed by established methods. Immunodecoration was eval-

uated by enhanced chemiluminescence (ECL). Densitometric scanning of ECL-blots was performed on a Molecular Dynamics 300S computing densitometer (Sunnyvale, CA) using ImageQuant V3.0 software.²⁴

Statistics

All data are displayed as means ± SEM (n = 5 to 8). For differences between glucose-fed and alcohol-treated rats on liquid diets, statistical evaluation was performed by Student's *t* test for paired samples. For differences between ad libitum-fed rats on the solid pelleted chow diet, and the other 2 groups of rats on liquid diets, post-hoc statistical evaluation was performed by Student's *t* test for unpaired samples.

RESULTS

Comparison Between Chow-Fed Rats and Liquid-Fed Groups

Mean body weights were as follows with ensuing data presented as means ± SEM with *P* values pertaining to differences between ad libitum-fed rats on a solid chow diet (n = 8) versus rats fed on a nutritionally complete liquid diet containing approximately one third of total dietary energy as either glucose (n = 8) or ethanol (n = 8). Body weights (g) were as follows: ad libitum solid chow-fed, 393 ± 9; glucose-fed rats on a liquid diet, 194 ± 4 (*P* < .001); and ethanol-treated rats on a liquid diet, 184 ± 3 (*P* < .001). Gastrocnemius weights (right muscle, g) were as follows: ad libitum solid chow-fed, 1.715 ± 0.025; glucose-fed rats on a liquid diet, 1.040 ± 0.038 (*P* < .001); and ethanol-treated rats on a liquid diet, 0.781 ± 0.071 (*P* < .001). Hind limb weights (right musculature, g) were as follows: ad libitum solid chow-fed, 9.361 ± 0.274 g; glucose-fed rats on a liquid diet, 5.073 ± 0.195 (*P* < .001); and ethanol-treated rats on a liquid diet, 3.981 ± 0.159 (*P* < .001). These results reaffirms the supposition that alcohol-fed rats have muscle weights lower than those from rats fed on a solid diet ad libitum and support the notion that a pair-feeding regimen should be used to investigate the putative effects of alcohol on muscle.

Muscle from ad libitum-fed rats on the solid chow diet were not subjected to detailed biochemical analysis as comparisons with rats on liquid diet are unsound. We reiterate the entirely different nature and composition of the solid chow diet in comparisons with the liquid feeds. In simple terms, any potential difference between muscles from rats fed on the solid or liquid diets may be due to any one of a multitude of dietary components (see Table 2).

On the other hand, it is also important to show that the liquid feeding protocol did not cause overt changes in muscle protein profiles (either via overt changes in relative abundance or molecular mass, etc.) and for this reason a general comparison between representative samples from ad libitum-fed rats on the solid chow, and rats on the glucose- or ethanol-containing liquid diets are displayed on the same gel. These gels show similar patterns of expression and molecular mass of immunopositive proteins in muscles from rats fed on solid or liquid diets.

Comparison Between Glucose-Fed Control and Alcohol-Treated Rats on Liquid Diets

Chronic ethanol feeding reduced body weight and skeletal muscle weight (Fig 1). There were almost identical propor-




Fig 1. Body and muscle weights in skeletal muscle of ethanol-fed rats. Rats were fed a nutritionally complete liquid diet containing ethanol as 35% of total calories. Controls were pair-fed the same diet in which ethanol was replaced by isocaloric glucose. At the end of the study, rats were killed and the muscles dissected out. Histograms represent the mean \pm SEM ($n = 8$ pairs). Notations above each histogram of the ethanol group refer to differences from the control groups. Differences between means were assessed with Student's *t* test for paired samples. For details on the changes in the muscle from rats on the solid chow diet, see Results.

tional changes in hind limb muscle and gastrocnemius consistent with previously reported changes in this model (Fig 1).

Since it is postulated that ethanol and/or its metabolites have

a modulatory effect on membrane fluidity and ion-regulatory membrane proteins, we determined the expression of key microsomal components of Ca^{2+} -handling from chronic alcohol exposed skeletal muscle. As illustrated in Fig 2, the Coomassie-stained protein band pattern of microsomal preparations from ad libitum-fed and glucose-fed control and myopathic muscle samples were relatively comparable. Since individual protein bands in Coomassie-stained gels may pertain to numerous molecular species, changes in specific isoforms of membrane proteins were identified using immunoblotting. Established antibodies to the α_1 - and α_2 -subunit of the dihydropyridine receptor (Figs 3 and 4, respectively), CSQ (Fig 5), SAR (Fig 6), the 90-kd junctional face membrane protein (Fig 7), the sarcoplasmic reticulum Ca^{2+} -ATPase (Fig 8), and the surface membrane marker β -dystroglycan (Fig 9) were employed. In conjunction with Fig 8, the enzyme activity of the sarcoplasmic reticulum Ca^{2+} pump was also determined (Fig 10).

Immunoblot analysis using monoclonal antibodies to the 2 main subunits of the transverse-tubular voltage sensor, the principal α_1 -subunit and the auxiliary α_2 -subunit of the DHPR revealed no alcohol-related changes in their expression. Neither the intensity of immunolabeling nor the relative position in comparison to molecular mass standard differed between normal and alcoholic myopathy specimens (Figs 3 and 4). The major Ca^{2+} -binding protein of the luminal sarcoplasmic reticulum, CSQ, seems to be only marginally affected by alcohol-induced modifications of muscle membrane systems (Fig 5). Evaluation of immunodecoration revealed a slight decrease in this terminal cisternae component. SAR of apparent 160 kd




Fig 2. Coomassie-stained gel of microsomal muscle protein fraction from control and ethanol-fed rats. Shown is a Coomassie-labeled gel, whereby lanes 1 and 2, 3 and 4, and 5 and 6 are the microsomal fraction isolated from rats representing the control-fed, ethanol-fed, and ad libitum animal groups, respectively. Muscle was obtained from rats fed on a solid diet ad libitum, or rats fed on a nutritionally complete liquid diet containing either glucose or ethanol as 35% of total dietary energy.

Alpha-1 Dihydropyridine (DHP) Receptor




Fig 3. Immunoblot: α_1 -subunit of the DHPR in muscle from control and ethanol-fed rats. Shown is an immunoblot labeled with an antibody to the α_1 -DHPR, whereby lanes 1 and 2, 3 and 4, and 5 and 6 are the microsomal fraction isolated from rats representing the control-fed, ethanol-fed, and ad libitum animal groups, respectively. The relative position of the immuno-decorated bands is marked by an arrow. The position of molecular mass standards ($\times 10^{-3}$) is indicated at the left. Histograms pertain to glucose-fed control and ethanol-fed rats and are displayed as the mean \pm SEM ($n = 5$ pairs). P values are displayed over the histograms. Detailed analysis of the muscle from rats on the solid chow diet was not carried out as explained in the main text (Methods and Discussion sections) and are displayed for comparative purposes only.

represents a Ca^{2+} -binding protein (and possibly luminal Ca^{2+} -transporter) predominantly localized to the longitudinal tubules of the sarcoplasmic reticulum and its expression was not significantly altered in microsomes from myopathic rats (Fig 6). Immunolabeling of the ubiquitous Ca^{2+} -binding protein CAL did not reveal distinct enough signals for proper statistical evaluation. However, our preliminary analysis indicates that the overall expression levels of this protein have not been changed in alcohol-exposed skeletal muscle microsomes (not shown). Since antibodies to the RyR1 isoform of the ryanodine receptor Ca^{2+} -release channel did not sufficiently cross-react with rat muscle for proper immuno decoration, we used the 90-kd junctional face membrane protein as a triad marker protein. This protein was previously shown to exist in a very close neighborhood relationship with the junctional Ca^{2+} -release channel complex and is therefore a suitable substitute for the RyR1 protein. The immunoblot analysis shown in Fig 7 demonstrates that chronic exposure to alcohol had no significant effect on this sarcoplasmic reticulum protein. The most

important Ca^{2+} -removal system during skeletal muscle relaxation is the sarcoplasmic reticulum Ca^{2+} -ATPase and immuno decoration of the fast-twitch isoform SERCA1 with monoclonal antibody IIIH11 clearly revealed an upregulation of this ion pump in myopathic specimens (Fig 8). Statistical analysis demonstrates an additional 65% increase in the relative expression level of SERCA1 ($P = .0113$). In agreement with the increased relative abundance of the SERCA1 isoform, there is an upregulation of its activity by approximately 15% ($P = .0357$; Fig 10). For internal standardization of our immunoblotting, we employed the abundant microsomal protein β -dystroglycan. As illustrated in Fig 9, normal control and myopathic membrane preparations exhibit very comparable levels of this sarcolemmal marker demonstrating equal loading in individual gel lanes. Thus, the increase in the expression of the Ca^{2+} -ATPase protein are not artefacts of gel electrophoresis procedures, protein transfers or immuno decoration methodology, but represent the effect of chronic alcohol exposure on key Ca^{2+} -regulatory membrane proteins.

DISCUSSION

Methodological Considerations

An important element of our study was the employment of a pair-feeding regimen, which facilitated the direct comparison

Alpha-2 Dihydropyridine (DHP) Receptor




Fig 4. Immunoblot: α_2 -subunit of the DHPR in muscle from control and ethanol-fed rats. Shown is an immunoblot labeled with an antibody to the α_2 -DHPR, whereby lanes 1 and 2, 3 and 4, and 5 and 6 are the microsomal fraction isolated from rats representing the control-fed, ethanol-fed, and ad libitum animal groups, respectively. For other details see legend to Fig 3.




Fig 5. Immunoblot: CSQ in muscle from control and ethanol-fed rats. Shown is an immunoblot labeled with an antibody to CSQ, whereby lanes 1 and 2, 3 and 4, and 5 and 6 are the microsomal fraction isolated from rats representing the control-fed, ethanol-fed, and ad libitum animal groups, respectively. For other details see legend to Fig 3.

between glucose-fed control and ethanol-fed treated rats. The purpose of employing a pair-feeding protocol pertains to the anorexic effects of consuming large amounts of alcohol as described previously.^{21,22} Rats fed alcohol-containing diets consume considerably less food than their counterparts fed diets without alcohol.²⁰⁻²² Thus, potentially, effects observed in alcohol-fed rats may be due to reduced dietary intakes rather than alcohol toxicity per se.²⁰⁻²² To resolve this, control rats were given identical amounts of the same liquid diet in which ethanol was replaced by isocaloric glucose.^{21,22} To confirm whether the pair feeding was necessary we analyzed muscle from a group of rats fed a solid chow diet ad libitum for the duration of the study. At the end of the 6 weeks, muscle from these rats were heavier than those from either the glucose-fed controls or ethanol-fed rats.

We are unable to explain why a study reported by another group showed no change in skeletal muscle Ca^{2+} -ATPase

activity in experimental chronic alcoholic myopathy.²⁹ Changes in enzyme activity in our model were small though statistically significant. However, consideration needs to be given to the fact that enzyme activities assayed in artificial conditions in vitro do not reflect the precise regulatory control mechanisms occurring in the intact cell in vivo. We are confident of an upregulation as much greater changes occurred after analysis of the SERCA1 protein using immunoblotting.

It is important to point out that changes in protein expression of SERCA1 and enzyme activity per se will not by themselves contribute to the myopathy as other processes are involved. Hitherto, we believed that disturbances in the rate of protein turnover were central steps in the etiology of the myopathy.³⁰⁻³³ Subsequently, we have shown that the changes are more complex, encompassing intervening steps such as protein adduct formation, altered protease activities, and increased proto-oncogene expression, free radicals, and other pathogenic factors such as membrane damage.^{1,2,34-38} Nevertheless, it is of interest to compare the magnitude of these changes in SERCA1 with




Fig 6. Immunoblot: SAR in muscle from control and ethanol-fed rats. Shown is an immunoblot labeled with an antibody to SAR, whereby lanes 1 and 2, 3 and 4, and 5 and 6 are the microsomal fraction isolated from rats representing the control-fed, ethanol-fed, and ad libitum animal groups, respectively. For other details see legend to Fig 3.




Fig 7. Immunoblot: 90-kd junctional face membrane protein in muscle from control and ethanol-fed rats. Shown is an immunoblot labeled with an antibody to JFP, whereby lanes 1 and 2, 3 and 4, and 5 and 6 are the microsomal fraction isolated from rats representing the control-fed, ethanol-fed, and ad libitum animal groups, respectively. For other details see legend to Fig 3.

other reported myopathologies (most of which show decreases in Ca^{2+} -ATPase activity or protein). Thus, in ischemia enzyme activities decrease in vitro, by about 25%.³⁹ In experimental heart failure, skeletal muscle SERCA protein (albeit the 2a isoform) has been reported to decrease by 16% in slow-twitch muscle at a significance level of $P < .001$.⁴⁰

We must also view the refractory nature of the non-SERCA proteins (Figs 3 through 7 and 9) in a more positive light. Remarkably, these proteins are conserved in both abundance and immunogenicity even in the presence of high levels of circulating ethanol and acetaldehyde (for example see Worrall et al³ and Niemala et al⁴). Previous studies have shown the presence of protein adducts, especially around the subsarcolemmal regions so a downregulation or loss of immunoreactivity of such non-SERCA proteins would be expected.

The abundance of SERCA1 is clearly increased in the alcohol-fed rats compared to the isocaloric glucose-fed rats (Fig 8). We have interpreted this observation to indicate that alcohol-treatment induced an upregulation of the sarcoplasmic reticulum Ca^{2+} -ATPase. However, it appears that the SERCA1 content in the alcohol-treated rats appears to be very similar to those rats fed ad libitum. An alternative interpretation of this data would be that alcohol treatment protected against the downregulation of the sarcoplasmic reticulum Ca^{2+} -ATPase

seen in those rats fed a restricted diet (ie, the isocaloric glucose-fed rats). However, it would be highly imprudent to make detailed biochemical comparisons with the ad libitum-fed rats with either of the 2 other groups. This is because the compositions of the solid chow and liquid diets are entirely different with respect to macro- and micronutrients, such as vitamins, minerals, and other essential and nonessential dietary elements. Differences between the rats fed the solid chow ad libitum and the animals on the liquid diets may thus be due to any one of several components. For example, dietary fat, amino acids, vitamins, and minerals and utilizable energy may all alter muscle biochemistry and composition at the gross biochemical or molecular level (for example, see previous works⁴¹⁻⁴⁶). This aspect of alcohol feeding has been the subject of a number of reviews including our own.²⁰⁻²² Thus, in comparing the 2 groups of rats fed the liquid diet, we can confidently argue that putative changes within ethanol-exposed skeletal muscle must be due to either ethanol, its metabolites or ethanol-induced metabolic disturbances per se rather than malnutrition. In simple terms, the increase in SERCA1 and Ca^{2+} -ATPase in this present study must have been due to alcohol rather than altered dietary intake.

**Sarcoplasmic-Endoplasmic Reticulum
 Ca^{2+} -ATPase (SERCA1)**




Fig 8. Immunoblot: SERCA1 Ca^{2+} -ATPase in muscle from control and ethanol-fed rats. Shown is an immunoblot labeled with an antibody to the SERCA1 isoform of the Ca^{2+} -ATPase, whereby lanes 1 and 2, 3 and 4, and 5 and 6 are the microsomal fraction isolated from rats representing the control-fed, ethanol-fed, and ad libitum animal groups, respectively. For other details see legend to Fig 3.




Fig 9. Immunoblot of β -dystroglycan in muscle from control and ethanol-fed rats. Shown is an immunoblot labeled with an antibody to β -dystroglycan, whereby lanes 1 and 2, 3 and 4, and 5 and 6 are the microsomal fraction isolated from rats representing the control-fed, ethanol-fed, and ad libitum animal groups, respectively. For other details see legend to Fig 3.

Membrane Damage in Alcoholism and Other Muscle Disorders

The toxic effects of alcohol appear to be accumulative and alcoholics often exhibit multiorgan involvement such as the heart and liver as well as skeletal muscle.⁴⁷ Perturbations in calcium homeostasis appear to be a common pathogenic mechanism. For example, reduced plasma membrane Ca^{2+} -ATPase activities are seen in hepatocyte and neutrophils in alcoholic liver disease⁴⁸ and also after carbon tetrachloride damage of liver plasma membranes.⁴⁹ These changes have been ascribed to the membrane disordering effects of ethanol.^{50,51} Alcoholic myopathology is also postulated to be triggered by alterations in membrane fluidity, ion channel activity, and ion pump regulation, thereby causing, putatively, an overall depression of muscle protein synthesis and muscle contractility.^{1,2,35}

We have recently argued that such membrane changes in alcohol myotoxicity may be related to the involvement of free

radicals.² This is also supported by the observation that the decrease in erythrocyte plasma membrane Ca^{2+} -ATPase due to alcohol exposure can be prevented by supplementation with the anti-oxidant vitamin E.⁵² However, studies investigating experimental vitamin E deficiency on skeletal muscle have shown markedly reduced SERCA activity.⁵³ Other pathologies in skeletal muscle reduce either SERCA Ca^{2+} -ATPase activity or mRNA such as ageing, muscle fatigue, and denervation.⁵⁴⁻⁵⁷ The directional changes of SERCA in these aforementioned pathologies directly contrast with the observed increase in SERCA1 due to alcohol feeding in the present study. These divergent results may be explained by the fact that the defects in alcohol-exposed muscle are distinctly complex, involving perturbations in protein synthesis, altered protein breakdown, DNA and RNA damage, and protein adduct formation.^{1,2,35} Such metabolic abnormalities all have the potential to influence steady-state levels of membrane proteins or their activities either indirectly (for example, via DNA damage) or directly (for example, via inactivation via covalent linkage with acetaldehyde).^{1,2,35} However, increases in SERCA1 protein and mRNA in type I fibers are observed in unloading,⁵⁸ which is compatible with the results obtained in muscle of alcohol-exposed rats.

In interpreting the significance of our observations, one must distinguish between acute and chronic ethanol exposure. Acutely, alcohol-induced fiber damage has been shown to trigger the leakage of the muscle marker creatine kinase, im-




Fig 10. Enzyme activity of the Ca^{2+} -ATPase in muscle from control and ethanol-fed rats. Rats were fed a nutritionally complete liquid diet containing ethanol as 35% of total calories. Controls were pair fed the same diet in which ethanol was replaced by isocaloric glucose. At the end of the study, rats were killed and the muscles dissected out. Histograms pertain to glucose-fed control and ethanol-fed rats and are displayed as the mean \pm SEM ($n = 8$ pairs). P values are displayed over the histograms. Detailed analysis of the muscle from chow fed rats was not performed as explained in the main text (Methods and Discussion sections).

plicating impairment in the membrane integrity of skeletal muscle cells.⁵⁹ However, plasma creatine kinase activities are not raised in chronic studies, possibly reflecting a component of adaptation or tolerance.^{30,60} Such a process of tolerance may be responsible for the upregulation of SERCA1, as opposed to a decrease in SERCA1 seen in most other muscle^{54-57,61} and alcohol-related pathologies.^{48,49}

Studies on human skeletal muscle biopsies from chronic alcoholics showed reduced myosin Ca²⁺-ATPase activities.⁶² However, it is important to emphasize that a distinction should be made between myofibrillary Ca²⁺-ATPase and the calcium regulatory enzymes associated with membranes.

The Contribution of Inactivity to Alcoholic Myopathy

There is a need to address the supposition that reduced physical activity in alcoholism may possibly contribute to the myopathy and muscular weakness. It is well known that reduced muscular activity leads to "disuse atrophy." However, the contribution of inactivity as a cause for the myopathy can be discounted for two reasons. First, disuse atrophy is primarily characterized by type I fiber atrophy and/or damage.⁶³⁻⁶⁶ Some studies have shown that disuse atrophy is associated with type II fiber atrophy, but these studies have not dissected out anorexia or other metabolic abnormalities which also cause type II fiber atrophy (for effects of starvation on different fibre-predominant muscles, see Preedy et al^{67,68}). Secondly, we have shown that increased activity (via swimming exercise) does not prevent the ethanol-induced type II fiber myopathy in experimental rats.⁶⁹ In the aforementioned studies, we used pair-feeding regimens for all treatment groups, which takes account of exercise-induced dietary changes.⁶⁹ This contrasts with studies using exercise regimens to prevent the myopathy without primary consideration of nutritional factors.⁷⁰

The Role of Centrally Mediated Depression

Centrally mediated depression may possibly contribute to the phenomena of alcoholic myopathy. Although there is little relevant work on the effects of centrally mediated depression on skeletal muscle calcium regulatory proteins, we can draw analogy with the depressive effects on muscle protein synthesis, which is a characteristic hallmark of alcoholic myopathy. Administration of morphine sulfate and morphine-6-glucuronide depresses skeletal muscle protein synthesis, possibly via respiratory depression.⁷¹ Currently, however, we have no evidence that such comparisons are strictly analogous to the present alcohol-feeding study due to the complexity of alcohol metabolism inducing free radicals, endocrine changes, formation of adducts, and other modes of damage.^{1,2,34,35} For example, acute morphine sulfate and morphine-6-glucuronide administration does not alter circulating insulin concentration, although other endocrine changes occur.⁷¹ In contrast, insulin levels increase in response to ethanol.³¹

Implications of the Study

Patients suffering from alcoholic myopathy may lose up to 20% of their musculature.^{1,35} Since muscle represents the most abundant mammalian tissue (40% of body weight), chronic alcoholic myopathy has profound implications for the physiology of the whole body.^{1,35} In addition, defects in muscle strength and contractility will contribute markedly to increased morbidity and impaired quality-of-life measures in affected subjects.^{1,35} It is tempting to speculate that disturbances in SERCA1 may be a contributing factor to this pathology.

ACKNOWLEDGMENT

We thank Dr Kevin P. Campbell (University of Iowa, Iowa City, IA) for supplying us with antibodies.

REFERENCES

1. Preedy VR, Paice A, Mantle D, et al: Alcoholic myopathy: Biochemical mechanisms. *Drug Alcohol Depend* 63:199-205, 2001
2. Preedy VR, Adachi J, Asano M, et al: Free radicals in alcoholic myopathy: Indices of damage and preventive studies. *Free Radic Biol Med* 32:683-687, 2002
3. Worrall S, Niemela O, Parkkila S, et al: Protein adducts in type I and type II fibre predominant muscles of the ethanol-fed rat: Preferential localisation in the sarcolemmal and sub-sarcolemmal region. *Eur J Clin Invest* 31:723-730, 2001
4. Niemela O, Parkkila S, Koll M, et al: Generation of protein adducts with malondialdehyde and acetaldehyde in muscles with predominantly type I or type II fibers in rats exposed to ethanol and the acetaldehyde dehydrogenase inhibitor cyanamide. *Am J Clin Nutr* 76:668-674, 2002
5. Paice AG: Molecular and cellular mechanisms in alcohol-induced muscle disease. PhD thesis. University of London, London, UK, 1998
6. Adachi J, Asano M, Ueno Y, et al: 7 α - and 7 β -hydroperoxycholest-5-en-3 β -ol in muscle as indices of oxidative stress: Response to ethanol dosage in rats. *Alcohol Clin Exp Res* 24:675-681, 2000
7. Nicolas JM, Antunez E, Thomas AP, et al: Ethanol acutely decreases calcium transients in cultured human myotubes. *Alcohol Clin Exp Res* 22:1086-1092, 1998
8. Melzer W, Herrmann-Frank A, Luttgau HC: The role of Ca²⁺ ions in excitation-contraction coupling of skeletal muscle fibres. *Biochim Biophys Acta* 1241:59-116, 1995
9. Murray BE, Froemming GR, Maguire PB, et al: Excitation-contraction-relaxation cycle: Role of Ca²⁺-regulatory membrane proteins in normal, stimulated and pathological skeletal muscle (review). *Int J Mol Med* 1:677-687, 1998
10. Catterall WA: Structure and function of voltage-gated ion channels. *Annu Rev Biochem* 64:493-531, 1995
11. Leong P, MacLennan DH: Complex interactions between skeletal muscle ryanodine receptor and dihydropyridine receptor proteins. *Biochem Cell Biol* 76:681-694, 1998
12. MacLennan DH: Ca²⁺ signalling and muscle disease. *Eur J Biochem* 267:5291-5297, 2000
13. Yano K, Zarain-Herzberg AL: Sarcoplasmic reticulum calsequestrins: Structural and functional properties. *Mol Cell Biochem* 135: 61-70, 1994
14. Ohkura M, Furukawa K, Fujimori H, et al: Dual regulation of the skeletal muscle ryanodine receptor by triadin and calsequestrin. *Biochemistry* 37:12987-12993, 1998
15. Leberer E, Timms BG, Campbell KP, et al: Purification, calcium binding properties, and ultrastructural localization of the 53,000- and 160,000 (sarcolumenin)-dalton glycoproteins of the sarcoplasmic reticulum. *J Biol Chem* 265:10118-10124, 1990
16. Michalak M, Mariani P, Opas M: Calreticulin, a multifunctional

Ca²⁺ binding chaperone of the endoplasmic reticulum. *Biochem Cell Biol* 76:779-785, 1998

17. Froemming GR, Ohlendieck K: The role of ion-regulatory membrane proteins of excitation-contraction coupling and relaxation in inherited muscle diseases. *Front Biosci* 6:D65-D74, 2001
18. Odermatt A, Taschner PE, Khanna VK, et al: Mutations in the gene-encoding SERCA1, the fast-twitch skeletal muscle sarcoplasmic reticulum Ca²⁺ ATPase, are associated with Brody disease. *Nat.Genet* 14:191-194, 1996
19. Ohlendieck K, Ervasti JM, Snook JB, et al: Dystrophin-glycoprotein complex is highly enriched in isolated skeletal muscle sarcolemma. *J Cell Biol* 112:135-148, 1991
20. Preedy VR, Duane P, Peters TJ: Biological effects of chronic ethanol consumption: A reappraisal of the Lieber-De Carli liquid-diet model with reference to skeletal muscle. *Alcohol Alcohol* 23:151-154, 1988
21. Preedy VR, McIntosh A, Bonner AB, et al: Ethanol dosage regimens in studies of ethanol toxicity: influence of nutrition and surgical interventions. *Addict Biol* 1:255-262, 1996
22. Lieber CS, DeCarli LM: The feeding of ethanol in liquid diets. *Alcohol Clin Exp Res* 10:550-553, 1986
23. Murray BE, Ohlendieck K: Cross-linking analysis of the ryanodine receptor and alpha1-dihydropyridine receptor in rabbit skeletal muscle triads. *Biochem J* 324:689-696, 1997
24. Froemming GR, Murray BE, Harmon S, et al: Comparative analysis of the isoform expression pattern of Ca(2+)-regulatory membrane proteins in fast-twitch, slow-twitch, cardiac, neonatal and chronic low-frequency stimulated muscle fibers. *Biochim Biophys Acta* 1466:151-168, 2000
25. Bradford MM: A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* 72:248-254, 1976
26. Culligan K, Banville N, Dowling P, et al: Drastic reduction of calsequestrin-like proteins and impaired calcium-binding in dystrophic mdx muscle. *J Appl Physiol* 92:435-445, 2002
27. Laemmli UK: Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 227:680-685, 1970
28. Towbin H, Staehelin T, Gordon J: Electrophoretic transfer of proteins from polyacrylamide gels to nitrocellulose sheets: Procedure and some applications. *Proc Natl Acad Sci USA* 76:4350-4354, 1979
29. Cardellach F, Taraschi TF, Ellingson JS, et al: Maintenance of structural and functional characteristics of skeletal-muscle mitochondria and sarcoplasmic-reticular membranes after chronic ethanol treatment. *Biochem J* 274:565-573, 1991
30. Preedy VR, Peters TJ: The effect of chronic ethanol feeding on body and plasma composition and rates of skeletal muscle protein turnover in the rat. *Alcohol Alcohol* 23:217-224, 1988
31. Preedy VR, Duane P, Peters TJ: Comparison of the acute effects of ethanol on liver and skeletal muscle protein synthesis in the rat. *Alcohol Alcohol* 23:155-162, 1988
32. Preedy VR, Peters TJ: Acute effects of ethanol on protein synthesis in different muscles and muscle protein fractions of the rat. *Clin Sci (Colch)* 74:461-466, 1988
33. Preedy VR, Peters TJ: The effect of chronic ethanol ingestion on protein metabolism in type-I- and type-II-fibre-rich skeletal muscles of the rat. *Biochem J* 254:631-639, 1988
34. Preedy VR, Seitz HK: Oxidative stress, cell damage and its possible prevention by alpha-tocopherol: Cardiovascular and other cells, in Watson RR (ed): *Alcohol and Heart Disease*. London, UK, Taylor & Francis, 2001, pp 120-145
35. Preedy VR, Peters TJ, Adachi J, et al: Pathogenic mechanisms in alcoholic myopathy, in Agarwal DP, Seitz HK (eds): *Alcohol in Health and Disease*. New York, NY, Marcel Dekker, 2003, pp 243-259
36. Preedy VR, Adachi J, Peters TJ, et al: Recent advances in the pathology of alcoholic myopathy. *Alcohol Clin Exp Res* 25:54S-59S, 2001
37. Koll M, Ahmed S, Mantle D, et al: Effect of acute and chronic alcohol treatment and their superimposition on lysosomal, cytoplasmic, and proteosomal protease activities in rat skeletal muscle in vivo. *Metabolism* 51:97-104, 2002
38. Paice AG, Hesketh JE, Towner P, et al: Alcohol increases c-myc mRNA and protein in skeletal and cardiac muscle. *Metabolism* 51:1285-1290, 2002
39. Tupling R, Green H, Senisterra G, et al: Ischemia-induced structural change in SR Ca²⁺-ATPase is associated with reduced enzyme activity in rat muscle. *Am J Physiol Regul Integr Comp Physiol* 281:R1681-R1688, 2001
40. Simonini A, Chang K, Yue P, et al: Expression of skeletal muscle sarcoplasmic reticulum calcium-ATPase is reduced in rats with postinfarction heart failure. *Heart* 81:303-307, 1999
41. Kjeldsen K, Everts ME, Clausen T: Effects of semi-starvation and potassium deficiency on the concentration of ouabain-binding sites and sodium and potassium contents in rat skeletal muscle. *Br J Nutr* 56:519-532[³H], 1986
42. Astier C, Rock E, Lab C, et al: Functional alterations in sarcoplasmic reticulum membranes of magnesium-deficient rat skeletal muscle as consequences of free radical-mediated process. *Free Radic Biol Med* 20:667-674, 1996
43. Tesseraud S, Peresson R, Lopes J, et al: Dietary lysine deficiency greatly affects muscle and liver protein turnover in growing chickens. *Br J Nutr* 75:853-865, 1996
44. Yoshizawa F, Kimball SR, Vary TC, et al: Effect of dietary protein on translation initiation in rat skeletal muscle and liver. *Am J Physiol* 275:E814-E820, 1998
45. Ohtsuka A, Kojima H, Ohtani T, et al: Vitamin E reduces glucocorticoid-induced oxidative stress in rat skeletal muscle. *J Nutr Sci Vitaminol (Tokyo)* 44:779-786, 1998
46. Samec S, Seydoux J, Dulloo AG: Post-starvation gene expression of skeletal muscle uncoupling protein 2 and uncoupling protein 3 in response to dietary fat levels and fatty acid composition: A link with insulin resistance. *Diabetes* 48:436-441, 1999
47. Estruch R, Nicolas JM, Villegas E, et al: Relationship between ethanol-related diseases and nutritional status in chronically alcoholic men. *Alcohol Alcohol* 28:543-550, 1993
48. Baffy G, Varga Z, Foris G, et al: Disturbed intracellular calcium-related processes of hepatocytes and neutrophils in human alcoholic liver disease. *Clin Biochem* 23:241-245, 1990
49. Parola M, Albano E, Autelli R, et al: Inhibition of the high affinity Ca²⁺-ATPase activity in rat liver plasma membranes following carbon tetrachloride intoxication. *Chem Biol Interact* 73:103-119, 1990
50. Kowalczyk CL, Stachecki JJ, Schultz JF, et al: Effects of alcohols on murine preimplantation development: Relationship to relative membrane disordering potency. *Alcohol Clin Exp Res* 20:566-571, 1996
51. Chen SY, Yang B, Jacobson K, et al: The membrane disordering effect of ethanol on neural crest cells in vitro and the protective role of GM1 ganglioside. *Alcohol* 13:589-595, 1996
52. Nanji AA, Sadrzadeh SM: Effect of fish oil and vitamin E on ethanol-induced changes in membrane ATPases. *Life Sci* 55:L245-L249, 1994
53. Nirdnoy W, Komararat P, Wilairat P: Comparison of sarcoplasmic reticulum Ca²⁺-adenosine triphosphatase from vitamin E-deficient dystrophic rabbit skeletal muscle with iron-ascorbate-treated and untreated enzyme. *J Biochem (Tokyo)* 103:309-312, 1988
54. Schulte L, Peters D, Taylor J, et al: Sarcoplasmic reticulum Ca²⁺ pump expression in denervated skeletal muscle. *Am J Physiol* 267:C617-C622, 1994

55. McKenna MJ, Harmer AR, Fraser SF, et al: Effects of training on potassium, calcium and hydrogen ion regulation in skeletal muscle and blood during exercise. *Acta Physiol Scand* 156:335-346, 1996
56. Viner RI, Ferrington DA, Williams TD, et al: Protein modification during biological aging: selective tyrosine nitration of the SERCA2a isoform of the sarcoplasmic reticulum Ca^{2+} -ATPase in skeletal muscle. *Biochem J* 340:657-669, 1999
57. Hunter SK, Thompson MW, Ruell PA, et al: Human skeletal sarcoplasmic reticulum Ca^{2+} uptake and muscle function with aging and strength training. *J Appl Physiol* 86:1858-1865, 1999
58. Peters DG, Mitchell-Felton H, Kandarian SC: Unloading induces transcriptional activation of the sarco(endo)plasmic reticulum Ca^{2+} -ATPase 1 gene in muscle. *Am J Physiol* 276:C1218-C1225, 1999
59. Amaladevi B, Pagala S, Pagala M, et al: Effect of alcohol and electrical stimulation on leakage of creatine kinase from isolated fast and slow muscles of rat. *Alcohol Clin Exp Res* 19:147-152, 1995
60. Martin F, Ward K, Slavin G, et al: Alcoholic skeletal myopathy, a clinical and pathological study. *Q J Med* 55:233-251, 1995
61. Loirat MJ, Lucas-Heron B, Ollivier B, et al: Calcium binding protein changes of sarcoplasmic reticulum from rat denervated skeletal muscle. *Biosci Rep* 8:369-378, 1988
62. Martin FC, Slavin G, Levi AJ, et al: Investigation of the organelle pathology of skeletal muscle in chronic alcoholism. *J Clin Pathol* 37:448-454, 1984
63. Herbison GJ, Jaweed MM, Ditunno JF: Muscle fiber atrophy after cast immobilization in the rat. *Arch Phys Med Rehabil* 59:301-305, 1978
64. Gibson JN, Halliday D, Morrison WL, et al: Decrease in human quadriceps muscle protein turnover consequent upon leg immobilization. *Clin Sci (Colch)* 72:503-509, 1987
65. Smith HK, Maxwell L, Martyn JA, et al: Nuclear DNA fragmentation and morphological alterations in adult rabbit skeletal muscle after short-term immobilization. *Cell Tissue Res* 302:235-241, 2000
66. Blakemore SJ, Rickhuss PK, Watt PW, et al: Effects of limb immobilization on cytochrome c oxidase activity and GLUT4 and GLUT5 protein expression in human skeletal muscle. *Clin Sci (Colch)* 91:591-599, 1996
67. Preedy VR, Garlick PJ: Protein synthesis in skeletal muscle of the perfused rat hemicorpus compared with rates in the intact animal. *Biochem J* 214:433-442, 1983
68. Preedy VR, Paska L, Sugden PH, et al: The effects of surgical stress and short-term fasting on protein synthesis in vivo in diverse tissues of the mature rat. *Biochem J* 250:179-188, 1988
69. Cook EB, Palmer TN, Peters TJ, et al: The imposition of moderate exercise on the development of biochemical abnormalities in alcoholic myopathy. *Clin Sci (Colch)* 86:26, 1994 (suppl 30, abstr)
70. Vila L, Ferrando A, Voces J, et al: Effect of chronic ethanol ingestion and exercise training on skeletal muscle in rat. *Drug Alcohol Depend* 64:27-33, 2001
71. Hashiguchi Y, Molina PE, Preedy VR, et al: Central effects of morphine and morphine-6-glucuronide on tissue protein synthesis. *Am J Physiol* 271:R619-R625, 1996