



Statistical modeling of rates and trends in Holocene relative sea level

Erica L. Ashe ^{a,b,c,*}, Niamh Cahill ^d, Carling Hay ^e, Nicole S. Khan ^f, Andrew Kemp ^g, Simon E. Engelhart ^h, Benjamin P. Horton ^{f,i,j}, Andrew C. Parnell ^{d,k}, Robert E. Kopp ^{b,c}



^a Department of Statistics and Biostatistics, Rutgers University, New Brunswick, NJ, United States

^b Department of Earth & Planetary Sciences, Rutgers University, New Brunswick, NJ, United States

^c Institute of Earth, Ocean & Atmospheric Sciences, Rutgers University, New Brunswick, NJ, United States

^d School of Mathematics and Statistics, Maynooth University, Kildare, Ireland

^e Department of Earth and Environmental Sciences, Boston College, Chestnut Hill, MA, United States

^f Asian School of the Environment, Nanyang Technological University, Singapore

^g Department of Earth and Ocean Sciences, Tufts University, Medford, MA, United States

^h Department of Geosciences, University of Rhode Island, Kingston, RI, United States

ⁱ Earth Observatory of Singapore, Nanyang Technological University, Singapore

^j Department of Marine and Coastal Sciences, Rutgers University, New Brunswick, NJ, United States

^k Hamilton Institute, Insight Centre for Data Analytics, Maynooth University, Kildare, Ireland

ARTICLE INFO

Article history:

Received 8 March 2018

Received in revised form

18 October 2018

Accepted 23 October 2018

Available online 5 December 2018

Keywords:

Hierarchical statistical modeling

Sea level

RSL

ABSTRACT

Characterizing the spatio-temporal variability of relative sea level (RSL) and estimating local, regional, and global RSL trends requires statistical analysis of RSL data. Formal statistical treatments, needed to account for the spatially and temporally sparse distribution of data and for geochronological and elevational uncertainties, have advanced considerably over the last decade. Time-series models have adopted more flexible and physically-informed specifications with more rigorous quantification of uncertainties. Spatio-temporal models have evolved from simple regional averaging to frameworks that more richly represent the correlation structure of RSL across space and time. More complex statistical approaches enable rigorous quantification of spatial and temporal variability, the combination of geographically disparate data, and the separation of the RSL field into various components associated with different driving processes. We review the range of statistical modeling and analysis choices used in the literature, reformulating them for ease of comparison in a common hierarchical statistical framework. The hierarchical framework separates each model into different levels, clearly partitioning measurement and inferential uncertainty from process variability. Placing models in a hierarchical framework enables us to highlight both the similarities and differences among modeling and analysis choices. We illustrate the implications of some modeling and analysis choices currently used in the literature by comparing the results of their application to common datasets within a hierarchical framework. In light of the complex patterns of spatial and temporal variability exhibited by RSL, we recommend non-parametric approaches for modeling temporal and spatio-temporal RSL.

© 2018 Elsevier Ltd. All rights reserved.

1. Introduction

The instrumental record of change in relative sea level (RSL, the difference between sea-surface height and land-surface height) is short, with the oldest tide-gauge record (Amsterdam, The Netherlands) dating to the 18th century (e.g., Van Veen, 1945;

Woodworth, 1999). Modern, quality-controlled measurements from Northern Hemisphere sites are available beginning in the early-to-mid 19th century, and globally from the mid 20th century onward. However, the geographic distribution of observations remains skewed to the Northern Hemisphere (PSMSL, 2017; Holgate et al., 2013; Pugh, 1987). RSL proxies are therefore required to infer RSL changes and the contribution of processes that operate over longer timescales (Bloom, 1964; Shennan, 1989; Törnqvist et al., 2008; Dutton et al., 2015). Whereas instrumental records are (near-)continuous with relatively small vertical uncertainty and negligible (minute-to-hour resolution) temporal uncertainties, RSL

* Corresponding author. Department of Earth & Planetary Sciences, Rutgers University, New Brunswick, NJ, United States.

E-mail address: erica.ashe@rutgers.edu (E.L. Ashe).

proxy data exhibit sample-specific vertical (inferential and measurement) and temporal uncertainties (e.g., Törnqvist et al., 2015; Woodroffe et al., 2015; Hibbert et al., 2016). Like the distribution of tide-gauge measurements, the distribution of RSL proxy data is sparse in time and space.

Quantifying the rates and spatial patterns of RSL change, on timescales ranging from decades to millennia, therefore involves piecing together sparse and noisy instrumental and/or proxy data (e.g., Kopp et al., 2009, 2016; Hay et al., 2015; Piecuch et al., 2017). Statistical models allow RSL records to be filtered for quality assurance (Düsterhus et al., 2016b) as well as fused in a consistent manner that allows rigorous quantification of multiple sources of uncertainty. Statistical models are needed to answer fundamental questions in sea-level research, such as quantifying rates of RSL change (e.g., Cahill et al., 2015a; Khan et al., 2015), assessing spatial variability of the extent and magnitude of high stands (e.g., Kopp et al., 2009; Khan et al., 2017; Vacchi et al., 2018), identifying the global-mean sea-level (GMSL) signal (e.g., Church and White, 2004; Jevrejeva et al., 2006; Kopp et al., 2009; Hay et al., 2015; Kopp et al., 2016), and improving estimates to constrain dominant physical processes, including ice-sheet behavior and glacio-isostatic adjustment (GIA; Engelhart et al., 2011; Mitrovica et al., 2011), based on their distinct spatial and temporal patterns (e.g., Milne et al., 2005; Dangendorf et al., 2017; Kopp et al., 2015a; Hay et al., 2015). Although statistical methods have for decades played a major role in reconstructing other paleoclimate variables (e.g., temperature; Visser and Molenaar, 1988; Fritts, 1991; Smith et al., 1996; Mann et al., 1998), their application to instrumental (e.g., Church and White, 2004; Jevrejeva et al., 2006; Hay et al., 2013; Kopp, 2013) and paleo (e.g., Parnell, 2005; Kopp et al., 2009; Cahill et al., 2016; Khan et al., 2015) RSL data is more recent.

Hierarchical statistical models, described in detail in Section 2, distinguish between a process level (representing, for example, the physics of RSL change) and a data level (representing, for example, the noisy recording of RSL by instruments or proxies) and cleanly distinguish between variability and uncertainties introduced at these different levels (Cressie and Wikle, 2015; Tingley et al., 2012). They are flexible, capable of accommodating missing data, and enable probabilistic inference about RSL over time and space. Viewing statistical models in a hierarchical framework, however, does not require a hierarchical computational implementation; the hierarchical perspective provides a valuable tool for dissecting and comparing models (Tingley and Huybers, 2010), regardless of implementation. Though only some authors have used hierarchical RSL models explicitly (e.g., Kopp et al., 2016; Khan et al., 2017), almost all statistical models of RSL can be reformulated hierarchically. Using a hierarchical framework to present *modeling choices* (i.e., how to characterize the relationships among variables¹) and *analysis choices* (i.e., how to implement a model structure) in a consistent manner, we present an integrated perspective on the choices made in analyzing temporal and spatio-temporal RSL datasets. Although this paper primarily concentrates on proxy data from the Holocene, the models are applicable to other timescales (e.g., Kopp et al., 2009). The appropriate modeling and analysis choices depend on the research questions asked, the type of data used, and the spatio-temporal scale (e.g., local to global, years to millennia) under consideration.

In the remainder of this paper, we first introduce hierarchical models and their application to RSL data (Section 2). We then describe different models representing the data-generating process (Section 3.1) by which RSL is linked to proxy records. At the

process level, we describe time-series models (Section 3.2) for representing RSL at a single site and spatio-temporal models (Section 3.3) for representing the temporal evolution of sea level across a regional or global domain. We then discuss different analysis techniques and their advantages and disadvantages (Section 4). To illustrate more concretely the similarities and differences between these approaches, we build case studies by applying models to common datasets including tide-gauges measurements (TGs), near-continuous RSL reconstructions (*continuous cores*), and *sea-level index points* (*SLIPs*) from locations along the Atlantic coast of the United States (Section 5). Finally, we make recommendations to help identify which methods to use to obtain temporal and spatio-temporal estimates of RSL and/or GMSL and rates of change based on the data being analyzed and the objective of the study (Section 6).

2. Hierarchical statistical modeling

In statistical nomenclature, *uncertainty* signifies an interval around which the true value is likely to fall, whereas *statistical error* is the (unknown) difference between the predicted value and the true value. *Residuals*, which can be analyzed to test assumptions about modeled errors, are the difference between an observed and a predicted value (Section 4). The *prior distribution* of a Bayesian model represents the knowledge about a given phenomenon before new data is observed, whereas the *posterior distribution* is the conditional probability that is assigned after the relevant new evidence (the observed data) is taken into account.

Hierarchical statistical models, which are frequently but not always implemented in a Bayesian framework, partition the multiple random effects that lead to individual observations into levels, thus clarifying the assumptions in a statistical analysis. They separate the underlying phenomenon of interest, such as sea level, and its variability, characterized at what is called the *process level*, from the noisy mechanism by which this underlying process is observed, characterized at the *data level*). Bayesian hierarchical models are based on conditional probabilities: observed data are regarded as conditional on a *latent* (unobserved) process, which is conditioned on unknown parameters and the assumptions in the model structure. Inverting the conditional probabilities allows probabilistic estimation of a time series or field, which can vary as a function of time and/or space. Each level of a hierarchical statistical model quantifies uncertainties separately; this can require more careful consideration of sources of uncertainties than approaches that pool the uncertainties from different levels together. Almost any statistical model can be reinterpreted as a hierarchical model; doing so increases transparency by explicitly making the distinction between modeling assumptions and analysis methods (or inference choices), as well as the difference between process variability and observation *noise*.

The primary goal in statistical analysis of RSL data is to estimate latent RSL (i.e., the noise-free time series or spatio-temporal field) and its uncertainty from observed, *noisy data*. At least three levels are defined in most RSL model hierarchies. The data level characterizes the relationship between RSL and the observed RSL data (instrumental and/or proxy) and incorporates measurement, inferential (e.g., from the conversion of a proxy's elevation to a distribution of RSL), and dating uncertainties. The process level models 'true' (i.e., noise-free) RSL and, in some cases, decomposes RSL into the underlying processes that comprise it. The parameter level captures key attributes of the data and process levels through unobserved parameters (e.g., characteristic temporal and spatial scales of variability). Hierarchical models estimate the posterior probability distribution of the noise-free RSL time series or field

¹ Italicized terms are defined in the glossary, Section 3.

(and its uncertainty), which enables probabilistic inference about RSL over time and space (see Cressie and Wikle, 2015, for further details on hierarchical models). We interpret published sea-level analyses within a hierarchical framework in order to compare modeling assumptions as well as analysis methods from these implementations.

Conditional probability distributions are the basic mechanism for modeling uncertainty in hierarchical models. The *conditional probability* distribution of A, given B, is denoted $p(A|B)$. Bayesian statistics uses *Bayes' theorem* (Laplace, 1812) to invert the conditional probability of the observed data, y , and calculate the conditional probability of unknown parameter(s) or process(es), θ , given the data, y :

$$p(\theta|y) = \frac{p(y|\theta)p(\theta)}{p(y)}. \quad (1)$$

The *likelihood* function, $p(y|\theta)$ (also known as a sampling or data distribution), is the probability of observing the data as described by the parameter(s) or process(es) of the fitted model. The prior distribution, $p(\theta)$, expresses a priori beliefs about the unknown parameter(s) or process(es), before data have been observed, and $p(y)$ is the *marginal likelihood* of the data, defined as the probability of observing y averaged across all possible processes or parameters. The conditional, posterior distribution, $p(\theta|y)$, is the resulting process or parameter distribution, given the observations. The parameters used to construct the prior distribution, known as *hyperparameters*, can be fixed, estimated, or have (*hyper*)prior distributions themselves. For the remainder of this paper, we will ignore the marginal likelihood, which is irrelevant provided the observations are static, and use the alternative form of Bayes' theorem that states the posterior is proportional to the likelihood times the prior:

$$p(\theta|y) \propto p(y|\theta)p(\theta). \quad (2)$$

In a simple hierarchical statistical model of RSL, the data model, $p(y|f, \theta_d)$, expresses the distribution of the RSL data, y , given the latent (unobserved) sea-level process, f , and the parameters of that

distribution, θ_d . Below the data level, the RSL process model, $p(f|\theta_s)$, incorporates scientific knowledge and uncertainty into the estimation of the true RSL process through its conditional parameters, θ_s . On the bottom level, the parameter model, $p(\theta_d, \theta_s)$, specifies the prior distribution of all unknown parameters and hyperparameters.

$$p(f, \theta_s, \theta_d|y) \propto \underbrace{p(y|f, \theta_d)}_{\text{posterior}} \cdot \underbrace{p(f|\theta_s)}_{\text{data model}} \cdot \underbrace{p(\theta_d, \theta_s)}_{\text{process model parameter model}} \quad (3)$$

Modeling choices refer to the relationships defined within a model and the assumptions made in constructing these relationships (e.g., a linear relationship between time and RSL), whereas analysis choices describe decisions about how to implement a specific model structure (e.g., using least-squares analysis, Aitken, 1934; likelihood maximization, Wilks, 1938; or fully Bayesian analysis with Monte Carlo sampling, Hastings, 1970).

Models are always simplified, imperfect versions of reality. It is therefore important to recognize that a model's estimate of the truth (e.g., the latent process, f) is conditional upon the assumptions of the model and the accuracy of the analysis approach. Because a perfect model of the world is also a uselessly intractable one (Borges, 1954), statistical estimates are useful, but imperfect, approximations. Consideration of alternative sets of structural modeling assumptions is an important part of characterizing the robustness of an estimate.

The hierarchical statistical framework accommodates a broad range of complexity in modeling and analysis choices, and most methods of statistical analysis used in sea-level science can be reframed as hierarchical models. For example, the structure of trends in RSL through time can be defined prior to analysis by explicitly assuming linear, polynomial, piecewise-linear, or other forms of the relationship between time and RSL at the process level. *Non-parametric* approaches, such as spline regression (e.g., Gharineiat and Deng, 2015) or models with *Gaussian process* priors (GP; Rasmussen and Williams, 2006; e.g., Kopp et al., 2009; Cahill et al., 2015a), can also be used to determine trends, without a predetermined functional form at the process level. A probabilistic

Table 1
Techniques table.

Technique	Analysis Methods	Modeling Choices	Data	Time Period	Illustrative Publications
Simple Linear Regression	Least squares	Temporally linear	TGs, CCs, SLIPs	≤ 3 ky	Shennan et al. (2002) Engelhart et al. (2009)
EIV Change-point	Errors-in-variable, Bayesian analysis	Change-point model	CCs, TGs, SLIPs	Common Era, Late Holocene	Kemp et al. (2013), Brain et al. (2015)
EIV IGP	Errors-in-variable, Bayesian analysis	Covariance functions, Proxy-systems model, Integrated GP	TGs, CCs, SLIPs	Common Era, Holocene	Gehrels and Woodworth (2013), Cahill et al. (2015a, 2015b)
Regional Averaging	Least squares, Ad hoc, Virtual station	Physical models	TGs, Altimetry data	Instrumental	Douglas (1991), Jevrejeva et al. (2009), Dangendorf et al. (2017)
EOF Regression	Least squares	EOFs	TGs, Altimetry data	Instrumental	Church and White (2006)
Probabilistic Ensembles	Particle filter	Physical models	SLIPs	LIG	Düsterhus et al. (2016)
Kalman Smoother	Multi-model KS	Spatio-temporal, Covariance functions, Physical models	TGs	Instrumental	Hay et al. (2013, 2015, 2017)
Gaussian processes	Bayesian Analysis, Empirical Bayesian analysis	Spatio-temporal, Covariance functions Physical models	TGs, CCs, SLIPs	Instrumental, Holocene, LIG, Common Era	Parnell (2005), Kopp (2013), Kopp et al. (2015), Khan et al. (2015, 2017)

Table includes common techniques, analysis methods, modeling choices, the type of data typically used, relevant time periods to which this approach has been applied, and some examples in publications. Sections 3.2 and 3.3 provide details on the modeling choices, and section 4 discusses specific analysis choices. TGs - tide gauges; CCs - continuous core records; SLIPs - sea-level index points; EIV - errors-in-variables; IGP - integrated Gaussian process; EOFs - empirical orthogonal functions.

ensemble approach (e.g., Düsterhus et al., 2016a,b), where each ensemble member is assigned an equal prior probability, is another option for modeling the process level. Table 1 presents pairs of modeling and analysis choices from past analyses, recast within a hierarchical framework, highlighting the time periods and data that have been analyzed in the literature using these models.

3. Modeling choices

3.1. Modeling the data level

RSL proxy data differ from instrumental data in their sources of uncertainty, which are modeled at the data level. Whereas instrumental data have negligible temporal uncertainty, proxy data have inherent temporal uncertainties (e.g., from radiometric dating; Polach, 1976; Stuiver and Polach, 1977; Reimer et al., 2013). In the broadest sense, there are two types of Holocene RSL proxy data. Discrete SLIPs constrain the position of RSL in time and space (geographic and vertical) and can be treated independently of one another at the data level in most circumstances (e.g., Shennan et al., 2002; Engelhart and Horton, 2012; Engelhart et al., 2015). In contrast, continuous cores are produced by analyzing a sequence of ordered samples from a single sediment core. These records constrain RSL change through time at a single geographic location (Gehrels et al., 2002; Varekamp et al., 1992), but the data points are not independent because a common age-depth model (e.g., Wright et al., 2017) is generally used to estimate sample age. A particular challenge when working with RSL proxy data is realistically characterizing the geochronological uncertainties that arise from the process of radiocarbon calibration (Reimer et al., 2013), which results in probability distributions for calendar ages that are often multi-modal and discontinuous. However, many models assume (explicitly or implicitly) normal uncertainties for calibrated radiocarbon ages for simplicity (although this is an oversimplification of reality). Some age-depth models (e.g., Parnell et al., 2008, 2015; Wright et al., 2017) handle these difficulties and return predicted age distributions with approximately normal uncertainties.

The data level of a hierarchical statistical model represents the relationship between uncertain observations and RSL. The specific type of data and associated uncertainty determine the form of this relationship and hence the form of the data-level model. For example, regression models often assume that the independent variable, time, t , has been measured exactly, and only account for uncertainty in time's functional relationship with RSL, f . This link between the observed data and the sea-level process can be represented as

$$y_i = f(t_i) + \epsilon_i, \quad (4)$$

where y_i is proxy or instrumental observation i and $f(t_i)$ is true RSL (under the assumptions of the model) at the time that y_i was observed. Many models assume measurement uncertainties are independent and normally distributed, such that $\epsilon_i \sim \mathcal{N}(0, \sigma_i^2)$, where σ_i is the assumed standard deviation of measurement and *inferential uncertainty* for observation i (e.g., Hijma et al., 2015). This assumption, however, typically ignores some biases (e.g., mis-calibration) and assumes an ideal measurement is taken, which is rarely achieved in reality. In analyses that do not incorporate measurement and inferential uncertainty specific to each observation, ϵ_i are typically assumed to be independent and identically distributed (iid) with Gaussian uncertainty and to pool data uncertainty with process variability not represented in the structure of $f(t)$. The data level of a spatio-temporal model is equivalent to that of a time-series model, where true sea-level, $f(\mathbf{x}_i, t_i)$, is dependent on both geographic location, \mathbf{x}_i , and time, t_i .

The distinction between measurement uncertainty and inferential uncertainty (the relationship between RSL and a proxy's position) can be explicit:

$$y_i = f(t_i) + \epsilon_i + \eta_i \quad (5)$$

where ϵ_i is the unobserved measurement error for observation i , and η_i is the indicative meaning (vertical relationship of a proxy to contemporary tide levels) uncertainty for the specific sample ($\eta = 0$ for direct instrumental observations of RSL), which may depend on time due to changes in tidal range. Hierarchical models for RSL proxies can be even more explicit in the representation of η_i ; whereas this is often specified in a database based upon an interpretation (which can introduce subjectivity and additional assumptions) conducted separately, it can be related directly and probabilistically to raw data, such as microfossil species abundances, in an additional level of the model (e.g., extending the approach used by Parnell et al., 2015 for paleoclimate data).

Temporal uncertainties in proxy data are separated from process noise at the data level:

$$\bar{t}_i = t_i + \delta_i, \quad (6)$$

where \bar{t}_i is the central point estimate of the calibrated age for radiocarbon dating, t_i is the true age (under the assumptions of the model), which is unknown and unobserved, and δ_i is unobserved temporal error, which is often incorporated as normal uncertainty within the analysis. These uncertainties can be incorporated in several different ways (see Section 4.6).

3.2. Process level: modeling the temporal sea-level process

RSL time-series models have a long history, beginning with hand-drawn curves (e.g., Lighty et al., 1982; Zong, 2004; Smith et al., 2011; Abdul et al., 2016) and evolving to include different forms of statistical regression (e.g., temporally linear, Shennan et al., 2002; change-point, Kemp et al., 2015b; GP, Kopp et al., 2009; EOF, Church and White, 2004). Some of these explicitly separate data uncertainty from process variability; others incorporate both data uncertainty and non-linear or high-frequency process variability into the error term, ϵ . Recasting these models in a hierarchical framework allows the separation of uncertainties of different types, providing a common basis for comparing modeling choices.

3.2.1. Temporally linear models

The simplest approach to estimating RSL and an average rate of RSL change is fitting a temporally linear model to observed data. As just two examples, Shennan and Horton (2002) and Engelhart et al. (2009) applied simple linear regression to discrete SLIPs and tide-gauge measurements to estimate the rate of RSL change during the past few thousand years, over which period the observations were qualitatively judged to be well approximated by a linear trend. In both instances, the authors performed linear regression on the midpoints of the SLIPs and did not account for inferential and measurement uncertainty (temporal and vertical). The process-level relationship is represented by

$$f(t) = mt + \beta, \quad (7)$$

where $f(t)$ is the modeled true RSL, m is the constant rate of change in RSL, and β is the intercept. The slope, m , and y -intercept parameter, β , can be estimated using many analysis methods, but are most typically analyzed using least-squares regression.

Temporally linear models are familiar to most researchers and

easy to use, and therefore provide a convenient way to find a first-order estimate (a rough approximation) of rates over time periods when they are expected to be roughly constant. However, temporally linear models can provide biased estimates of the slope parameters, due to their sensitivity to the temporal distribution of data. For example, intervals with a high concentration of data exert an undue influence on rate estimates. In addition, the linearity assumption is rigid; temporally linear models lack the ability to model any evolution in rates of RSL change. Temporally linear regression also assumes stationarity of errors (errors do not change over time or among observations). Temporally linear models are appropriate for modeling a first-order estimate, but are generally inappropriate for more in-depth analyses.

3.2.2. Change-point models

Change-point models represent a single time series as separate, continuous, temporally linear sections and are generally employed to estimate the timing of changes in trend. For example, Kemp et al. (2015a) estimated when modern rates of RSL change began in Connecticut using change-point models. Long et al. (2014) used a change-point model to analyze whether there was an acceleration in RSL change in the UK over the past 300 years. At the process level, with m change points,

$$f(t) = \begin{cases} \alpha_1 + \beta_1(t - \gamma_1), & \text{when } t < \gamma_1 \\ \alpha_{j-1} + \beta_j(t - \gamma_{j-1}), & \text{when } \gamma_{j-1} < t < \gamma_j \\ \alpha_m + \beta_{m-1}(t - \gamma_m), & \text{when } \gamma_m \leq t \end{cases} \quad (8)$$

where γ_k is the change point and α_k is the expected value of RSL at the change point (with a continuity constraint, such that $\alpha_k = \alpha_{k-1} + \beta_{k-1}(\gamma_k - \gamma_{k-1})$), and β_j is the rate of RSL change for each of the $m + 1$ segments. The parameters of change-point models can be estimated using a range of analysis approaches, including non-linear least squares and empirical Bayes (Section 4.3), but in the paleo sea-level modeling literature, these models generally follow Cahill et al. (2015b) in employing a change-point process model using a *Bayesian Hierarchical Model* (BHM; Section 4.4) within an *errors-in-variable* (EIV) framework (Section 4.6).

Change-point models attempt to address a primary goal in sea-level research, identifying accelerations in RSL and GMSL change (e.g., Church and White, 2006; Jevrejeva et al., 2008; Kopp, 2013), and they improve upon simple linear models by allowing for varying rates of RSL change and are relatively simple to implement. However, the linear constraints on each section are still fairly rigid and often do not represent the true physical behavior of RSL. When there is a clear pattern of phase changes in the data and variability around the trends is *white noise* (signal having serially uncorrelated random variation), change-point models may be appropriate for estimating the timing of these phase changes; however, they cannot estimate the magnitude of accelerations because they assume acceleration is instantaneous. Additionally, the white noise assumption, which can be tested with analyses of residuals, is frequently violated. If the model accounts for every change point in the regression lines, the assumption of iid errors can be met. Alternative, less parametric approaches, such as *Kalman Smoother* (KS) or GP models, are more flexible in representing RSL time series when the data exhibit fluctuations that cannot be adequately captured by a small number of change points; as such, they can help to answer questions about accelerations in a manner that recognizes that accelerations may occur gradually rather than abruptly. For a more complete overview of change-point models, see Ducré-Robitaille et al. (2003).

3.2.3. Gaussian process models

A Gaussian process (GP) is a generalization of the Gaussian (normal) probability distribution in continuous time (and space) (Rasmussen and Williams, 2006). To our knowledge, GP modeling was introduced into sea-level analysis by Parnell (2005) and into the paleo-sea-level literature by the Last Interglacial analysis of Kopp et al. (2009). In a GP, the relationship among any arbitrary set of points (e.g., in time, or in space and time) is a multivariate normal distribution defined by a mean vector and a covariance matrix. A temporal GP is fully defined by its mean function, $\mu(t)$, and covariance function, $K(t, t')$, where t is an input variable, which here represents time (but can be extended without loss of generality to higher dimensions, for example to include geographic location in spatial sea-level modeling; see Section 3.3). When RSL, $f(t)$, is a GP this is expressed as

$$f(t) \sim \mathcal{GP}(\mu(t), K(t, t')). \quad (9)$$

The covariance function, $K(t, t')$, defines prior expectations about the variance of the process about its mean and the correlation between points in time (and space), and thus about the way in which information is shared between time points.

In a GP model, the sea-level function, $f(t)$, is non-parametric (i.e., its form is not predetermined). Accordingly, GP time-series models have much more flexibility than temporally linear or change-point models. The shape of the curve is driven by the covariance matrix, which is estimated conditional upon the data, as opposed to a predetermined functional form. A key assumption of a GP model is that values among any set of given points assume a normal distribution.

A variant type of GP model is an Integrated Gaussian Process (IGP) model, which places a GP prior on the rate process rather than the sea-level process (see Holsclaw et al., 2013 for more details and justification). The IGP approach was introduced into the RSL literature by Cahill et al. (2015a), following the methodology of Holsclaw et al. (2013), who presented a new method of obtaining the derivative process by viewing this procedure as an inverse problem. At the process level, IGP regression models the RSL rate process, $f'(t)$, as a GP. The underlying RSL process, $f(t)$, is the integral of the rate process plus a constant intercept, α :

$$f'(t) \sim \mathcal{GP}(\mu(t), K(t, t')), \quad (10)$$

$$f(t) = \alpha + \int_0^t f'(u) du, \quad (11)$$

where t is time. For example, Cahill et al. (2015a) estimated the continuous and dynamic evolution of RSL change in North Carolina from sediment cores using IGP models.

Assuming a stationary covariance for the rates of RSL change produces a non-stationary covariance for RSL. The Bayesian approach allows regularization (introducing additional information in order to prevent over-fitting or solving an ill-fit problem), which reduces issues with identifiability (the theoretical possibility of learning the true values of a model's underlying parameters after obtaining an infinite number of observations from it). One limitation of the IGP is that the sea-level function needs to be twice differentiable; unlike a GP model of levels, this does not allow abrupt changes of rate, but instead requires that any change of rate happens through a gradual acceleration. For example, any RSL trend that is well-represented by a change-point model, which assumes instantaneous acceleration, cannot be represented well by an IGP model. Another drawback of the IGP is that it is not immediately clear how to extend it to define a derivative process in multiple dimensions (e.g., Holsclaw et al., 2013), such as for applying it to

spatial datasets, although some insights could be gleaned from spatial first-difference methods, commonly applied in the econometrics literature (e.g., Juodis, 2018).

In both GP and IGP models, the covariance functions can take a range of functional forms (Section S1). The form and parameters of the covariance function – called hyperparameters, because they set assumptions that inform the non-parametric representation of $f(t)$ – define how abruptly modeled RSL may change with temporal (or spatio-temporal) distance. Scale hyperparameters express prior beliefs about the amplitude of variability over time. Range (or characteristic length scale) hyperparameters set the distance over which the correlation between two sites or times decays toward zero (e.g., Rasmussen and Williams, 2006). Smoothness hyperparameters determine the speed of decay in the correlation in time or space (e.g., the degree of differentiability). For fixed hyperparameters, GP posterior distributions are analytically tractable (i.e., no approximation or sampling is necessary) when data uncertainty is represented as normally distributed; statistically speaking, this reflects the fact that the normal distribution is self-conjugate. Covariance functions for GP priors can be constructed by summing different terms with different characteristic scales of variability; however, linking these different terms with distinct physical processes requires incorporating process knowledge through deterministic physical models.

GP and IGP models are appropriate for many applications because of their flexibility and ability to incorporate physical knowledge regarding ranges and scales of variability through their covariance functions (Section S1). However, they do have several key disadvantages. GP models generally assume that the covariance function is stationary – e.g., that prior expectations about the relationship between RSL at 10 ka and 8 ka are the same as those between 4 ka and 2 ka. This is a rough approximation, although still more flexible than parametric approaches. IGP models generally make the same assumption about rates as opposed to levels, which is a potentially more accurate approximation.

GP models are considerably more difficult to implement than linear or change-point models, although an increasing number of tools are available to assist in their implementation. For example, Kopp (2016) makes documentation and code publicly available for implementing a spatio-temporal version of a GP model, and Cahill (2018) provides code to implement an IGP model.

These models exhibit relatively long analysis times (e.g., see Table 2 for analysis times of illustrative analyses). Statistical modeling is an iterative process of model development, model fitting, and model criticism, and slow analysis methods can be a hindrance to this process. Moreover, for some methods the computational time can scale rapidly with the number of data points. The time to invert a covariance matrix for a GP analysis scales with the cube of the number of data points, and the computational time of a model that both inverts a covariance matrix and samples temporal uncertainty (e.g., the EIV-IGP) scales with the number of data points to the fourth power. Although there are techniques to estimate the covariance matrix in order to make it more easily invertible, these models are currently not scalable to large datasets using full covariance matrices.

3.2.4. Summary of time series models

Each modeling choice has advantages and disadvantages. Temporally linear models are sensitive to the temporal distribution of data and influential data points. However, when uncertainties in the data are incorporated into the model, linear regression provides an easy, fast, and appropriate way to determine first-order rates of change in processes that are approximately constant. Change-point models assume that phases of persistent sea-level behavior are approximated by linear trends, which may not accurately represent

the underlying physics of RSL change and mask (to some degree) the continuous evolution of RSL through time (Cahill et al., 2015a). Drawbacks of these simpler approaches motivated Cahill et al. (2015a) to develop a non-parametric (EIV-IGP) methodology for estimating rates of RSL change from multiple types of proxy data at a single site (Sections 5.1 and 5.2) and Kopp et al. (2009) and Kopp (2013) to develop spatio-temporal GP models (Sections 5.2 and 5.3), which share information from nearby geographical sites to overcome the limited length of records in certain locations. However, GP models also have several drawbacks, including their less intuitive nature, complexity of implementation, longer computation time, and stationarity assumptions.

3.3. Process level: modeling the spatio-temporal sea-level process

Spatio-temporal models allow information to be shared among sites based on their proximity or physical relations, and they also enable estimation of RSL and its rates of change at sites where there are no data. In addition, spatio-temporal models support the estimation of multi-site metrics, such as change in global-mean sea level (GMSL), which is defined as the spatial average of RSL or sea-surface height (SSH) over the ocean (Gornitz et al., 1982). Most spatio-temporal models implemented in the literature are not explicitly hierarchical, but – as with time-series models – they can be recast in this way in order to facilitate comparison.

Spatio-temporal RSL models represent a continuum from purely statistical to purely physical models. At the purely statistical end of this spectrum, the priors of the process level relating RSLs from different locations to one another are based solely on their spatial and temporal proximity, and the bounds on the hyperparameters (before they are optimized) are typically based on knowledge of the variability of the processes they attempt to capture. At the purely physical end of the spectrum, a deterministic model (e.g., a GIA model) is used to estimate the RSL field; probabilistic ensembles are just one example. Intermediate formulations incorporate physical information into the construction of prior distributions.

3.3.1. RSL represented with single or multiple GP priors

The simplest and most fully statistical models place a single GP prior on RSL, as in equation (9) (with the mean and covariance functions dependent on both time and geographic location), conditioning on RSL proxy or instrumental data to yield a posterior distribution of RSL in time and space. The covariance function in this context may be spatially and temporally separable, in which case it is represented as the product of a temporal covariance function and a spatial covariance function. The former describes prior expectations about scales of change in time, the latter about scales of change in space. The analysis of a spatio-temporal GP is amenable to the same approaches as a temporal GP.

A single GP with a parametric covariance function is rarely implemented in the spatio-temporal RSL modeling literature, because a single scale of temporal variability and a single scale of spatial variability is too simple to capture physical behavior. More physical insight recognizes that RSL should have multiple spatio-temporal scales of variability, and can therefore be represented as the sum of multiple terms with GP priors. Kopp (2013) introduced this approach into the spatio-temporal RSL literature to model tide-gauge data along the east coast of the United States in order to determine whether there was an acceleration in local RSL. His process model employed nine separate terms with GP priors, combining three spatial scales of variability (global, regional, and local) with three temporal scales of variability (low, medium, and high frequency). Lower resolution RSL proxy data frequently require a simpler process level. For example, several studies (e.g., Kopp et al., 2016; Khan et al., 2017) employ models of the form:

$$f(\mathbf{x}, t) = g(t) + l(\mathbf{x}, t) + s(\mathbf{x}, t), \quad (12)$$

where $g(t)$ represents a global term that is common to all sites and could include (in a global analysis) the global-mean effects of thermal expansion and changing land ice volume; $l(\mathbf{x}, t)$ is a regional linear term, which might represent processes like GIA, ocean/atmosphere dynamics, and the static-equilibrium effects of land-ice mass changes; and $s(\mathbf{x}, t)$ might capture smaller-scale region (or local) processes, like tectonics or natural sediment compaction.

Although covariance functions can be used to capture specific physical processes that influence RSL, they have fundamental limitations. Using covariance functions alone, based on scales of variability, can fail to represent the processes intended, in contrast to using physical models. For example, many common covariance functions will miss spatial teleconnections such as those associated with sea-level fingerprints or large-scale climate modes, because they assume that correlation decays with distance. [Hay et al. \(2015\)](#) used GPs but worked around this by using physical models to estimate the covariance, as did [Kopp et al. \(2009\)](#). However, the assumption that the distribution of physically probable outcomes is normal is restrictive, so this approach is also imperfect.

3.3.2. Empirical orthogonal functions

Empirical orthogonal function (EOF) regression fits sparse observations with a set of spatial patterns (EOFs) that characterize the maximum amount of variation in a relatively dense, complementary dataset. These spatial patterns are derived through EOF decomposition (equivalent to principal component analysis [PCA]), which decomposes the dense dataset into orthogonal patterns. For example, [Church and White \(2004\)](#), [Domingues et al. \(2008\)](#), and [Ray and Douglas \(2011\)](#) used EOF decomposition to calculate the dominant spatial patterns of (high-frequency) variability in GIA-corrected SSH from altimetry observations, and applied these patterns to fit tide-gauge data and estimate longer-term GMSL change. Although EOF decomposition incorporates no direct or explicit knowledge of physical processes, many of the dominant EOFs are associated with known physical modes of variability (e.g., North Atlantic Oscillation, Atlantic Multi-decadal Oscillation, Pacific Decadal Oscillation, El Niño Southern Oscillation). The process level in EOF regression can be represented as:

$$f(\mathbf{x}, t) = g(t) + \sum_i U_i(\mathbf{x})\alpha_i(t) + GIA(\mathbf{x}, t - t_0) \quad (13)$$

Here, $g(t)$ is a global ‘mode’ that is constant over space, each U represents a leading spatial EOF, α is a time series of amplitudes of the EOFs (also known as a principal component of the associated EOF), and $GIA(\mathbf{x}, t - t_0)$ represents the GIA term (implemented through a correction from a single, selected GIA model). The solution, including the amplitudes of the leading EOFs, models the change in RSL from one time step to the next (e.g., monthly averages for [Church and White, 2004, 2011](#)).

An advantage of EOFs is that they learn about correlations from the observations, which allows for complex patterns with teleconnections. The assumption that dominant spatial patterns are constant over time, across frequencies of variability, and over the changing selection of tide gauges may lead to biases, however, because of the sensitivity of EOFs to the choice of spatial domain and time period. Additionally, features of physical modes can be mixed between EOFs, and there is no guarantee that an EOF pattern has physical meaning; instead, the patterns can represent noise (see [Calafat et al., 2014](#) for a detailed critique).

3.3.3. Incorporating physics-based models

More physical knowledge can be incorporated at the process

level by building physics-based models into the covariance structure or by using a probabilistic ensemble approach. For example, rather than optimizing hyperparameters of a covariance function based on the data, [Kopp et al. \(2009\)](#) used physical models of glacial-isostatic adjustment (GIA) to help define the prior covariance structure of a spatio-temporal GP for an analysis of GMSL and RSL change during the Last Interglacial. This approach still assumed a GP prior; more complex priors can be represented through more direct use of physics-based models or emulators thereof.

Although the analysis methods (Section 4) used in the implementations differ, the process models of [Hay et al. \(2015\)](#) and [Dangendorf et al. \(2017\)](#), used to analyze the instrumental record, are similar variants of:

$$\begin{aligned} f(\mathbf{x}, t) = & g(t) + \sum_j FP_j(\mathbf{x})M_j(t) + DSL(\mathbf{x}, t) + GIA(\mathbf{x})(t - t_0) \\ & + NL(\mathbf{x}, t) + w(\mathbf{x}, t). \end{aligned} \quad (14)$$

Here, the spatio-temporal RSL field is split into several component fields. A globally uniform term, $g(t)$, includes global thermal expansion and unmodeled sources of change. (It is not, however, representative of GMSL, as several of the other terms have non-zero global means). FP_j and M_j are the static-equilibrium fingerprint and melt, respectively, for each ice sheet/glacier source regions, indexed by j . $DSL(\mathbf{x}, t)$ is dynamic sea-level change, estimated using information from atmospheric/ocean global climate models. $GIA(\mathbf{x})$ is the local contribution to RSL from GIA, estimated using information from GIA process models, $NL(\mathbf{x}, t)$ is other non-linear signals not captured by the other, and $w(\mathbf{x}, t)$ is process noise.

An advantage to incorporating knowledge of processes through physical models is that they add potential information in the open ocean, far from tide gauge sites, whereas purely statistical models lose power away from the data. They also allow for teleconnections, rather than assuming informativeness always decreases with distance. A disadvantage is that they can be more complex to implement and may be overly rigid and rely on a small number of interpretations of physical processes. Current implementations also use discrete inputs and outputs, without comprehensively accounting for uncertainties in the parameters that determine the process or using a continuous parameter space.

Emulation of complex physical models, including 1-D and 3-D GIA models and ice-sheet models, with statistical models (e.g., Gaussian process emulators; [Kennedy et al., 2006](#); [Rougier, 2008](#)) or simplified physical models (e.g., [Urban and Keller, 2010](#)) can provide a faster, more flexible way of explicitly embedding this knowledge in a hierarchical framework. Statistical emulation reduces the processing time of these physical models, which are computationally intensive; it produces continuous output, in contrast to the discrete sea-level curves that are output for each set of discrete input parameters; and it enables probabilistic conclusions about the input parameters driving the physical models.

4. Analysis choices

Analysis methods used in the sea-level modeling literature include least-squares analysis (e.g., [Church and White, 2004](#); [Shennan and Horton, 2002](#); [Engelhart et al., 2009](#)), ad hoc approaches such as ‘virtual station’ averaging (detailed in section 4.2; e.g., [Jevrejeva et al., 2006](#); [Dangendorf et al., 2017](#)), empirical Bayesian analysis (e.g., [Kopp et al., 2009](#); [Khan et al., 2017](#); [Meltzner et al., 2017](#)), fully Bayesian analysis (e.g., [Parnell, 2005](#); [Cahill et al., 2016](#)), Kalman smoother (KS) algorithms (e.g., [Hay et al., 2015](#)), and direct and approximate methods for incorporating temporal uncertainty in proxy data into statistical models. Simple process models can be implemented with almost any analysis choice, while more complex models may require non-linear least squares or a

Bayesian approach.

4.1. Least squares

Least-squares analysis optimizes a model by minimizing the sum of squared deviations between the observed RSL and a RSL process model function (Legendre, 1805). It can be used with functions as simple as a line (i.e., simple linear regression, Section 3.2.1) or as complex as in EOF regression (Section 3.3.2). Least-squares analysis is included with most statistical software (e.g., R, MATLAB, SAS) and is easy to implement with many modeling choices. However, ordinary least-squares (OLS; Aitken, 1934) analysis does not include implementation of a data level, and therefore typically excludes explicit measurement and inferential uncertainties. It also assumes errors are independent and identically distributed.

Slightly more advanced solutions than OLS include weighted least squares (WLS) and generalized least squares (GLS). WLS addresses the problem of *heteroscedastic* (unequal) variances, and GLS additionally addresses the problem of autocorrelation among variances, both of which are common characteristics of data used in sea-level analyses. GLS estimators can be more efficient than OLS estimators (Goldberger, 1962). OLS, WLS, and GLS all require parametric linear models (though note that a linear model need not be a linear function of time). Total least squares (Golub, 1973; Golub et al., 1999) is a generalization of the least-squares approximation method and incorporates uncertainty in both the independent and dependent variable, and non-linear least squares uses optimization algorithms to maximize the fit of more complex models.

4.2. Ad hoc approaches: regional averaging, virtual stations, pre-processing

We define ‘ad hoc’ approaches as analysis methods constructed without an underlying statistical theory. Modern estimates of GMSL change apply various versions of these ad hoc approaches, including regional averaging, ‘virtual stations,’ and pre-processing to different subsets of tide gauges. The results of these techniques exhibit various GMSL curves (Fig. 1).

Regional averaging effectively removes the contributions of some processes, such as those included in the regional and local terms of purely statistical models. Definitions of the number of regions and how the averaging is implemented vary by study. Jevrejeva et al. (2006, 2009, 2016) attempted to address the spatial heterogeneity of tide gauges separated by geographic regions through a ‘virtual station’ approach, which iteratively averages rates between stations to estimate a regional average and then averages across all regions to find a global average (Fig. 1). Dangendorf et al. (2017) adopted the general idea of Jevrejeva’s ‘virtual station’ technique and weighted each regional estimate by its approximate area in relation to the entire ocean (Fig. 1).

Regardless of the model, many analyses ‘correct’ for physical processes prior to analysis (e.g., Cahill et al., 2015a; Tamisiea and Mitrovica, 2011; Church and White, 2011) by removing sites that do not meet desired criteria and by subtracting out signals from physics-based process models prior to analysis (i.e., pre-processing). For example, within regional averaging implementations, Douglas (1991, 1997) and Holgate (2007) corrected for the effects of GIA using single GIA models and screened out tide-gauge stations deemed to include a sizable tectonic contribution. Dangendorf et al. (2017) corrected each tide gauge, prior to analysis, according to the static-equilibrium fingerprints of assumed melt components, GIA, and vertical land motion, which were each estimated by physical process models. For more details on the ‘virtual station’ approach, see Section S2.1.




Fig. 1. Comparison of GMSL curves based on different subsets of tide gauges, process model choices, and analysis methods, including KS, GP, virtual stations, and EOF regression (Jevrejeva et al., 2008; Church et al., 2013; Hay et al., 2015).

4.3. Empirical Bayesian analysis

Empirical Bayesian analysis, employing *Empirical Hierarchical Models* (EHMs; see Cressie and Wikle, 2015; Gelman et al., 2013 for background), uses point estimates of the parameters based on the RSL data (e.g., Kopp et al., 2016; Hay et al., 2015). Maximum likelihood estimates (MLEs, $\hat{\theta}$) are optimal point estimates found by maximizing the likelihood $p(y|\theta)$ of the model, given fixed data. An EHM yields a posterior distribution of RSL, conditional on the data and the optimal parameters $p(f|y, \hat{\theta}_s, \hat{\theta}_d)$. Although explicit bounds are usually set on hyperparameters for MLEs, there is no explicit prior distribution on the parameters. Instead, the parameter level describes the optimization or estimation of the data and process parameters, θ_d and θ_s , respectively.

$$\underbrace{p(f|y, \hat{\theta}_s, \hat{\theta}_d)}_{\text{posterior}} \propto \underbrace{p(y|f, \hat{\theta}_d)}_{\text{likelihood}} \cdot \underbrace{p(f|\hat{\theta}_s)}_{\text{prior}}. \quad (15)$$

Almost all published implementations of RSL process models with spatio-temporal GP priors applied to RSL proxy data use empirical Bayesian analysis (e.g., Kopp et al., 2015b; Khan et al., 2017). For instrumental data, Hay et al. (2015, 2017) demonstrated an EHM with GP priors alongside the KS approach (section 4.5) to estimate GMSL, the spatio-temporal RSL field, and the components contributing to RSL globally at decadal intervals from tide gauge records. Meltzner et al. (2017) implemented an empirical GP model using coral microatoll proxy data from the mid-Holocene in Southeast Asia to estimate rates of RSL change by incorporating a periodic term to capture the 18.6-year tidal cycle.

GMSL reconstructions fusing proxy and instrumental data are possible using empirical Bayesian analysis, although they have rarely been implemented. Kopp et al. (2016) provide the only example of using both instrumental and proxy data to construct an empirical GMSL reconstruction over the past 2500 years using spatio-temporal modeling with empirical Bayesian analysis (Fig. 2).

EHMs generally require fewer computational resources than fully Bayesian techniques; however, like fully Bayesian approaches, empirical GP implementations require computation of the inverse of a full covariance matrix (over all times and space), the computational demands of which are more strenuous than a state-space model (a model that is defined by a system of first-order difference equations of state variables), which estimates a covariance matrix at each time step. For this reason, EHM analyses (and BHM analyses) do not scale to large datasets as easily as other

approaches.

4.4. Fully Bayesian analysis

Another analysis choice, fully Bayesian analysis, gives rise to Bayesian Hierarchical Models (BHMs; see Cressie and Wikle, 2015; Gelman et al., 2013 for background). A fully Bayesian analysis requires that all model parameters have prior probability distributions, allowing parameters to take on a range of probable values. These prior distributions may incorporate informative prior knowledge or may be uninformative, vague priors. Priors are typically sampled using *Markov Chain Monte Carlo* techniques (MCMC: algorithms used to approximate random samples from complex probability distributions, e.g., Gelman et al., 2011); however, for a limited set of likelihood and *conjugate prior* distributions, combined with relatively simple model structures and known hyperparameters, they can be solved analytically.

The output of a BHM is the posterior distribution, $p(f, \theta_s, \theta_d|y)$, of the sea-level process, f , and the parameters, θ_s and θ_d , given the observed data, y . This posterior is proportional to the product of the likelihood of the model, $p(y|f, \theta_d)$, the prior distribution of the model, $p(f|\theta_s)$, and the prior of the parameters, $p(\theta_d, \theta_s)$, where θ_d and θ_s are the data and sea-level process hyperparameters, respectively:

$$\underbrace{p(f, \theta_s, \theta_d|y)}_{\text{posterior}} \propto \underbrace{p(y|f, \theta_d)}_{\text{likelihood}} \cdot \underbrace{p(f|\theta_s)}_{\text{prior}} \cdot p(\theta_d, \theta_s). \quad (16)$$

As with empirical Bayesian analysis, fully Bayesian analysis can be implemented with virtually any process model (e.g., Parnell et al., 2015; Cahill et al., 2015b, 2016; Piecuch et al., 2017). In general, it is more computationally demanding than an empirical analysis but provides more thorough estimates of relative uncertainties (e.g., Piecuch et al., 2017).

4.5. Kalman smoother techniques

The Kalman smoother can combine process-based models of the drivers of sea-level change with spatially and temporally sparse observations to estimate a temporal or spatio-temporal model. Implementation of the KS is based on the Kalman filter (Kalman, 1960), a data assimilation technique that iteratively performs a least-squares analysis whenever observations are available, but in the absence of observations relies on model dynamics to compute the best estimate of the state vector. The Kalman filter method assumes that the state at time k evolves linearly from the state at

$k - 1$. The approach is similar to a Bayesian updating process, occurring one time step at a time, and is equivalent to a linear Gaussian state-space model. (For examples of non-linear, non-Gaussian state-space models, see Parnell et al., 2015; Cahill et al., 2016). The KS extends the Kalman filter so that estimates at any given point in time are informed by observations in its future as well as in its past. For example, Hay et al. (2013, 2015) used the Kalman smoother to implement a model similar to that in equation (14) and thus model GMSL, the field of RSL change, and different driving processes (see Section S2.2 for more details on this implementation).

The KS approach is flexible in terms of process models that can be represented. Because it is recursive, it is computationally faster than approaches (such as EHMs or BHMs) that require estimating all spatio-temporal points simultaneously; the KS scales linearly with the number of data points (Grewal and Andrews, 2001). It is therefore especially valuable for estimating GMSL and RSL over the instrumental period, as it enables analysis of data at a higher temporal resolution than non-recursive analysis methods with comparable modeling choices. However, the KS approach fails for low data density (which can be shown analytically; see Hay et al., 2017; Gelb et al., 1974), does not readily handle temporal uncertainty (Kalman, 1960; Visser and Molenaar, 1988), and therefore has not yet been implemented in the literature using proxy data to estimate trends over longer timescales. Moreover while multiple methods have been used to incorporate temporal uncertainties within models with GP priors, including errors-in-variable (EIV) and *noisy-input GP* (NIGP) methods (Section 4.6); the KS has not been applied to RSL data with temporal uncertainties.

4.6. Incorporation of temporal uncertainty

The temporal uncertainty of RSL proxy data has been incorporated into models in various manners. An EIV framework, which has been implemented in temporal IGP (e.g., Kemp et al., 2013; Gehrels and Woodworth, 2013; Brain et al., 2015; Cahill et al., 2015a) and change-point (e.g., Cahill et al., 2015b) models and in a spatio-temporal GP model (Kopp et al., 2009), incorporates temporal uncertainty directly through MCMC sampling of the distributions. Because of its use of MCMC sampling, the EIV framework is generally employed together with a fully Bayesian analysis approach.

An alternative approach with less computational demand is to approximate and recast temporal noise as RSL uncertainty. The NIGP method of McHutchon and Rasmussen (2011) has been implemented to do this in temporal and spatio-temporal empirical




Fig. 2. GMSL estimate with 67% and 90% credible intervals over the last ~2500 years from Kopp et al. (2016) using a model with GP priors, applied to RSL proxy data and instrumental data in an empirical Bayesian analysis.

models with GP priors (e.g., Miller et al., 2013; Kopp et al., 2015b; Khan et al., 2017). The NIGP uses the first-order Taylor-series approximation – a linear expansion about each input point – to translate errors in the independent variable, time, into equivalent errors in the dependent variable, RSL, such that temporal uncertainty is recast as sea-level uncertainty proportional to the squared gradient of the GP posterior mean.

5. Illustrative analyses

There are a number of modeling and analysis choices that can be used to evaluate a given scientific question. To illustrate the advantages and disadvantages of specific implementations in RSL modeling, we apply several models to common datasets. We focus on pairs of modeling and analysis choices that commonly occur together in the literature and organize them from simple to more complex. First, we demonstrate the differences between several time-series models – temporally linear models with ordinary least-squares analysis (e.g., Shennan and Horton, 2002; Engelhart et al., 2009), change-point and EIV-IGP models with fully Bayesian analyses (e.g., Brain et al., 2015; Cahill et al., 2015b; Cahill, 2018), and temporal models with GP priors with empirical Bayesian analysis (ET-GP) – to estimate RSL change from continuous cores over the Common Era. The quasi-linearity of RSL over this period warrants an evaluation of linear and change-point models. Next, we demonstrate a spatio-temporal GP model with empirical Bayesian analysis (EST-GP; e.g., Khan et al., 2017; Kopp et al., 2016), which characterizes spatial and temporal variability in RSL change over the Holocene using proxy data. This is the only technique currently used in the RSL literature that accommodates both temporal uncertainties and spatial correlations. Therefore, to illustrate the spatio-temporal approach, we compare the results of the EST-GP model to a site-by-site ET-GP. Last, we analyze tide-gauge data with a physically-informed KS model (e.g., Hay et al., 2013) and an EST-GP model (e.g., Hay et al., 2017), as they are capable of estimating both GMSL and the spatio-temporal fields of RSL and its rates of change with uncertainties. The type of data, time period of interest, and relevant scientific question determines which techniques are appropriate. Table 2 outlines the implementations applied in Sections 5.1, 5.2, and 5.3, and the detailed descriptions of each model can be found in Section S3.

5.1. Estimating rates of RSL change from continuous cores (Common Era)

Attempting to answer scientific questions about the timing of

RSL accelerations in relation to climatic drivers requires RSL proxy data because instrumental records are frequently too short. The near-continuous records from single cores of salt-marsh sediment (continuous cores) are well-suited to capturing the onset of modern rise because they provide a longer record than instrumental data, and they possess sufficient vertical and temporal resolution to provide a meaningful estimate of the timing of accelerations in sea-level rise. The data used in the following analyses include previously published data from continuous cores collected at two sites in New Jersey (Leeds Point and Cape May Courthouse; Kemp et al., 2013, 135 data points) and one site in North Carolina (Sand Point; Kemp et al., 2011, 109 data points), where the New Jersey sites are assumed to be independent of the North Carolina site. We applied the linear, change-point, ET-GP, and EIV-IGP models, described in Section S3.1, to the data (Section S4).

Fig. 3 shows estimated RSL and rates of RSL change for each model. The change-point, ET-GP, and EIV-IGP models more appropriately characterize the uncertainties of the data by incorporating temporal and vertical uncertainties. Conversely, the rigidity of the temporally linear model does not accommodate the underlying process(es) influencing the data. The ET-GP and Bayesian EIV-IGP models yield similar mean estimates, although the Bayesian EIV-IGP makes somewhat more precise predictions with smaller uncertainties (Fig. 3) in this particular application, which may be due to the choice of covariance functions or their hyperparameters (see Section S1 for discussion of the details of covariance functions and their parameters).

Each of the models implemented assumes that the data error terms are independent and normally distributed, while the temporally linear model further assumes that the errors are identically distributed (iid). In order to test these assumptions (see Sections 3.2 and 4 for a more thorough description of model assumptions), we present a plot of the residuals as a function of the predicted RSL values and an autocorrelation function (ACF) plot, which shows the autocorrelation in residuals as a function of lag, the number of time steps between predictions (Fig. 4). If all assumptions are met, the errors should be random, meaning there is no pattern or correlation in the residuals. The residuals of the linear regression model display a non-random, temporally-dependent pattern, indicating that the model does not fit this dataset well and the temporally linear model is inconsistent with the temporal evolution in the data. The non-linear models show less structure in the residuals, as well as smaller residuals than the linear model. Although the non-linear models show smaller and less structured residuals than the linear model, it remains valuable

Table 2
Illustrative analyses table.

Implementation	Analysis Approach	Assumptions	Illustrative Dataset	Run Time
Time series analyses				
Temporally linear regression	general least-squares	Linear signal, Gaussian uncertainties, does not distinguish measurement uncertainty and process nonlinearity	Continuous core	2 s
Change-point regression	fully Bayesian	Segment-wise linear, Gaussian uncertainties, does not distinguish measurement uncertainty and process nonlinearity	Continuous core	24 min
Empirical temporal Gaussian process model (ET-GP)	empirical Bayes	Gaussian uncertainties, Stationary covariance of sea level correctly characterizes temporal variability, Once-differentiable sea-level signal	Continuous core	60 s
Bayesian error-in-variable integrated Gaussian process model (EIV-IGP)	fully Bayesian	Gaussian uncertainties, Stationary covariance of sea-level rate (can lead to non-stationary sea level covariance) correctly characterizes temporal variability, Twice-differentiable sea-level signal	Continuous core	6.1 h
Spatio-temporal analyses				
Empirical spatio-temporal Gaussian process model (EST-GP)	empirical Bayes	Spatio-temporal covariance correctly characterizes variability	Holocene Tide gauge	1.2 h 26 min
Linear state-space model (KS)	Kalman smoother	Based on the physics-based models used; linear in prediction from state to state	Tide gauge	4 h

Run times of each implementation are reported on a standard laptop computer (2015 MacBook Pro, 2.3 GHz Intel Core i5).




Fig. 3. Common Era comparison of linear model results (column 1), change-point analysis (column 2), temporal empirical GP model results (column 3), and Bayesian EIV-IGP model results (column 4), where input data are continuous cores, and there is no spatial correlation, in New Jersey (a,b), and North Carolina (c,d). Output includes estimates of RSL (a,c) and rates of RSL change (b,d), which are each shown with mean and 95% confidence or credible intervals, where each model covers more than 95% of the data, except for the linear regression model.




Fig. 4. Linear model (a,c) and all models (b,d) residuals plotted against the predicted RSL values for each site (New Jersey: a,b, North Carolina: c,d). (e) Autocorrelation function plot of residuals for each model, where the EIV-IGP North Carolina model and all three New Jersey models violate the independence assumption, because they show significant autocorrelations (more than 5% of which are above or below the blue 95% significance line). (For interpretation of the references to colour in this figure legend, the reader is referred to the Web version of this article.)

to check whether they are consistent with the assumptions of independent errors. An ACF plot can determine whether this assumption is met or violated. For example, the ACF plot in Fig. 4 demonstrates that the analyses of the North Carolina dataset do not violate the independence assumption. Conversely, the change-point, ET-GP, and EIV-IGP analyses of the New Jersey dataset do, shown by the significant (at 5% significance level; blue lines in Fig. 4) autocorrelation in residuals. This signifies that the residuals may contain additional information that is not included in the model structure.

Some of the differences between the ET-GP and EIV-IGP models are associated with the conventional choices of covariance functions used. The squared-exponential covariance function, used in the EIV-IGP, is slightly smoother in this implementation than the Matérn covariance function (see Section S1) in the ET-GP (Fig. 3). However, the choice of time length over which the rate of RSL change is averaged in the ET-GP model affects the smoothness of (and uncertainty in) the rate curve; a linear transformation on the predicted RSLs is performed to calculate the rate curve and its uncertainty. The ET-GP method, as employed, enables more complexity for various processes through multiple separate covariance functions, summed together to create the composite RSL process, but either method could incorporate various covariance functions.

In addition to making different assumptions, these four implementations produce distinct results about the probability and timing of accelerations in RSL. A temporally linear model will never predict an acceleration in RSL because of its inherent assumption of a constant rate of RSL change, whereas a change-point model is designed to detect slight changes in rate, but assumes instantaneous acceleration. The ET-GP and EIV-IGP, conversely, produce continuous posterior distributions on rates of RSL change over time. Any inflection points in the rate curves (Fig. 3b,d) can be interpreted as changes in trend, but their significance must be evaluated. For example, the ET-GP model estimates a significant (at 5% significance level) difference in the rate of RSL change in New Jersey between 1270 and 1795. Inflections can also be observed in the rate

curve for the New Jersey record around ~170 and 570 CE, indicating changes in rate, but the differences are not significant. Alternatively, taking the derivative of the rate curve, for either the ET-GP or EIV-IGP model, would produce a probabilistic estimate of acceleration (or deceleration) over time. Hence, the non-parametric nature of the ET-GP and EIV-IGP leads to more flexible inference about the evolution of RSL.

5.2. Characterizing spatial and temporal variability in RSL change from proxy data (Holocene to present)

Attempting to answer scientific questions about the regional-scale patterns of RSL associated with different driving physical processes requires combining information from various sources and locations in a spatio-temporal model. We apply the EST-GP to proxy data to illustrate the only model in the literature that accommodates both temporal uncertainties and spatial correlations.

We compiled data from previously published studies (Engelhart and Horton, 2012; Kemp et al., 2013, 2014, 2015a, 2017a, 2017b; Khan et al., 2017) along the Atlantic coast of the United States and the circum-Caribbean (latitudes 24.95–44.68°N, longitudes 67.38–81.73°W) from 11 ka to present (Supplemental S4). We employ 450 SLIPs spanning from 8 ka to present from Engelhart and Horton (2012), 66 SLIPs from 11 ka to present from Khan et al. (2017), and 498 continuous core data points and 28 SLIPs from 3 ka to present from Kemp et al. (2013, 2014, 2015a, 2017a, 2017b).

We applied the EST-GP model to the whole spatio-temporal dataset, and also applied the ET-GP model on a site-by-site basis. The models implemented are described in Section S3.2. Other modeling and analysis choices reviewed within this manuscript (e.g., KS or EOF) have not been implemented in the literature with proxy data, and therefore are not included in this illustrative analysis.

Whereas the ET-GP model only predicts RSL and rates of RSL change at sites with data (because the model is temporal only and runs independently for each site), the EST-GP model can make predictions at any point in space and time. Fig. 5a shows




Fig. 5. Spatio-temporal Holocene model (EST-GP) results, including maps of mean estimated RSL (a), rate of RSL change (b), and standard deviation of RSL (c), averaged over 4000-year periods. (d) Stars on the map show the locations of two sites with estimated RSL curves (e,f). Comparison of predicted RSL and its rates of change are shown with 95% credible intervals at North Carolina for the EST-GP (includes spatial correlations) and ET-GP models (e). The EST-GP makes predictions at a site, Inner Delaware, where no data were used as input; data are shown for comparison purposes only (f). (g) Two models demonstrate the difference in uncertainties when the model is close to data (Outer Delaware, orange) versus far from data (Merritt Island, blue). (For interpretation of the references to colour in this figure legend, the reader is referred to the Web version of this article.)

the temporal evolution of RSL estimated by the EST-GP along the Atlantic coast of the U.S. and the geographic distribution of the data used in the model over time. Fig. 5b and c show the temporal evolution of the rate of RSL change and uncertainties (standard deviation of the posterior RSL estimate), respectively. Fig. 5d shows the locations of the sites used in the comparison between the spatio-temporal and temporal-only models (North Carolina; Fig. 5e) and the site used for model validation (Inner Delaware; Fig. 5f). Data from the Inner Delaware validation site were not incorporated into the analysis. Although the EST-GP prediction at Inner Delaware is based solely on correlations in RSL with other sites, the predictions fall very close to the omitted data (shown by red boxes in Fig. 5f). At this site, only 2 out of 28 data points (7%) fall outside of the 95% credible interval model prediction of RSL. Generally, both ET-GP and EST-GP models are excessively conservative, covering more than 95% of the data that are used as input to the models within their 95% credible intervals.

One notable difference between the ET-GP and the EST-GP is the spatial correlation within the EST-GP model. The form of the RSL curve in North Carolina (Fig. 5e) is influenced by this correlation. Whereas the ET-GP model produces higher RSL at 11 ka than 10 ka, the EST-GP uses information from the correlation

with other sites to predict increasing RSL throughout the Holocene. The EST-GP also maintains fairly constant uncertainties throughout the period of interest, whereas the ET-GP has less precision when data are sparse, due to the assumption of independence between sites. However, at times and locations farther away from the data, the uncertainty increases in the EST-GP model, as well (Fig. 5c). Predicted uncertainty in RSL is greater at sites that are far from data (e.g., Merritt Island, FL; Fig. 5g) and in the early-to mid-Holocene (e.g., Fig. 5c), whereas uncertainties decrease by up to ~80% at times and locations with precise data (e.g., Outer Delaware; Fig. 5g).

In these specific implementations, another notable difference is the process level model of the EST-GP, which has three distinct terms capturing common, regional, and local signals; the regional term for each site incorporates information primarily from other sites within about 700–750 km, based on the optimized length scale parameter for that term, while the local term incorporates information primarily from only 10–12 km distance. These terms can be separated and analyzed (Fig. 6), resulting in maps of the spatio-temporal signal for each term (Fig. 6a) and plots of each term for specific sites. The common signal (which is uniform over the entire domain) absorbs a majority of the signal (Fig. 6b), whereas the regional and local signals explain the variation




Fig. 6. (a) Maps of the regional RSL signal, averaged over 4000-year periods, where the size of the white dots are proportional to the number of data points at each location, (b) the regional and local signals for three sites along the Atlantic Coast, (c) the common signal, which applies to the entire area of study as well as the regional and local signals at the same sites in b, plotted concurrently for comparison (note: these signals are not plotted on the same scale).

between sites. The higher RSL heights along the southern coast in the early- and mid-Holocene (12 ka to 4 ka) are evident in the maps (Fig. 5) and in the regional curve for South Carolina (Fig. 6b), which is shown to demonstrate distinctive patterns in regional signals in comparison to Eastern Maine and New Jersey. The Eastern Maine site has a lower regional signal, and slight differences in these sites are represented in the local signal (Fig. 6b). The regional term picks up differences among sites associated with GIA, a dominant regional process; however, the common signal absorbs a significant portion of the GIA signal, because of the similarity over the sites in the study area. Including a physical model at the process level may provide more insight into the relative contributions of other physical processes acting over different spatial scales.

5.3. Estimating spatio-temporal RSL and GMSL from instrumental data (1900 to present)

Attempting to answer scientific questions about GMSL change in the recent past requires instrumental records. During the instrumental period, data include satellite altimetry measurements and tide gauges. These data are inherently different from proxy data, as they have negligible temporal uncertainty and smaller vertical

uncertainties, and thus allow different methods. Estimating GMSL through time and interpolating the spatio-temporal field of RSL change from instrumental records in the past are challenges well-suited to KS and GP model techniques. Both techniques are implemented here using multiple tide-gauge records obtained from the Permanent Service for Mean Sea Level (PSMSL, 2017; Holgate et al., 2013), with results shown at two sites: Atlantic City, New Jersey, (39.4°N , 74.4°W) and Wilmington, North Carolina (34.2°N , 78.0°W) (Fig. 7). The models implemented are described in Section S3.3.

Both techniques can compute posterior estimates of GMSL (Fig. 7) as well as reconstruct the spatio-temporal sea-level field, conditioned on observed data, but their implementations are very different. The KS approach (Section 4.5; described in more detail in Supplemental S2.2) steps through a forward filtering pass and a backward smoother pass for each time step, enabling computation of the covariance for a smaller subset of points and thus faster solution times (~ 45 seconds for a single KS run at the tide-gauge sites only, and ~ 4 hours for the entire multi-model (see Supplemental S2.2 for details of this model) implementation globally). Conversely, the EST-GP conditions on all observations concurrently. In Hay et al. (2015), both KS and GP implementations use output from physical process models.




Fig. 7. Comparison of KS and EST-GP predictions at two sites (a) based on annually-averaged tide gauges, which are shown in red. Both models include a spatial component and produce maps of RSL rates of change, for the KS (b) and EST-GP (c). (d) GMSL time series obtained using the KS applied to subsets of tide gauges from previous studies, with 1σ uncertainties (Hay et al., 2017). (For interpretation of the references to colour in this figure legend, the reader is referred to the Web version of this article.)

However, in the current implementation, the EST-GP model has no physical-model input and is purely statistical, based solely on the data. See Hay et al. (2015) for a more complete treatment of these two analysis approaches.

We compare estimated RSL and uncertainty for the models at two sites in Fig. 7a and the estimated rates of change in RSL and their uncertainties (standard deviations) in Fig. 7b. The spatial field computed by Hay et al. (2017) is less refined because of their modeling choice to compute the global field on a standard 5° grid. A higher-resolution field can be computed with the KS; however, this will be accompanied by longer model run time. Embedded in the KS spatial maps are dynamic sea level fields from several global climate models. When less data are available, for example earlier in time when the tide gauges are sparser, the KS predicts a much rougher sea-level time series for each location, despite the fact that

there are tide gauges at these particular sites, whereas the EST-GP has larger uncertainties when there are no tide gauges as input at a specific site (Fig. 7a).

Because of the differences in implementation of the KS and EST-GP, there are some drawbacks and advantages to each. The inversion of the full covariance matrix (over all space and time points for data and predictions), which is required for the EST-GP, makes the resolution of annual tide-gauge data difficult to handle when attempting to model these data on a global scale. As a solution, lower-resolution (e.g., decadal) averages can be used as input (Hay et al., 2015). Conversely, the KS becomes unstable during the backward smoothing pass when persistent data gaps are present in the records. The KS therefore requires a subset of tide gauges which ensures observation availability over time.

The KS model has also been tested on various subsets of tide gauges (Church and White, 2011; Holgate, 2007; Ray and Douglas, 2011), which can slightly influence the results (Fig. 7d, Hay et al., 2017). However, the results with a single analysis technique are more similar than when the data and analyses are both different. When the research question relates to estimating GMSL, fully Bayesian methods may be too computationally intensive for the datasets; however, ad hoc (Section 4.2) choices may lead to different conclusions than the KS or GP techniques, especially when analyzing different sets of instrumental data.

6. Discussion and conclusions

Hierarchical statistical modeling frameworks are conceptual tools that provide a transparent approach for separating modeling choices at the data and process level from analysis choices. Appropriate modeling and analysis choices in sea-level research depend on the type of data and the scientific question(s) being addressed. We suggest that non-parametric approaches are generally best suited for analyzing RSL and rates of RSL change.

One goal in sea-level research is identifying changes in rates of RSL and GMSL change (e.g., Church and White, 2006; Jevrejeva et al., 2008; Kopp, 2013). We recommend temporal GP and IGP models, using fully Bayesian or empirical Bayesian analyses when the dataset includes instrumental or proxy data from a single site. These types of models enable probabilistic inference about accelerations and rates of change, and within a specific study, one can test hypotheses about the timing and magnitude of changes in rates.

Identifying the physical processes that explain patterns of spatial variability in RSL is a further objective of the sea-level community. Spatio-temporal approaches are required to address this problem. To date, spatio-temporal models of proxy data have generally used covariance functions that represent different scales of variability but do not tie these scales of variability to specific physical processes. Such models are useful for addressing this objective, but require post-inference comparison to physics-based models to interpret their results in terms of physical drivers. An alternative approach, yet to be demonstrated with proxy data, is to employ models with GP priors that incorporate different processes through physics-based models (as in Hay et al., 2015). This approach could be powerful for certain processes (e.g., GIA or the elastic spatial fingerprints of ice-sheet mass changes) but in other cases (e.g., dynamic sea level change over multiple millennia) adequate physics-based models are lacking, so purely statistical process-level terms will remain necessary. For constraining global sea-level change during the instrumental period, state-space analysis approaches, such as the KS, can manage large data sets more efficiently than empirical or Bayesian approaches that require simultaneously estimating all space-time points.

Improving estimates of GIA is a related goal to explaining spatial variability because it is the dominant driver of spatial variability in RSL change over mid-to-late Holocene timescales (Peltier et al., 2015). Traditionally, this is done through an iterative, manual process, where data from specific sites are compared to different versions of physical GIA models. However, alternative approaches include using a suite or probabilistic ensemble of GIA models (e.g., Hay et al., 2015; Düsterhus et al., 2016a,b) or using a single GIA model as a mean prior estimate and fitting the mismatch with a Gaussian process (e.g., Kopp et al., 2016; Vacchi et al., 2018). Both approaches can permit further constraints on the role of spatially-variable GIA, while appropriately characterizing uncertainties, although the former approach requires that the truth be represented in the

probabilistic ensemble or suite of models.

An important area of development for statistical sea-level models is more comprehensive and accurate use of data. Most proxies do not conform to normal distributions, so techniques for incorporating non-Gaussian likelihoods, such as integrating proxy systems models into spatio-temporal models, have the potential to make use of proxies that have been too approximately interpreted. A general approach has been developed by Parnell et al. (2015), which readily applies to RSL. Although they are usually carried out prior to process modeling, integrating proxy systems models into full statistical models (e.g., Cahill et al., 2016) is a key goal for the next generation of palaeo-RSL models. Data compilation efforts (e.g., Düsterhus et al., 2016a,b) aim to standardize and synthesize RSL data, which will enhance the comparability and accessibility of information to improve both physical models and statistical reconstructions. The accuracy and consistency of all statistical models will be enhanced when databases are standardized.

A key methodological challenge is scaling spatio-temporal hierarchical modeling approaches for paleo-sea level data to large, yet still temporally noisy, datasets. Unlike Gaussian process models, whose computational complexity grows in proportion with the cube of the number of data points, the computational complexity of a Kalman smoother grows linearly. For example, conducting the EST-GP analysis on 5000 proxy data points with temporal uncertainty (on a standard laptop) would lead to a computational time for a single model iteration of about 35 days; the fully Bayesian EIV-IGP analysis on the same dataset and same computational platform would – without improvements in computational efficiency – take about 1,300 years. Adapting the Kalman smoother for temporally noisy data may provide an approach to overcoming these scaling problems and thus allow the simultaneous analysis of much larger proxy datasets. Recent work in the machine-learning literature exploring the translation between Gaussian process models and linear-Gaussian state space models (e.g., Hartikainen and Särkkä, 2010) may prove useful here. There are also several approximation and estimation techniques in the GP and machine-learning literature that have not yet been applied in a sea-level context, such as variational inference (Blei et al., 2017), which could speed up analyses and improve resolution with large datasets.

Model validation and criticism are additional areas in statistical sea-level research that can be greatly improved with standardization of both data compilation and the tools used to evaluate models. Statistical methods should be validated using cross-validation techniques, bootstrapping, or simulations to precisely determine whether a model achieves what it is designed to. Having a standard set of tools to evaluate the replicability and interrogation of structural assumptions is a clear area for growth in the sea-level community to improve the reproducibility of statistical analysis.

Acknowledgements

This work was supported by National Science Foundation grants OCE-1458904 (ELA, REK, and BPH), OCE-1458903 (SEE), OCE-1458921 (ACK), OCE-1702587 (ELA and REK), Singapore Ministry of Education Academic Research Fund Tier 2 MOE218-T2-1-030, the National Research Foundation of Singapore, and the Singapore Ministry of Education under the Research Centres of Excellence initiative (NSK and BPH), Science Foundation Ireland Career Development Award grant number 17/CDA/4695 (ACP), and is a contribution to IGCP Project 639, INQUA Project CMP1601P “HOLSEA” and PALSEA3.

Glossary

Table 3

Definitions of relevant terms.

Term	Meaning
<i>ad hoc</i>	analysis methods constructed without an underlying statistical theory
<i>analysis choices</i>	decisions in how to implement a specific model structure (e.g., least-squares, likelihood maximization, empirical Bayesian analysis, fully Bayesian analysis (MCMC), Kalman Smoother, ad hoc, EIV, NIGP)
<i>Bayesian Hierarchical Model (BHM)</i>	uses fully Bayesian analysis, which approximates complicated distributions through sampling, usually using MCMC sampling
<i>conditional probability</i>	the distribution of a random quantity, given a particular value of another (unknown) random quantity; expresses uncertainty in hierarchical models
<i>conjugate prior</i>	prior distribution that comes from the same family of distributions as the likelihood distribution, so as to enable an analytically-tractable solution for the posterior
<i>continuous core</i>	near-continuous records from a single core of salt-marsh sediment or a single coral head
<i>covariance function</i>	function defining prior beliefs about the relationship between one or more variables in a Gaussian process, as a measure of how much they change together
<i>data level</i>	hierarchical level that characterizes the relationship between the phenomenon to be modeled and the observed data (instrumental and/or proxy) and incorporates measurement, inferential, and dating uncertainty
<i>Empirical Hierarchical Model (EHM)</i>	uses empirical Bayesian analysis, which estimates parameters with point estimates, usually by maximizing their likelihoods, as opposed to a BHM, which samples the prior distributions on parameters
<i>errors-in-variable (EIV)</i>	framework that accounts for the measurement uncertainty in the independent variables by assuming that uncertainties in both variables are independent of one another
<i>errors-in-variables integrated Gaussian process (EIV-IGP)</i>	model implemented on time series data, modeling the rate of RSL change, deriving the RSL curve from the rate curve; incorporates uncertainty in the independent (time) variable and the dependent variable through EIV framework
<i>empirical orthogonal function (EOF)</i>	regression technique used to find the dominant spatial patterns in a dataset; when analyzing sea level, used to find the dominant patterns in sea surface height (SSH) from satellite altimetry measurements and apply to tide gauges in order to estimate GMSL change
<i>error</i>	the difference between a measurement and the true value, for a particular data point; one can model the error as a random draw from an uncertainty distribution
<i>empirical spatio-temporal Gaussian process (EST-GP)</i>	model with Gaussian process priors, which incorporates spatial and temporal covariance functions to produce the fields of RSL and rate of RSL change as maps; solved using an empirical methodology that maximizes the likelihood of the model conditional upon the parameters of the prior
<i>empirical temporal Gaussian process (ET-GP)</i>	model using Gaussian process priors, which is independent in space (no spatial component) and solved using an empirical methodology that maximizes the likelihood of the model conditional upon the parameters of the prior
<i>Gaussian process (GP)</i>	a generalization of the multi-variate Gaussian distribution to continuous time (and space), which is fully defined by its mean function and covariance function; GP regression provides an analytically-tractable solution when adopting the assumption of normality for all distributions
<i>hyperparameter</i>	parameter of a prior distribution
<i>hyperprior</i>	prior distribution on a hyperparameter
<i>inferential uncertainty</i>	the quantified dispersion that arises from the data-generation process from true RSL to the creation of a RSL proxy
<i>Kalman Smoother (KS)</i>	iterative method that comprises a forward filtering pass and a backward smoother pass; used in a multi-model implementation to compute posterior estimates of GMSL and spatio-temporal RSL fields, conditioning prior estimates from physical models of several processes on observations
<i>latent</i>	unobserved or hidden (e.g., the true values of RSL)
<i>likelihood</i>	the probability of observing the data as described by the fitted model; also known as the sampling or data distribution; a conditional distribution that is a function of unknown parameters for observed data and incorporates the form of uncertainty in the data (e.g., measurement and/or inferential)
<i>marginal distribution</i>	unconditional probability distribution of a random quantity, found by integrating over all values of the conditional distribution in Bayesian analyses
<i>Markov Chain Monte Carlo (MCMC)</i>	techniques used to generate random variables, perform complicated calculations, and simulate complicated distributions through sampling in Bayesian hierarchical models (common algorithms include Gibbs sampling, Metropolis-Hastings, Metropolis within Gibbs, importance sampling)
<i>modeling choices</i>	decisions that define the relationships in a model, usually at the process level; in sea-level analysis, the relationship between time, space and RSL (e.g., linear, polynomial, change-point, GP (integrated), incorporation of physical models)
<i>noisy-input Gaussian process (NIGP)</i>	a method for incorporating uncertainty in the independent variable within a Gaussian process model; using a Taylor expansion about each input point to recast input noise as output noise proportional to the squared gradient of the GP posterior mean (McHutchon and Rasmussen, 2011); in sea-level analysis, geochronological uncertainty is recast as proportional uncertainty in RSL
<i>noise</i>	error; statistical noise refers to unexplained variation or randomness
<i>noisy data</i>	error-prone data that have been corrupted by known or unknown processes
<i>non-parametric</i>	not involving any assumptions as to the functional form

Table 3 (continued)

Term	Meaning
<i>posterior distribution</i>	the probability distribution of an unknown quantity, conditional on observed data; in sea-level analysis, estimates (for example) the true RSL time series or field of RSL with uncertainties, given proxy or instrumental data
<i>prior distribution</i>	the information about an uncertain parameter or process that is combined with the probability distribution of new data to yield the posterior distribution; can be subjective, based on a priori beliefs, or uninformative, which minimizes the impact on inference
<i>process level</i>	hierarchical level at which the phenomenon of interest is modeled and in some cases, decomposed; includes process variability
<i>residuals</i>	the difference between an observed and a modeled or predicted value
<i>sea-level index point (SLIP)</i>	discrete proxy data that constrain RSL at a single point in time and space
<i>uncertainty</i>	parameter characterizing the range of values within which a measured value can be said to lie with a specified probability
<i>white noise</i>	serially uncorrelated random variation (zero mean and finite variance)

Appendix A. Supplementary information

Code for select models herein can be found at <https://doi.org/10.5281/zenodo.1643438>. Supplementary information to this article can be found online at <https://doi.org/10.1016/j.quascirev.2018.10.032>.

References

- Abdul, N.A., Mortlock, R.A., Wright, J.D., Fairbanks, R.G., 2016. Younger Dryas sea level and meltwater pulse 1B recorded in Barbados reef crest coral *Acropora palmata*. *Paleoceanography* 31, 330–344. <https://doi.org/10.1002/2015PA002847>.
- Atkien, A.C., 1934. On least-squares and linear combinations of observations. In: *Proceedings of the Royal Society of Edinburgh*, 55, pp. 43–48.
- Blei, D.M., Kucukbir, A., McAuliffe, J.D., 2017. Variational inference: a review for statisticians. *J. Am. Stat. Assoc.* 112, 859–877.
- Bloom, A.L., 1964. Peat accumulation and compaction in a Connecticut coastal marsh. *J. Sediment. Res.* 34, 599–603.
- Borges, J.L., 1954. On Exactitude in Science.
- Brain, M.J., Kemp, A.C., Horton, B.P., Culver, S.J., Parnell, A.C., Cahill, N., 2015. Quantifying the contribution of sediment compaction to late Holocene salt-marsh sea-level reconstructions, North Carolina, USA. *Quat. Res.* 83, 41–51.
- Cahill, N., 2018. IGP (Version 1.0) [Computer Software] Retrieved. <http://github.com/ncahill89/IGP>. (Accessed 25 February 2018).
- Cahill, N., Kemp, A.C., Horton, B.P., Parnell, A.C., 2016. A Bayesian hierarchical model for reconstructing relative sea level: from raw data to rates of change. *Clim. Past* 12, 525–542. <https://doi.org/10.5194/cp-12-525-2016>.
- Cahill, N., Kemp, A.C., Horton, B.P., Parnell, A.C., et al., 2015a. Modeling sea-level change using errors-in-variables integrated Gaussian processes. *Ann. Appl. Stat.* 9, 547–571.
- Cahill, N., Rahmstorf, S., Parnell, A.C., 2015b. Change points of global temperature. *Environ. Res. Lett.* 10, 084002.
- Calafat, F.M., Chambers, D.P., Tsimplis, M.N., 2014. On the ability of global sea level reconstructions to determine trends and variability. *J. Geophys. Res.: Oceans* 119, 1572–1592.
- Church, J.A., Monselesan, D., Gregory, J.M., Marzeion, B., 2013. Evaluating the ability of process based models to project sea-level change. *Environ. Res. Lett.* 8, 014051. <https://doi.org/10.1088/1748-9326/8/1/014051>.
- Church, J.A., White, N.J., 2004. Estimates of the regional distribution of sea level rise over the 1950–2000 period. *J. Clim.* 17, 2609–2625.
- Church, J.A., White, N.J., 2006. A 20th century acceleration in global sea-level rise. *Geophys. Res. Lett.* 33 <https://doi.org/10.1029/2005GL024826>.
- Church, J.A., White, N.J., 2011. sea-level rise from the late 19th to the early 21st century. *Surv. Geophys.* 32, 585–602. <https://doi.org/10.1007/s10712-011-9119-1>.
- Cressie, N., Wikle, C.K., 2015. *Statistics for Spatio-temporal Data*. John Wiley & Sons, Ltd Sons.
- Dangendorf, S., Marcos, M., Wöppelmann, G., Conrad, C.P., Frederikse, T., Riva, R., 2017. Reassessment of 20th century global mean sea level rise. *Proc. Natl. Acad. Sci. Unit. States Am.* 114, 5946–5951.
- Domingues, C.M., Church, J.A., White, N.J., Gleckler, P.J., Wijffels, S.E., Barker, P.M., Dunn, J.R., 2008. Improved estimates of upper-ocean warming and multi-decadal sea-level rise. *Nature* 453. <https://doi.org/10.1038/nature07080>.
- Douglas, B.C., 1991. global sea-level rise. *J. Geophys. Res. Oceans* 96, 6981–6992. <https://doi.org/10.1029/91jc00064>.
- Douglas, B.C., 1997. Global sea rise: a redetermination. *Surv. Geophys.* 18, 279–292. <https://doi.org/10.1023/A:1006544227856>.
- Ducré-Robitaille, J.F., Vincent, L.A., Boulet, G., 2003. Comparison of techniques for detection of discontinuities in temperature series. *Int. J. Climatol.* 23, 1087–1101. <https://doi.org/10.1002/joc.924>.
- Düsterhus, A., Rovere, A., Carlson, A.E., Horton, B.P., Kleemann, V., Tarasov, L., Barlow, N.L.M., Bradwell, T., Clark, J., Dutton, A., Gehrels, W.R., Hibbert, F.D., Hijma, M.P., Khan, N., Kopp, R.E., Sivan, D., Törnqvist, T.E., 2016a. Palaeo-sea-level and palaeo-ice-sheet databases: problems, strategies, and perspectives. *Clim. Past* 12, 911–921. <https://doi.org/10.5194/cp-12-911-2016>.
- Düsterhus, A., Tamisiea, M.E., Jevrejeva, S., 2016b. Estimating the sea level highstand during the last interglacial: a probabilistic massive ensemble approach. *Geophys. J. Int.* 206, 900–920.
- Dutton, A., Carlson, A.E., Long, A.J., Milne, G.A., Clark, P.U., DeConto, R., Horton, B.P., Rahmstorf, S., Raymo, M.E., 2015. Sea-level rise due to polar ice-sheet mass loss during past warm periods. *Science* 349. <https://doi.org/10.1126/science.aaa4019>.
- Engelhart, S., Peltier, W., Horton, B., 2011. Holocene relative sea-level changes and glacial isostatic adjustment of the U.S. Atlantic coast. *Geology* 39, 751–754. <https://doi.org/10.1130/g31857.1>.
- Engelhart, S.E., Horton, B.P., 2012. Holocene sea level database for the Atlantic coast of the United States. *Quat. Sci. Rev.* 54, 12–25. <https://doi.org/10.1016/j.quascirev.2011.09.013>.
- Engelhart, S.E., Horton, B.P., Douglas, B.C., Peltier, W.R., Törnqvist, T.E., 2009. Spatial variability of late Holocene and 20th century sea-level rise along the Atlantic coast of the United States. *Geology* 37, 1115–1118.
- Engelhart, S.E., Vacchi, M., Horton, B.P., Nelson, A.R., Kopp, R.E., 2015. A sea-level database for the Pacific coast of central North America. *Quat. Sci. Rev.* 113, 78–92. <https://doi.org/10.1016/j.quascirev.2014.12.001>.
- Fritts, H., 1991. *Reconstructing Large-scale Climatic Patterns from Tree-ring Data*. The University of Arizona Press, Tucson.
- Gehrels, W.R., Belknap, D.F., Black, S., Newnham, R.M., 2002. Rapid sea-level rise in the Gulf of Maine, USA, since AD 1800. *Holocene* 12, 383–389.
- Gehrels, W.R., Woodworth, P.L., 2013. When did modern rates of sea-level rise start? *Global Planet. Change* 100, 263–277. <https://doi.org/10.1016/j.gloplacha.2012.10.020>.
- Garber, A., Kasper, J., Nash, R., Price, C.F., Sutherland, A., 1974. *Applied Optimal Interpolation*. M.I.T. Press, Cambridge.
- Gelman, A., Brooks, S., Jones, G.L. (Eds.), 2011. *Handbook of Markov Chain Monte Carlo*. Chapman & Hall/CRC Press.
- Gelman, A., Carlin, J.B., Stern, H.S., Dunson, D.B., Vehtari, A., Rubin, D.B., 2013. *Bayesian Data Analysis*, third ed. Chapman & Hall/CRC Press, London.
- Gharneiat, Z., Deng, X., 2015. Application of the multi-adaptive regression splines to integrate sea level data from altimetry and tide gauges for monitoring extreme sea level events. *Mar. Geodes.* 38.
- Goldberger, A.S., 1962. Best linear unbiased prediction in the generalized linear regression model. *J. Am. Stat. Assoc.* 57, 369–375.
- Golub, G., 1973. Some modified matrix eigenvalue problems. *SIAM Rev.* 15, 318–344.
- Golub, G., Hansen, J., O'Leary, D., 1999. Tikhonov regularization and total least squares. *SIAM J. Matrix Anal. Appl.* 21, 185–194.
- Gornitz, V., Lebedeff, S., Hansen, J., 1982. global sea level trend in the past century. *Science* 215, 1611–1614.
- Grewal, M.S., Andrews, A.P., 2001. *Kalman Filtering: Theory and Practice Using MATLAB*, second ed. John Wiley & Sons, Inc.
- Hartikainen, J., Särkkä, S., 2010. Kalman filtering and smoothing solutions to temporal Gaussian process regression models. In: *IEEE International Workshop on Machine Learning for Signal Processing (MLSP)*.
- Hastings, W., 1970. Monte Carlo sampling methods using Markov chains and their application. *Biometrika* 57, 97–190.
- Hay, C.C., Morrow, E., Kopp, R.E., Mitrovica, J.X., 2013. Estimating the sources of global sea level rise with data assimilation techniques. *Proc. Natl. Acad. Sci. U.S.A.* 110, 3692–3699.
- Hay, C.C., Morrow, E., Kopp, R.E., Mitrovica, J.X., 2015. Probabilistic reanalysis of twentieth-century sea-level rise. *Nature* 517, 481–484.
- Hay, C.C., Morrow, E., Kopp, R.E., Mitrovica, J.X., 2017. On the robustness of Bayesian fingerprinting estimates of global sea-level change. *J. Clim.* 30, 3025–3038.
- Hibbert, F.D., Rohling, E.J., Dutton, A., Williams, F.H., Chutcharavan, P.M., Zhao, C., Tamisiea, M.E., 2016. Coral indicators of past sea-level change: a global

- repository of U-series dated benchmarks. *Quat. Sci. Rev.* 145, 1–56. <https://doi.org/10.1016/j.quascirev.2016.04.019>.
- Hijma, M., Engelhart, S.E., Törnqvist, T.E., Horton, B.P., Hu, P., Hill, D.F., 2015. Handbook of Sea-level Research. John Wiley & Sons, Ltd., p. 34. <https://doi.org/10.1002/9781118452547.ch34> chapter A protocol for a geological sea-level database.
- Holgate, S.J., 2007. On the decadal rates of sea level change during the twentieth century. *Geophys. Res. Lett.* 34. <https://doi.org/10.1029/2006GL028492>.
- Holgate, S.J., Matthews, A., Woodworth, P.L., Rickards, L.J., Tamisiea, M.E., Bradshaw, E., Foden, P.R., Gordon, K.M., Jevrejeva, S., Pugh, J., 2013. New data systems and products at the permanent Service for mean sea level. *J. Coast Res.* 29, 493–504.
- Holsclaw, T., Sansó, B., Lee, H.K.H., Heitmann, K., Habib, S., Higdon, D., Alam, U., 2013. Gaussian process modeling of derivative curves. *Technometrics* 55, 57–67. <https://doi.org/10.1080/00401706.2012.722391>.
- Jevrejeva, S., Grinsted, A., Moore, J.C., 2009. Anthropogenic forcing dominates sea level rise since 1850. *Geophys. Res. Lett.* 36. <https://doi.org/10.1029/2009GL040216>.
- Jevrejeva, S., Grinsted, A., Moore, J.C., Holgate, S.J., 2006. Nonlinear trends and multiyear cycles in sea level records. *J. Geophys. Res. Oceans* 111.
- Jevrejeva, S., Jackson, L.P., Riva, R.E.M., Grinsted, A., Moore, J.C., 2016. Coastal sea level rise with warming above 2 °C. *Proc. Natl. Acad. Sci. Unit. States Am.* 113, 13342–13347.
- Jevrejeva, S., Moore, J.C., Grinsted, A., Woodworth, P.L., 2008. Recent global sea level acceleration started over 200 years ago? *Geophys. Res. Lett.* 35. <https://doi.org/10.1029/2008GL033611>.
- Judis, A., 2018. First difference transformation in panel var models: robustness, estimation, and inference. *Econom. Rev.* 37, 650–693. <https://doi.org/10.1080/07474938.2016.1139559>.
- Kalman, R., 1960. A new approach to linear filtering and prediction problems. In: *Transactions of the ASME – Journal of Basic Engineering*, 82, pp. 35–45.
- Kemp, A., Hawkes, A., Donnelly, J., Vane, C.H., Horton, B.P., Hill, T., Anisfeld, S., Parnell, A., Cahill, N., 2015a. Relative sea-level change in Connecticut (USA) during the last 2200 years. *Earth Planet Sci. Lett.* 428, 217–229.
- Kemp, A.C., Bernhardt, C.E., Horton, B.P., Kopp, R.E., Vane, C.H., Peltier, W.R., Hawkes, A.D., Donnelly, J.P., Parnell, A.C., Cahill, N., 2014. Late Holocene sea-and land-level change on the US southeastern Atlantic coast. *Mar. Geol.* 357, 90–100.
- Kemp, A.C., Dutton, A., Raymo, M.E., 2015b. Paleo constraints on future sea-level rise. In: *Current Climate Change Reports*, pp. 205–215.
- Kemp, A.C., Hill, T.D., Vane, C.H., Cahill, N., Orton, P., Talke, S.A., Parnell, A., Sanborn, K., Hartig, E.K., 2017a. Relative sea-level trends in New York City during the past 1500 years. *Holocene* 27, 1169–1186.
- Kemp, A.C., Horton, B., Donnelly, J.P., Mann, M.E., Vermeer, M., Rahmstorf, S., 2011. Climate related sea-level variations over the past two millennia. *Proc. Natl. Acad. Sci. Unit. States Am.* 108, 11017–11022.
- Kemp, A.C., Horton, B.P., Vane, C.H., Corbett, D.R., Bernhardt, C., Engelhart, S.E., Anisfeld, S.C., Parnell, A.C., Cahill, N., 2013. Sea-level change during the last 2500 years in New Jersey, USA. *Quat. Sci. Rev.* 81, 90–104.
- Kemp, A.C., Kegel, J.J., Culver, S.J., Barber, D.C., Mallinson, D.J., Leorri, E., Bernhardt, C.E., Cahill, N., Riggs, S.R., Woodson, A.L., Mulligan, R.P., Horton, B.P., 2017b. Extended late Holocene relative sea-level histories for North Carolina, USA. *Quat. Sci. Rev.* 160, 13–30.
- Kennedy, M.C., Anderson, C.W., Conti, S., O'Hagan, A., 2006. Case studies in Gaussian process modelling of computer codes. *Reliab. Eng. Syst. Saf.* 91, 1301–1309. <https://doi.org/10.1016/j.ress.2005.11.028>.
- Khan, N.S., Ashe, E.L., Horton, B.P., Dutton, A., Kopp, R.E., Brocard, G., Engelhart, S.E., Hill, D.F., Peltier, W., Vane, C.H., Scatena, F.N., 2017. Drivers of Holocene sea-level change in the caribbean. *Quat. Sci. Rev.* 155, 13–36.
- Khan, N.S., Ashe, E.L., Shaw, T.A., Vacchi, M., Walker, J., Peltier, W., Kopp, R.E., Horton, B.P., 2015. Holocene relative sea-level changes from near-, intermediate-, and far-field locations. In: *Current Climate Change Reports*, 1, pp. 247–262.
- Kopp, R.E., 2013. Does the mid-Atlantic United States sea level acceleration hot spot reflect ocean dynamic variability? *Geophys. Res. Lett.* 40, 3981–3985. <https://doi.org/10.1002/grl.50781>.
- Kopp, R.E., 2016. CESL-STEHM-GP (Version 1.0) [Computer Software] Retrieved. <http://github.com/bobkopp/CESL-STEHM-GP>. (Accessed 25 February 2018).
- Kopp, R.E., Hay, C.C., Little, C.M., Mitrovica, J.X., 2015a. Geographic variability of sea-level change. In: *Current Climate Change Reports*, 1, pp. 192–204.
- Kopp, R.E., Horton, B.P., Kemp, A.C., Tebaldi, C., 2015b. Past and future sea-level rise along the coast of North Carolina, United States. *Climatic Change* 132, 693–707.
- Kopp, R.E., Kemp, A.C., Bittermann, K., Horton, B.P., Donnelly, J.P., Gehrels, W.R., Hay, C.C., Mitrovica, J.X., Morrow, E.D., Rahmstorf, S., 2016. Temperature-driven global sea-level variability in the Common Era. *Proc. Natl. Acad. Sci. Unit. States Am.* 113, E1434–E1441. <https://doi.org/10.1073/pnas.1517056113>.
- Kopp, R.E., Simons, F.J., Mitrovica, J.X., Maloof, A.C., Oppenheimer, M., 2009. Probabilistic assessment of sea level during the last interglacial stage. *Nature* 462, 863–867.
- Laplace, P.S., 1812. Mémoire sur les formules qui sont fonctions de tres grands nombres et sur leurs applications aux probabilités. *Mémoires de l'Academie des sciences de Paris*.
- Legendre, A.M., 1805. *On the Method of Least Squares*.
- Lighty, R.G., Macintyre, I.G., Stuckenrath, R., 1982. Acropora palmata reef framework: a reliable indicator of sea level in the western atlantic for the past 10,000 years. *Coral Reefs* 1, 125–130. <https://doi.org/10.1007/BF00301694>.
- Long, A., Barlow, N., Gehrels, W., Saher, M., Woodworth, P., Scaife, R., Brain, M., Cahill, N., 2014. Contrasting records of sea-level change in the eastern and western North Atlantic during the last 300 years. *Earth Planet Sci. Lett.* 388, 110–122.
- Mann, M.E., Bradley, R.S., Hughes, M.K., 1998. Global-scale temperature patterns and climate forcing over the past six centuries. *Nature* 392, 779–787.
- McHutchon, A., Rasmussen, C.E., 2011. Gaussian process training with input noise. In: Shawe-Taylor, J., Zemel, R.S., Bartlett, P.L., Pereira, F., Weinberger, K.Q. (Eds.), *Advances in Neural Information Processing Systems*, vol. 24. Curran Associates, Inc., pp. 1341–1349.
- Meltzner, A.J., Switzer, A.D., Horton, B.P., Ashe, E.L., Qiu, Q., Hill, D.F., Bradley, S.L., Kopp, R.E., Hill, E.M., Majewski, J.M., Natawidjaja, D.H., Suwargadi, B.W., 2017. Half-metre sea-level fluctuations on centennial timescales from mid-Holocene corals of Southeast Asia. *Nat. Commun.* 8.
- Miller, K.G., Kopp, R.E., Horton, B.P., Browning, J.V., Kemp, A.C., 2013. A geological perspective on sea-level rise and its impacts along the U.S. mid-Atlantic coast. In: *Earth's Future*, 1, pp. 3–18.
- Milne, G.A., Long, A.J., Bassett, S.E., 2005. Modelling Holocene relative sea-level observations from the caribbean and South America. *Quat. Sci. Rev.* 24, 1183–1202. <https://doi.org/10.1016/j.quascirev.2004.10.005>.
- Mitrovica, J.X., Gomez, N., Morrow, E., Hay, C., Latychev, K., Tamisiea, M.E., 2011. On the robustness of predictions of sea level fingerprints. *Geophys. J. Int.* 187, 729–742. <https://doi.org/10.1111/j.1365-246X.2011.05090.x>.
- Parnell, A., Haslett, J., Allen, J., Buck, C., Huntley, B., 2008. A flexible approach to assessing synchrony of past events using Bayesian reconstructions of sedimentation history. *Quat. Sci. Rev.* 27, 1872–1885.
- Parnell, A.C., 2005. *The Statistical Analysis of Former Sea Level*. Ph.D. thesis. University of Sheffield.
- Parnell, A.C., Doan, T.K., Salter-Townshend, M., Allen, J.R.M., Huntley, B., Haslett, J., 2015. Bayesian inference for palaeoclimate with time uncertainty and stochastic volatility. In: *Journal of the Royal Statistical Society. Series C (Applied Statistics)*, 64.
- Peltier, W.R., Argus, D.F., Drummond, R., 2015. Space geodesy constrains ice age terminal deglaciation: the global ICE 6G C (VM5a) model. *J. Geophys. Res.: Solid Earth* 120, 450–487.
- Piecuch, C.G., Huybers, P., Tingley, M.P., 2017. Comparison of full and empirical Bayes approaches for inferring sea-level changes from tide-gauge data. *J. Geophys. Res.: Oceans* 122, 2243–2258.
- Polach, H. (Ed.), 1976. Radiocarbon Dating as a Research Tool in Archaeology - Hopes and Limitations, *Symposium on Scientific Methods of Research in the Study of Ancient Chinese Bronzes and South East Asian Metal and Other Archaeological Artifacts*. Australian National University.
- PSMSL, 2017. Tide gauge data. <http://www.psmsl.org/data/obtaining/>.
- Pugh, D., 1987. *Tides, Surges and Mean Sea-level: a Handbook for Engineers and Scientists*. Wiley, Chichester.
- Rasmussen, C.E., Williams, C.K.I., 2006. *Gaussian Processes for Machine Learning*. MIT Press.
- Ray, R.D., Douglas, B.C., 2011. Experiments in reconstructing twentieth-century sea levels. *Prog. Oceanogr.* 91, 496–515. <https://doi.org/10.1016/j.pocean.2011.07.021>.
- Reimer, P., Bard, E., Bayliss, A., Beck, J., Blackwell, P., Ramsey, C., Buck, C., Cheng, H., Edwards, R., Friedrich, M., Grootes, P., Guilderson, T., Haflidason, H., Hajdas, I., Hatté, C., Heaton, T., Hoffmann, D., Hogg, A., Hughen, K., Kaiser, K., Kromer, B., Manning, S., Niu, M., Reimer, R., Richards, D., Scott, E., Southon, J., Staff, R., Turney, C., van der Plicht, J., 2013. IntCal13 and Marine13 radiocarbon age calibration curves 0–50,000 years yrs BP. *Radiocarbon* 55, 1869–1887.
- Rougier, J., 2008. Efficient emulators for multivariate deterministic functions. *J. Comput. Graph. Stat.* 17, 827–843. <https://doi.org/10.1198/106186008X384032>.
- Shennan, I., 1989. Holocene crustal movements and sea-level changes in great britain. *J. Quat. Sci.* 4, 77–89. <https://doi.org/10.1002/jqs.3390040109>.
- Shennan, I., Horton, B., 2002. Holocene land-and sea-level changes in Great Britain. *J. Quat. Sci.* 17, 511–526.
- Shennan, I., Peltier, W.R., Drummond, R., Horton, B., 2002. Global to local scale parameters determining relative sea-level changes and the post-glacial isostatic adjustment of Great Britain. *Quat. Sci. Rev.* 21, 397–408.
- Smith, D., Harrison, S., Firth, C., Jordan, J., 2011. The early Holocene sea level rise. *Quat. Sci. Rev.* 30, 1846–1860.
- Smith, T.M., Reynolds, R.W., Livezey, R.E., Stokes, D.C., 1996. Reconstruction of historical sea surface temperatures using empirical orthogonal functions. *J. Clim.* 9, 1403–1420.
- Stuiver, M., Polach, H., 1977. Reporting of ¹⁴C data - discussion. *Radiocarbon* 19, 355–363.
- Tamisiea, M.E., Mitrovica, J.X., 2011. The moving boundaries of sea level change: understanding the origins of geographic variability. *Oceanography* 24, 24–39.
- Tingley, M.P., Craigmile, P.F., Haran, M., Li, B., Mannshardt, E., Rajaratnam, B., 2012. Piecing together the past: statistical insights into paleoclimatic reconstructions. *Quat. Sci. Rev.* 35, 1–22.
- Tingley, M.P., Huybers, P., 2010. A bayesian algorithm for reconstructing climate anomalies in space and time. Part I: development and applications to paleoclimate reconstruction problems. *J. Clim.* 23, 2759–2781.
- Törnqvist, T.E., Rosenheim, B.E., Hu, P., Fernandez, A.B., 2015. Handbook of Sea-level Research. John Wiley & Sons, Ltd., pp. 347–360. <https://doi.org/10.1002/9781118452547> chapter Dating Methods: Radiocarbon dating and calibration.
- Törnqvist, T.E., Wallace, D.J., Storms, J.E.A., Wallinga, J., van Dam, R.L., Blaauw, M., Derkzen, M.S., Klerks, C.J.W., Meijneken, C., Snijders, E.M.A., 2008. Mississippi

- Delta subsidence primarily caused by compaction of Holocene strata. *Nat. Geosci.* 1, 173–176.
- Urban, N.M., Keller, K., 2010. Probabilistic hindcasts and projections of the coupled climate, carbon cycle and atlantic meridional overturning circulation system: a bayesian fusion of century-scale observations with a simple model. *Tellus Dyn. Meteorol. Oceanogr.* 62, 737–750. <https://doi.org/10.1111/j.1600-0870.2010.00471.x>.
- Vacchi, M., Engelhart, S.E., Nikitina, D., Ashe, E.L., Peltier, W.R., Roy, K., Kopp, R.E., Horton, B.P., 1992. Postglacial relative sea-level histories along the Northeastern Canadian coastline. *Quat. Sci. Rev.*
- Van Veen, J., 1945. Bestaat er een geologische bodemdaling te Amsterdam sedert 1700?. In: *Tijdschrift Koninklijk Nederlandsch Aardrijkskundig Genootschap*, 1.
- Varekamp, J., Thomas, E., Van de Plassche, O., 1992. Relative sea-level rise and climate change over the last 1500 years. *Terra. Nova* 4, 293–304.
- Visser, H., Molenaar, J., 1988. Kalman filter analysis in dendroclimatology. *Biometrics* 44, 929–940.
- Wilks, S.S., 1938. The large-sample distribution of the likelihood ratio for testing composite hypotheses. *Ann. Math. Stat.* 9, 60–62.
- Woodroffe, S.A., Long, A.J., Punwong, P., Selby, K., Bryant, C.L., Marchant, R., 2015. Radiocarbon dating of mangrove sediments to constrain Holocene relative sea-level change on Zanzibar in the southwest Indian Ocean. *Holocene* 25, 820–831. <https://doi.org/10.1177/0959683615571422>.
- Woodworth, P.L., 1999. High waters at Liverpool since 1768: the UK's longest sea level record. *Geophys. Res. Lett.* 26, 1589–1592.
- Wright, A.J., Edwards, R.J., van de Plassche, O., Blaauw, M., Parnell, A.C., van der Borg, K., de Jong, A.F.M., Roe, H.M., Selby, K., Black, S., 2017. Reconstructing the accumulation history of a saltmarsh sediment core: which age-depth model is best? *Quat. Geochronol.* 39, 35–67.
- Zong, Y., 2004. Mid-holocene sea-level highstand along the Southeast coast of China. *Quat. Int.* 117, 55–67. [https://doi.org/10.1016/S1040-6182\(03\)00116-2](https://doi.org/10.1016/S1040-6182(03)00116-2).