



## A systematic exploration of a perinatal wellbeing framework through women's experiences of lumbo-pelvic pain

Franziska Wadephul<sup>1,\*</sup>, Lesley Glover, Julie Jomeen<sup>2</sup>, Nicola Hanefeld

Faculty of Health Sciences, University of Hull, Hull, United Kingdom



### ARTICLE INFO

#### Keywords:

Maternal wellbeing  
Perinatal wellbeing  
Back pain  
Pelvic girdle pain  
Systematic review

### ABSTRACT

**Background:** Women's wellbeing during the perinatal period has received increasing attention in research, policy and practice, but is often poorly defined and conceptualised. We have developed a framework of perinatal wellbeing (PWB) which we will refine further in this review, using the example of lumbo-pelvic pain (LPP). Perinatal LPP, which includes lower back pain (LBP) and pelvic girdle pain (PGP), is common and can significantly affect women's wellbeing.

**Aim:** The aims of this review are (1) to synthesise research into women's experiences of LPP and (2) to use these findings to contribute further to developing our framework of PWB.

**Designs and methods:** A systematic search of online databases was conducted for qualitative studies exploring women's experiences of LPP linked to the perinatal period; 15 papers describing 11 studies were identified. A framework synthesis approach (Carroll et al., 2011; Carroll et al., 2013) was used to synthesise studies, using the PWB framework as the *a priori* framework.

**Findings:** The review highlights the impact of LPP on all areas of women's lives and their functioning at every level, as well as the impact of a range of factors on women's experiences. Only one study explored women's experiences of LBP, all others focused on PGP. Findings illustrate how multi-faceted women's wellbeing is in the context of LPP, particularly the importance of relationships and support, but also the role played by wider socio-cultural discourses of pregnancy and motherhood and by women's individual circumstances and characteristics. Findings underline the interconnectedness of physical, emotional and psychological experiences. The review largely confirmed, and further elaborated, the domains of the original framework, but also led to some changes, notably the inclusion of an 'individual factors' domain describing women's individual circumstances and characteristics. The limited discussion of LPP during labour and birth was notable.

**Conclusions and implications:** Findings support the framework, but also provide evidence for some changes, thus further refining the framework. Women's wellbeing in the perinatal period (with regards to LPP, other issues, or generally) should not be considered in isolation, but needs to take account of women's life context. The perinatal period should be considered a continuum, rather than seeing each part in isolation. For clinical practice, the review underlines the importance of distinguishing between PGP and LBP and offering appropriate, individualised support.

### Introduction and background

There has been an increasing acknowledgement that the wellbeing of women during the perinatal period is important, not just for women themselves, but also for their babies and families, both in the short- and long-term. Consequently, perinatal wellbeing (PWB) has become an important concept within research, policy, and clinical practice. How-

ever, PWB is frequently poorly defined and often relates to only physical and/or psychological wellbeing, rather than taking a more comprehensive, multidimensional approach. If we want to be able to explore, assess, or support PWB we need to be clear about what exactly we mean by it (Ayers and Olander, 2013). In a review of theoretical discussions of PWB ((Wadephul et al., 2020)), we proposed a tentative conceptual framework of perinatal wellbeing (Fig. 1) consisting of domains pertain-

\* Corresponding author.

E-mail address: [franziska.wadephul@mu.ie](mailto:franziska.wadephul@mu.ie) (F. Wadephul).

<sup>1</sup> Current post: Department of Psychology, University of Maynooth, Ireland.

<sup>2</sup> Current post: School of Health and Human Sciences, Southern Cross University, Australia.




Fig. 1. PRISMA 2009 Flow Diagram.

ing to society and culture, community, immediate environment, and individual factors, and encompassing physical, emotional, and cognitive experiences of wellbeing. As this framework was largely based on theoretical discussion, we consider it important to explore how well it fits with women’s lived experiences. This review aims to do this by examining women’s experiences of a specific condition, lumbo-pelvic pain, within the context of the original framework. As LPP is a relatively common problem in the perinatal period and can have a serious and often long-lasting impact on women’s wellbeing, it is a pertinent example to use to further explore the original PWB framework.

Lumbo-pelvic pain (LPP) includes lower back pain (LBP) and pelvic girdle pain (PGP). Women in the perinatal period are particularly vulnerable to LPP due to hormonal and physiological changes, problems arising from birth, and the often demanding conditions of new motherhood. About 50% of pregnant women (Vleeming et al., 2008; Wu et al., 2004) and 25% of postpartum women experience postpartum LPP (Wu et al., 2004). Globally, non-specific lower back pain is the leading cause of disability (Buchbinder et al., 2013). It is a complex condition defined by the location of pain; in the majority of cases no specific cause can be identified (Hartvigsen et al., 2018).

LBP usually refers to pain between the twelfth rib and the gluteal fold (Vleeming et al., 2008); intensity varies considerably, but can have a considerable impact on women’s wellbeing and functioning. PGP is characterised by often severe pain between the posterior iliac crest and the gluteal fold, particularly around the sacroiliac joint (Vleeming et al., 2008; Wu et al., 2004). Women have described PGP as an intense, often stabbing pain (Wu et al., 2004). PGP can severely limit mobility and may necessitate the use of crutches or a wheelchair (Gutke et al., 2018). Even moderate PGP has a negative impact on women’s quality of life and daily functioning, and plays a large role in sick leave. PGP often resolves with birth, but continues postnatally in approximately a third of women (Gutke et al., 2011).

Two recent papers have reviewed women’s experiences with pregnancy-related pelvic girdle pain (Mackenzie et al., 2018; Varley and Hunter, 2019). The primary aim of this review is not to synthesise women’s experiences of LPP as such, but to explore the utility of the framework in this context. Applying the PWB framework to a specific condition experienced by many women in the perinatal period allowed us to ‘test’ the framework’s fit with women’s experiences. It also enabled elaboration of the domains and subdomains, which were relatively ab-

**Table 1**  
Search terms.

back pain OR low back pain OR lumbar pain OR lumbar pelvic pain OR lumbo pain OR symphysis pubis dysfunction OR pelvic girdle pain AND perinatal OR postnatal OR prenatal OR antenatal OR postpartum OR maternal OR pregnant OR pregnancy OR labour OR birth AND qualitative
--

stract in the original framework. In this review, through a synthesis of women's experiences of LPP, we aim:

- to evaluate how well the initial framework fits with women's lived experience of lumbo-pelvic pain,
- to refine the initial framework and add detail to the domains, and
- to explore how domains and sub-domains are related.

## Methods

### Search and selection of papers

Online databases (Academic Search Premier, CINAHL, Medline, PsycINFO) were searched using the search terms in Table 1. Reference lists were searched for additional papers. Fig. 2 shows the number of papers retrieved and retained at each stage. Papers were included if they used a qualitative methodology and focused on women's experiences of lower back pain or pelvic girdle pain arising during the perinatal period. Papers which focused predominantly on women's experiences of treatment for LPP, such as osteopathy or physiotherapy, were excluded. The search was carried out in December 2019. It was repeated in September 2019; no new papers were identified.

### Quality assessments

The quality of included papers was assessed using the CASP checklist for qualitative research (Critical Appraisal Skills Programme, 2017). This was carried out by the first author, with additional checks by another author (LG). Ratings were broadly similar between the two assessors; any divergences were discussed and mutually agreed. No papers were excluded; quality ratings are available in supplementary files.

### Synthesis

The synthesis of the included papers was based on framework synthesis (Carroll et al., 2011; Carroll et al., 2013), a deductive approach which uses an *a priori* framework against which to map and code extracted data. This allows the existing framework to act 'as the basis for the synthesis and could be built-upon, expanded upon, reduced or added to by these new data' (Carroll et al., 2011, p. 4). Data extracts from the results sections of included papers were coded, using the domains and sub-domains of the original PWB framework as the *a priori* framework. Themes, and where necessary sub-themes, were developed for each domain or sub-domain. Data which did not fit into the existing framework was coded into additional themes. Coding and the initial development of themes was carried out in NVivo by two of the authors (FW and LG).

After completion of data mapping and coding, the fit of the initial framework with the coded data from the review studies was examined to establish to what extent the initial framework was supported by the data. Changes were then made to the original framework, leading to the development of the revised framework.

## Findings

### Included studies

The search identified 15 papers, describing 11 studies, which met the inclusion criteria (Table 2). Studies originated in Norway, Sweden,

Ireland, and the UK. Even though the search terms included lower back pain and lumbo-pelvic pain in general, all studies, with one exception (Close et al., 2016), focused exclusively on the experiences of women with PGP. Close et al studied the experiences of women with pregnancy-related LPP, including GPG, LBP, and combined pain.

Almost all used interviews to gather qualitative data; one (Close et al., 2016) used focus groups and Fredriksen et al. (2008) analysed women's contributions to an online discussion forum. Data were analysed using a range of qualitative approaches (Table 2). While some studies included only pregnant women (Clarkson and Adams, 2018; Elden et al., 2013, 2014; Persson et al., 2013) or only postnatal women (Elkins-Bushnell and Boyle, 2019; Engeset et al., 2014; Gutke et al., 2017; Shepherd, 2005; Wuytack et al., 2015a, 2015b), two included both (Close et al., 2016; Fredriksen et al., 2008). In a longitudinal study (Crichton and Wellock, 2008; Wellock and Crichton, 2007a, 2007b) women were interviewed in pregnancy and postnatally. Shepherd (2005) interviewed postnatal women at two time points. For postnatal participants, the length of time at interview was not always specified. Crichton and Wellock interviewed women soon after birth (6 weeks); several other studies also included women with more persistent postnatal LPP, up to, or more than, ten years (Elkins-Bushnell and Boyle, 2019; Engeset et al., 2014; Gutke et al., 2017). One study (Wuytack et al., 2015a, 2015b) only included primiparous women. The other studies either did not specify or included both primiparous and multiparous women.

### Themes within the original framework

Table 3 shows how the themes and sub-themes were mapped with the domains and sub-domains of the original framework. Examples illustrating each theme and sub-theme are provided in the supplementary files. These themes and sub-themes further elaborate many of the domains and sub-domains which were less well defined and more abstract in the original framework.

There was some explicit evidence for the *society and culture* domain in the included papers. Support for this domain, however, was largely implicit and was commonly found in other domains, particularly other people's and women's own attitudes to LPP, pregnancy, and motherhood, for example in terms of what is considered normal. Discourses of normality in the perinatal period, particularly when used by health care professionals, work colleagues or those close to them, significantly affected women's psychological and emotional experiences and availability of care.

In the *community* domain, which in the original framework was concerned with areas of women's lives reaching beyond immediate family and friends, the review identified three elements: work or study, health professionals, and other people, including strangers and other women with LPP.

Within the *immediate environment* domain, the review identified relationships with partners, the baby and older children, other family members, and friends as relevant for women's wellbeing. These relationships could be both supportive and unsupportive, and their nature often changed due to LPP. The findings of this review highlight that the *individual* domain in the original framework had not been sufficiently clearly defined. The review identified themes relating to women's daily lives, but considerable changes have been made to this domain as a result of this review (see below).

**Table 2**  
Overview of included studies

Aims	Population	Methods	Key findings
<p><b>Clarkson &amp; Adams 2018</b> (UK) Exploring views and experiences of women with pregnancy-related PGP</p>	8 pregnant women diagnosed with PGP	Interviews (at 21 to 30 weeks gestation) Interpretive thematic data analytic approach Focus on experiences and impact of PGP, management of pain, views on treatment	Themes: <ul style="list-style-type: none"> <li>• Reality of PGP pain</li> <li>• Key mechanisms of support</li> <li>• Impact of knowledge</li> </ul>
<p><b>Close et al 2016</b> (UK) Exploring experiences of women with pregnancy-related LPP</p>	14 women: 12 postnatal (6 weeks to 9 months), 2 pregnant (36 weeks gestation) Self-selecting from participants who had taken part in reflexology RCT 8 with combination of lower back pain and pelvic pain, 5 LBP only, 1 pelvic pain only	Focus groups (3) Thematic analysis, guided by Newell and Burnard <sup>1</sup> framework Focus on: women's experiences and management of LPP	Themes: <ul style="list-style-type: none"> <li>• Physical and emotional impact on women's lives</li> <li>• Women's attitudes to, and knowledge of, LPP</li> <li>• Women's use of treatments and dissatisfaction with standard advice and treatment</li> </ul>
<p><b>Crichton &amp; Wellock 2008</b> (UK) Exploring the impact of PGP on women's lives and relationships during pregnancy and the first six weeks after giving birth (same study as <a href="#">Wellock &amp; Crichton 2007a,b</a>)</p>	28 women with PGP (diagnosed by physiotherapist) Pregnancy until 6 weeks postnatal Primi- and multi-gravida	Interviews (at initial diagnosis, 36 weeks gestation and 6 weeks after birth; not all women took part in 3 interviews) Heidggerian phenomenological approach Focus on impact on women's roles	Themes: <ul style="list-style-type: none"> <li>• Effects on personal role</li> <li>• Effects on maternal role</li> <li>• Effects on sexual relationship role</li> <li>• Effects on housekeeping role</li> </ul>
<p><b>Wellock &amp; Crichton 2007a</b> (UK) To explore women's experiences of PGP in terms of pain, impact on quality of life, and treatment by health professionals (same study as <a href="#">Crichton &amp; Wellock 2008</a> and <a href="#">Wellock &amp; Crichton 2007b</a>)</p>	28 women diagnosed with PGP (recruited in pregnancy)	Interviews at diagnosis, 36 weeks gestation and 6 weeks after birth (not all women gave 3 interviews) Phenomenological approach, analysis using Colaizzi's framework <sup>2</sup> Focus on women's experiences	Themes: <ul style="list-style-type: none"> <li>• Perceptions of pain</li> <li>• Coping with and management of PGP</li> <li>• Living with PGP</li> </ul>
<p><b>Wellock &amp; Crichton 2007b</b> (UK) Explore experiences with health professionals of women with PGP (in pregnancy and soon after birth) (same study as <a href="#">Crichton &amp; Wellock 2008</a> and <a href="#">Wellock &amp; Crichton 2007a</a>)</p>	28 women diagnosed with PGP (recruited in pregnancy)	Interviews at diagnosis, 36 weeks gestation and 6 weeks after birth (not all women gave 3 interviews) Phenomenological approach, analysis using Colaizzi's framework <sup>2</sup> Focus on women's experiences	Themes: <ul style="list-style-type: none"> <li>• Interaction with midwives</li> <li>• Interaction with doctors</li> <li>• Interaction with physiotherapists</li> <li>• Subthemes: <ul style="list-style-type: none"> <li>• Pain</li> <li>• Negative labelling</li> <li>• Dismissive staff</li> <li>• Feelings of dissatisfaction</li> </ul> </li> </ul>
<p><b>Elden et al 2013</b> (Sweden) Describing pregnant women's experiences of PGP in daily life (same study as <a href="#">Elden et al 2014</a>)</p>	27 pregnant women with PGP Recruited from participants in craniosacral RCT, all had received craniosacral therapy	Interviews Qualitative content analysis Focus on: PGP in daily life	Categories: <ul style="list-style-type: none"> <li>• PGP affects ability to cope with everyday life</li> <li>• Coping with motherhood</li> <li>• Relationships between partners often reached breaking point</li> <li>• Questioning one's identity as defined by profession/work</li> <li>• Lessons from living with PGP</li> </ul>

(continued on next page)

Table 2 (continued)

Aims	Population	Methods	Key findings
<p><b>Elden et al 2014</b> (Sweden) Exploring and describing pregnant women's experiences of severe PGP physically and relating to the healthcare system (same study as Elden et al 2013)</p>	<p>27 pregnant women with PGP All had craniosacral therapy (part of RCT)</p>	<p>Interviews Qualitative content analysis Focus on: experiences of PGP during pregnancy</p>	<p>Categories:</p> <ul style="list-style-type: none"> <li>• A strange body</li> <li>• The body on guard</li> <li>• Relation and support from health care</li> <li>• Acceptance of PGP</li> </ul>
<p><b>Elkins-Bushnell &amp; Boyle 2019</b> (UK) Exploring the occupational difficulties experienced by women with postnatal PGP and how they participate in activity</p>	<p>5 women with postnatal PGP With diagnosis of PGP or receiving treatment for PGP Between 1 and 9 years after last birth</p>	<p>Interviews Hermeneutic theory, thematic analysis Focus on: women's viewpoints, their everyday experiences</p>	<p>Themes:</p> <ul style="list-style-type: none"> <li>• Activity affected by PGP</li> <li>• Factors restricting participation in activity</li> <li>• Factors promoting participation in activity</li> <li>• Emotional impact of a change in participation</li> </ul>
<p><b>Engeset et al 2014</b> (Norway) Exploring how postnatal PGP influences women's daily life</p>	<p>5 women diagnosed with postnatal PGP Between 4 months and 11 years after last birth</p>	<p>Interviews Phenomenological-hermeneutical design</p>	<p>Themes:</p> <ul style="list-style-type: none"> <li>• Activity and pain</li> <li>• Lack of acknowledgement of pain and disability</li> <li>• Changed roles</li> </ul>
<p><b>Fredriksen et al 2008</b> (Norway) Exploring women's perspectives on PGP in pregnancy</p>	<p>Women with experience of PGP in pregnancy (contributions to online forum) Number of participants not known; data collected over one year</p>	<p>Women's contributions to an online discussion forum Qualitative text analysis (symbolic interactionist perspective) Focus: perspectives on PGP</p>	<p>Themes:</p> <ul style="list-style-type: none"> <li>• New bodily sensations</li> <li>• Fear</li> <li>• How much to endure?</li> <li>• Lack of acknowledgement</li> </ul>
<p><b>Gutke et al 2017</b> (Sweden) Exploring women's experiences of living with long-term pregnancy-related PGP</p>	<p>9 women with persistent pregnancy-related PGP Between 2 and 13 years after birth Self-reported LBP and clinical evaluation of pregnancy-related PGP</p>	<p>Interviews Analysed using empirical phenomenological psychological method Focus on: experiences of living with PGP</p>	<p>Two typologies:</p> <ul style="list-style-type: none"> <li>• The ongoing struggle against the pain</li> <li>• Adaptation and acceptance</li> <li>• Constituents:</li> <li>• Importance of the body for identity</li> <li>• Understanding of pain</li> <li>• Stages of change</li> </ul>
<p><b>Persson et al 2013</b> (Sweden) Investigating experiences of women living with PGP during pregnancy</p>	<p>9 pregnant women with diagnosed PGP Primi- and multipara; all on sick leave Last trimester</p>	<p>Interviews Analysed using a grounded theory approach Focus: experiences of PGP in current pregnancy</p>	<p>Core category: Struggling with daily life and enduring pain</p> <ul style="list-style-type: none"> <li>• actions caused by PGP: <ul style="list-style-type: none"> <li>○ grasping the incomprehensible</li> <li>○ balancing support and dependence</li> <li>○ managing the losses</li> </ul> </li> <li>• consequences of PGP: <ul style="list-style-type: none"> <li>○ enduring pain</li> <li>○ being a burden</li> <li>○ calculating the risks</li> <li>○ abdicating as a mother</li> </ul> </li> <li>• consequences regarding pregnancy / future pregnancy: <ul style="list-style-type: none"> <li>○ paying the price and reconsidering the future</li> </ul> </li> </ul>

(continued on next page)

Table 2 (continued)

Aims	Population	Methods	Key findings
<p><b>Shepherd 2005</b> (UK)</p> <p>Describe experiences of women with PGP in pregnancy and first 3 months after birth</p>	<p>9 women with PGP, recruited in third trimester</p> <p>Primi- and multi-gravida</p>	<p>Interviews 1 and 3 months after birth</p> <p>Heideggerian phenomenology, analysis using Colaizzi's framework for phenomenological analysis<sup>2</sup></p> <p>Focus: experiences of pelvic pain</p>	<p>Themes:</p> <ul style="list-style-type: none"> <li>• Pain</li> <li>• Lifestyle adaptation</li> <li>• Emotions</li> <li>• Health professionals' support and information</li> </ul>
<p><b>Wuytack et al 2015a</b> (Ireland)</p> <p>To explore primiparous women's experiences of persistent PGP and its impact on postpartum lives (same study as <a href="#">Wuytack et al 2015b</a>)</p>	<p>23 primiparous women with PGP (onset in pregnancy, persisting for at least 3 months postpartum)</p>	<p>Interviews (3-12 months after birth)</p> <p>Thematic analysis</p> <p>Focus on experience of living with PGP (for this paper)</p>	<p>Themes:</p> <ul style="list-style-type: none"> <li>• Putting up with the pain: coping with everyday life</li> <li>• I don't feel back to normal</li> <li>• Unexpected</li> <li>• What next?</li> </ul>
<p><b>Wuytack et al 2015b</b> (Ireland)</p> <p>To explore the health-seeking behaviours of primiparous women with persistent PGP (same study as <a href="#">Wuytack et al 2015a</a>)</p>	<p>23 primiparous women with PGP (onset in pregnancy, persisting for at least 3 months postpartum)</p>	<p>Interviews</p> <p>Thematic analysis</p> <p>Focus on coping strategies, care/support offered, help/advice sought (for this paper)</p>	<p>Themes:</p> <ul style="list-style-type: none"> <li>• They didn't ask, I didn't tell <ul style="list-style-type: none"> <li>○ Lack of follow-up after birth</li> <li>○ Healthcare professionals ignore it</li> </ul> </li> <li>• Seeking advice and support <ul style="list-style-type: none"> <li>○ Talking to others</li> <li>○ Triggers to seek help</li> <li>○ Barriers to getting help</li> </ul> </li> <li>• Coping strategies <ul style="list-style-type: none"> <li>○ Self-management strategies</li> <li>○ Pain medication</li> </ul> </li> </ul>

<sup>1</sup> Newell, R., & Burnard, P. (2006). *Research for evidence-based practice*. Oxford: Blackwell.

<sup>2</sup> Colaizzi, P. F. (1978). Psychological research as the phenomenologist views it. In: Valle, R., & King, M. (Eds.) *Existential phenomenological alternatives for psychology*. Oxford: Oxford University Press.




Fig. 2. Original perinatal wellbeing framework.

Women’s *physical experiences* of LPP were dominated by pain and the impact on mobility and functioning, which tended to affect many aspects of women’s lives as well as how women view and relate to their bodies. Their *emotional experiences* were characterised by worry and concern about their condition and its impact on their lives, as well as sadness, disappointment and frustration. However, there were also positive experiences of joy, hope, and reassurance. LPP affected women’s *psychological experiences* in a number of ways, most notably their sense of control, identity and purpose.

The dimension of *time* was very evident. Women described both day-to-day and longer-term changes in wellbeing linked to LPP. The review also confirmed the importance of considering the perinatal period as a continuum, and the importance of events prior to pregnancy and the impact of perinatal experiences on wellbeing afterwards. In the theoretical review (Wadephul et al., 2020) we have suggested that space

and/or the environment may be another significant aspect of wellbeing. This review provides some supporting evidence, especially in terms of the extent to which LPP restricts women’s ability to move around.

A number of themes which did not appear to fit within the original framework were included in the ‘additional themes’ category. These included: women’s expectations, attitudes, knowledge, and understanding of pregnancy, birth, motherhood, and LPP; and how women adapted to and coped with LPP. These were subsequently integrated into the refined framework (see below).

#### Refined PWB framework

##### Societal and cultural discourses and Structures, policies, and laws

The original domain ‘culture and society’ has been expanded and elaborated in the revised framework (Fig. 3). The new *culture and soci-*



**Table 3**  
Themes and sub-themes mapped against domains and sub-domains of the original framework.

Original domains/sub-domains	Themes and sub-themes
<b>Society and culture</b>	Pregnancy as normal/natural LPP as part of pregnancy
<b>Community</b>	
Health professionals	Knowledge and information Knowledgeable vs not knowledgeable Providing information Attitudes and relationship Feeling listened to Feeling dismissed Trust, feeling safe Health care system Routine appointments, support Availability of treatment
Work and study	Impact on ability to work Difficulties/challenges Making changes, sick leave Wanting to work/study Financial implications Relationships with colleagues and employers Feeling left out Understanding/lack of understanding
Other people / strangers	Feeling judged / lack of understanding Helpful Other women with LPP Giving/receiving advice Not feeling alone
<b>Immediate environment</b>	Avoiding social interactions, negative impact Changing relationships Support from others Attitudes Baby and older children
<b>Individual</b>	Time for self/leisure Daily functioning
<b>Experiences</b>	
Physical/emodied experiences	Awareness of body Changed view of body Acknowledging physical limitations Staying active, exercise, rest Pain Physical changes Sleep, exhaustion Physical functioning, immobility Recovery
Emotional/affective experiences	Concerns, worries, anxieties, fear About the baby, labour/birth, future pregnancy About other people's reactions About pain and physical impact About uncertainty Happiness, joy, hope Feeling lonely, isolated Reassurance, relief Feeling sad, depressed, disappointed About impact on maternal role, older children About pain and loss of functioning Because others don't understand Depression About lack of information or care Self-blame, guilt, feeling inadequate or embarrassed Short-tempered, frustrated, angry
Psychological/cognitive experiences	Sense of control, self-efficacy  Self-compassion, accepting help Identity, self-perception, self-esteem Sense of purpose, attitude to life Vulnerability, feeling dependent
<b>Time and change</b>	Perinatal period as a continuum Birth as end-point Impact of birth Impact on birth and postnatal choices Life course / pre-/post-perinatal period LPP in previous pregnancy Future pregnancy Body before pregnancy/LPP Thoughts about the future Changes In pain and mobility In wellbeing
<b>Additional themes</b>	Space Dealing with LPP/adapting Adapting, making changes, coping Medication, treatment Self-care, self-management Knowledge of LPP Knowing, not knowing Information about LPP Making sense of LPP Uncertainty Women's attitudes to LPP

ety domain comprises two sub-domains: *societal and cultural discourses and structures, policies, and laws*. The former relates to the wider socio-cultural influences on women's wellbeing which underlie many other themes within the framework. For example, discourses around what a 'normal pregnancy' is (e.g. Elden et al 2013) and assumptions around LPP as a normal part of pregnancy (e.g. Wellock and Crichton 2007b, Wuytack et al 2015b) affect not only women's experiences, attitudes, and expectations, but also the availability of help and support from others, including health professionals and employers. They are also likely to influence the availability and type of treatments. Socio-cultural discourses underpin women's own attitudes and expectations of what is normal in pregnancy and what they 'should' be feeling or able to do, which can induce feelings of guilt and self-blame due to physical limitations. Not only were women sometimes told by health professionals that backpain was 'part of pregnancy ... just get on with it' (Close et al., 2016, p.5), but some women also seemed to have internalised these expectations: '... pregnancy is not a disease. Everything ought to be as before the pregnancy' (Persson et al., 2013, p. 6).

Women's experiences of wellbeing are also affected by wider socio-cultural, political, and economic structures, policies, and laws. In this review, for example, this includes the impact of sick leave, and the availability of health care and treatments. These elements were included in the 'Community' and 'immediate environment' domains in the original

framework, under health professions and employment. It became apparent during the review these elements relate to overarching structures like policies and laws, and they were moved to the wider society and culture domain.

### Relationships

In the original framework, the 'community' and 'immediate environment' domains were about different 'areas' of women's lives, e.g. work and health professionals ('community') and family and friends ('immediate environment'). However, the review suggests that this domain was primarily concerned with the nature and type of relationships with different groups of people and individuals. We have therefore combined 'community' and 'immediate environment' into a *relationship* domain.

The *relationship* domain comprises a wide range of relationships varying in nature and in significance. For example, relationships with immediate family are likely to be more significant for wellbeing than relationships with acquaintances. Consequently, there is a gradient within this domain with respect to the importance and closeness of relationships. Furthermore, the nature and quality of a relationship can change during the perinatal period; for example, friends without children may become less close, whereas those with children (or new friends with babies due at the same time) may become more significant.






Fig. 3. Revised perinatal wellbeing framework.

Women described a wide range of relationships affecting their wellbeing in positive or negative ways. For example, while some women described relationships with health professionals characterised by trust and feeling safe and listened to, others felt that their feelings and concerns were dismissed. Similarly, relationships with employers and colleagues could be supportive and understanding, or unhelpful and leading to considerable amounts of stress. Many women described support, both practical and emotional, from family members and friends as crucial for their wellbeing. On the other hand, some women also experienced unhelpful relationships and a lack of support: 'I have been so mad at him as he's not helping me out at home; in some way it's expected that I should manage' (Persson et al., 2013, p. 4). A number of women described how their experience of LPP had negatively affected relationships, particularly with friends, as they were not able to socialise like before: 'It has made me the most miserable anti-social person ... cos I'm in too much pain' (Clarkson & Adams, 2018, p. 341).

#### Individual factors

The 'individual' domain in the original framework was not very clearly defined. The nature of this domain became more apparent with this review; we redefined it as *individual factors* as it concerns women's characteristics and circumstances which are likely to affect wellbeing. Several of the 'additional themes' were also included in this domain, including women's expectations, knowledge and attitudes to LPP, and their coping strategies. Essentially, this domain is about women's characteristics and circumstances, i.e. what they bring with them. In the context of this review, this includes: their knowledge of, and attitude to, LPP; their knowledge and attitudes with regard to pregnancy, birth and motherhood; their domestic situation and whether they are caring for older children; whether they are working (or studying); and how they cope with, and adapt to, LPP.

Findings by Gutke et al. (2017) illustrate individual factors and differences between women in terms of how they adapt to, and cope with,

PGP: while some women have accepted the condition and adapted to it, others, for a variety of reasons continued to struggle to do so.

#### Experiences: physical, emotional, psychological

This domain encompasses women's physical, emotional, and psychological experiences of wellbeing. It remains the same as in the original framework, but this review has provided further nuances and richer details and has highlighted the inter-connections between physical, emotional, and psychological experiences. LPP led to considerable physical limitations, affecting activities and daily functioning. Pain stood out in intensity, affecting not only physical, but also emotional and psychological experiences. The impact of sleep also affects all three facets of experiences. Many women felt anxious or concerned about a wide range of issues, including birth and being able to look after the baby and older children. Experiencing LPP also affected women's sense of identity and how they saw themselves, particularly in terms of the maternal role and a sense of dependence on others.

There are many differences in how individual women experience wellbeing, largely depending on their specific characteristics and circumstances, i.e. the individual factors domain. As proposed in the original framework, these experiences can be both positive and negative. However, in this review negative experiences tended to dominate due to the focus on LPP and its symptoms and consequences.

#### Time

The importance of a temporal dimension was confirmed by this review: fluctuations in wellbeing over time, the perinatal period as a continuum, and the significance of a life course perspective. Perinatal wellbeing is not static but changes over time, both in the short-term and over a longer time period. Changes in wellbeing were often linked to the impact of activity levels on LPP.

Women's experiences of wellbeing with respect to LPP confirm that the perinatal period needs to be considered as a continuum. Wellbeing in each of the different parts of the perinatal period (pregnancy,

labour/birth, postnatal period) does not occur isolation, but is affected by the other periods. For example, what happens in pregnancy can affect experiences in labour and postnatally, and what happens during labour and birth often has consequences postnatally. Furthermore, women think beyond just one period. The most common example of this was women thinking ahead to labour and implications of LPP for birth or postnatally.

What happens before the pregnancy affects wellbeing during the current perinatal period and what happens during the current perinatal period affects women's wellbeing later on in life. This includes how women think ahead to future pregnancies, e.g. not wanting another pregnancy because of concerns about pain.

#### *Space and environment*

In the theoretical review (Wadehul et al., 2020) we suggested that space and the environment might be another significant factor in perinatal wellbeing. In the context of this review, space appears to be relevant in terms of restrictions on women's mobility and their ability to go out.

The environment women live in may also be relevant for their wellbeing, for example in how easy it is for them to move around, including the availability of public transport and whether they can walk to work, shops, or their children's school. One woman commented that living in an environment where her young child was safe to go out and play because neighbours would keep an eye made it easier to cope with LPP. On the other hand, issues of the wider environment may be included in the wider society and culture domain.

## **Discussion**

### *Lumbo-pelvic pain*

It is striking that the focus of the included studies is almost exclusively on PGP; only one study explores women's experiences of LBP (Close et al., 2016). While LBP is relatively common, it is also less well defined than PGP and often remains formally un-diagnosed. Furthermore, LBP seems to have been normalised in pregnancy, and possibly postnatally. A key difference between PGP and LBP was identified by the one paper which included LBP (Close et al., 2016): while women expected LBP during pregnancy and therefore had an attitude of normality towards it, they were not aware of PGP, which often seemed to surprise them and was described by one woman as 'frightening'. Pelvic girdle pain tends to be more painful, acute and alarming than LBP and can therefore have a considerable impact on functioning. Given that the majority of studies focused on PGP, overall it is difficult to know to what extent insights from PGP studies are applicable in terms of women's experiences to LBP. It is quite likely that the intense pain and unexpected nature of PGP may have a more significant impact on wellbeing compared to LBP. However, the focus of this review was to explore ways in which a relatively common adverse situation could inform us about women's wellbeing. While there are certainly differences between the conditions as stated above, they both have a considerable impact on wellbeing.

### *The revised framework: 'fit' with existing research*

Pregnancy, birth and early motherhood are not just physical, biological experiences, but experiences which take place within, and are shaped by, the social and cultural context. They are therefore influenced by discourses within this context (Miller, 2007) and, in turn, women's wellbeing is affected by these discourses. Review findings suggest that discourses around what is normal in pregnancy affect women's experiences: several of the included studies report that women were told that back pain in pregnancy is normal and that they would just have to put up with it. Bessett (2010) describes a discourse of maternal sacrifice involving 'cultural pressures to "suffer nobly" the symptoms of pregnancy, no

matter how uncomfortable' (Bessett, 2010, p. 370). She suggests that going against this discourse, for example by complaining about discomfort and pain, threatens women's identity as a 'good mother'. This may have a direct impact on women's emotional wellbeing and a physical impact if women do not seek help and support as needed. This exemplifies how socio-cultural discourses and assumptions around pregnancy and motherhood can affect women's experiences of wellbeing. Other discourses are likely to be pertinent in other contexts.

The 'pregnancy as health' discourse (Fredriksen et al., 2010), which relates to expectations among pregnant women that they will remain healthy, fit, and able to work during pregnancy, is also reflected in findings. This can result in women being reluctant to slow down or take sick leave and may lead to feelings of failure and inadequacy.

Supportive and protective social and employment policies have the potential to increase maternal wellbeing (Tsai and Tai, 2018). This review illustrates how national policies can affect wellbeing during the perinatal period, particularly paid sick leave policies. It is notable that in the included papers discussion of taking sick leave usually referred to pregnancy. All papers originated in countries which have relatively generous policies on maternity leave; in this context, taking sick leave postnatally is less relevant. The extent to which women are able to make adaptations at work, including more flexible working, working part time, or making changes to what they do at work, can also have a significant impact on their wellbeing.

### *Relationships*

The significance for PWB of relationships is evident. What appears to matter is the *quality* of a relationship and whether it is positive, supportive, and understanding, or has a negative impact on wellbeing. The emphasis in research and clinical practice tends to be on relationships with a partner; however, other relationships are also significant, including relationships with children, close family and friends, neighbours, work colleagues, extended family, and health professionals. The quality and relative significance of particular relationships is dynamic and changes throughout the perinatal period. For example, in the included papers many of the women talked about how their relationships with friends changed: as they transition to parenthood, they may become less close to friends without children, while other pregnant women and mothers with children at a similar age become more significant (Jones et al., 2014).

### *Individual factors*

This review highlights individual differences between women and the impact this may have on their experiences of wellbeing, particularly in terms of coping strategies, expectations, attitude, and knowledge, but also their work and family circumstances. There is some research in this area, suggesting, for example, that coping strategies (e.g. George et al., 2013; Lafarge et al., 2013) and expectations (Henshaw et al., 2014) can affect aspects of perinatal wellbeing. However, overall these factors, which may explain how different women experience similar situations in different ways, is under-researched. Other individual factors which may affect wellbeing include women's personality, personal history, and women's environment (where they live and their housing situation).

### *Experiences of wellbeing*

Women's experiences of wellbeing are affected by the outer sections of the framework; these provide the *context* of wellbeing, while the experiences can be considered the *core* of wellbeing. This review demonstrates how closely linked different aspects of experiences of wellbeing are. The distinction between physical, emotional, and cognitive experiences is, to some extent, artificial; it is not always easy to make a clear distinction between these elements. This reflects the mind-body dualism which is so dominant in research and practice. However, while we would argue that it is important to consider the holistic nature of wellbeing, it is arguably difficult to capture this in research.

### Time

Both the context and the core of wellbeing have a temporal dimension: they change over time and experiences at different time periods have an impact on other time periods. Studies and clinical assessments of perinatal wellbeing, or aspects of wellbeing, tend to capture only a snapshot of wellbeing at a particular point of in time. However, it is clear that wellbeing is dynamic and fluctuates across the perinatal period (Newham and Martin, 2013) and over shorter periods of time. Capturing the dynamic, fluctuating nature of wellbeing may be enhanced by the use of frequent 'ecological momentary assessments' with the aid of diaries or digital technology (Newham and Martin, 2013). Changes in wellbeing in the perinatal period also mean that it is important to report and take into account the time period to which an assessment of wellbeing refers; for example, there can be considerable differences between early and late pregnancy, or the immediate days after birth or several month postpartum.

This review illustrates several ways in which the different parts of the perinatal period can be related such as women's concerns during pregnancy about giving birth with LPP or about the impact of labour on their postnatal wellbeing. It is notable that discussion of labour and birth was largely absent otherwise. This may be because during labour LPP is less of a focus. However, research into women's experiences, captured retrospectively, of LPP during labour and birth would widen our understanding of the issues around LPP.

The perinatal period does not exist within a vacuum, but needs to be located within the wider temporal context of women's lives. In the studies included in this review, women talked about this with respect to previous or future pregnancies and births. For example, women compared their current experiences of PGP to those in a previous pregnancy. Several women also talked about how the experiences of pain due to PGP affected their decisions about future pregnancies. A life course perspective may provide further insights into women's experiences and needs.

### Space/environment

We feel that consideration of the impact of space and environment on wellbeing would benefit from further exploration. Aspects of it may fit in within the experiences of wellbeing and individual factors domain as well as the wider socio-cultural context. It is evident from this review that LPP can restrict women's mobility and therefore ability to travel, including commuting, going shopping, and taking children to school. While this relates to where women live (individual factors) and is part of women's physical experience of wellbeing, it is also linked to the wider infrastructure and facilities at a societal level: the wider environment can make it harder, or easier, for women to deal with their lack of mobility. In this sense, the environment, in its widest sense, can be as disabling as the symptoms of LPP itself.

### Using the framework: research, practice, policy

The aim of this review was to explore the utility of the conceptual framework of wellbeing and to develop it by applying it to a particular perinatal situation which challenges wellbeing, i.e. lower back pain. By utilising framework analysis we were able to explore the ways in which women's own experiences fit the framework. Our findings suggest that the proposed multi-dimensional framework has a good fit; women's experiences have allowed us to develop and refine the framework to more closely reflect the actual lived experience of women. It allowed us to understand the challenges to wellbeing as well as those factors which support it. Arguably this gives a fuller understanding of wellbeing and allows a more elaborate development of the framework.

While this review aimed to employ the revised framework with respect to a very specific example, perinatal wellbeing in the context of LPP, the framework can be used more widely. It can be applied to perinatal wellbeing in specific contexts, conditions, or groups of women. On the other hand, it can also be used to illustrate, explore, and explain

perinatal wellbeing in a more general sense. As such, the framework can be used in a practical, applied way, or in a more theoretical way.

Individual research studies are likely to focus on separate aspects of wellbeing; however, awareness of the wider context and the influence of women's individual circumstances is important. This framework enables a comprehensive view of wellbeing during the perinatal period, including contributing factors and how women experience wellbeing. It also addresses the wider socio-cultural context and the importance of women's individual characteristics and circumstances.

Within clinical practice, the framework highlights the importance of an awareness and understanding of the wider context of women's lives and of their specific individual circumstances, characteristics, and needs. This is of significance not just to how health professionals care for women, but also for how maternity care is organised. In terms of policy, an understanding of all factors which can affect women's wellbeing is vital, including sick leave, maternity leave, and health care. Finally, the framework also has the potential to assist in the development of interventions designed to support women's wellbeing, underlining the importance of a comprehensive, holistic approach.

### Strengths and limitations

The geographical spread of included studies was limited to north-western Europe. In other countries, particularly lower and middle income countries, other issues are likely to be relevant. However, the aim of the framework is to identify domains rather than specific issues for women so while the content of the domains may vary across contexts and cultures, it is expected that the broad areas will remain the same, although different domains may come to the fore in different contexts.

This review uses a concrete example (LPP) to illustrate conceptual work on PWB. It builds on previous work (Wadehul et al., 2020), using a thorough and methodical approach to further develop and refine the initial framework. The refined framework of PWB is firmly grounded in the research literature. The review moves the development of the framework closer to women's experiences and allows it to be informed by women's voices rather than solely abstract concepts.

### Future directions

It is important to evaluate how well the framework applies to other situations or groups of women. Of particular note are the geographical limitations, as noted above; the differences, and similarities, of experiences of women in different cultures and/or less economically developed countries needs to be explored.

While the revised framework builds on the experiences of women who participated in the included studies, this was done within the context of the original framework. We feel it is important to further validate, and potentially revise, the framework using women's experiences and their own conceptualisations of perinatal wellbeing as a starting point. A mixed methods study with a large qualitative element with this aim has recently been completed. This also includes the perspectives of health professionals involved in the care of perinatal women.

### Conclusions

Pelvic girdle pain related to the perinatal period can have a profound impact on women's wellbeing in a number of ways; this is affected by a wide range of factors. The review highlights the lack of research into women's experiences of lower back pain, indicating a need for further research.

This review has illustrated the complex nature of wellbeing by using LPP as an example. It has demonstrated the utility and appropriateness of the PWB framework. It has allowed us to add detail to the framework and refine its components. Further work is required to evaluate how well the framework fits with women's experiences of perinatal wellbeing in general and how appropriate the framework is in contexts which are

different from the rather narrow geographical settings of the included studies.

The revised framework can be used to map existing research and guide future research, including issues which need further exploration with respect to how they affect perinatal wellbeing, such as socio-cultural discourses, the role of the environment, and women's past experiences. The framework has the potential to form the basis for further theoretical work on the concept of perinatal wellbeing. We anticipate, and welcome, further changes to the framework based on new and existing research into women's experiences.

#### Declaration of Competing Interest

None declared.

#### Ethical approval

Not applicable.

#### Funding sources

The first author's post was funded by the University of Hull; there was no specific, additional funding for this project.

#### Supplementary materials

Supplementary material associated with this article can be found, in the online version, at doi:[10.1016/j.midw.2021.103031](https://doi.org/10.1016/j.midw.2021.103031).

#### References

- Ayers, S., Olander, E.K., 2013. What are we measuring and why? Using theory to guide perinatal research and measurement. *J. Reprod. Infant Psychol.* 31 (5), 439–448. doi:[10.1080/02646838.2013.834041](https://doi.org/10.1080/02646838.2013.834041).
- Bessett, D., 2010. Negotiating normalization: The perils of producing pregnancy symptoms in prenatal care. *Soc. Sci. Med.* 71 (2), 370–377. doi:[10.1016/j.socscimed.2010.04.007](https://doi.org/10.1016/j.socscimed.2010.04.007).
- Buchbinder, R., Blyth, F.M., March, L.M., Brooks, P., Woolf, A.D., Hoy, D.G., 2013. Placing the global burden of low back pain in context. *Best Pract. Res. Clin. Rheumatol.* 27 (5), 575–589. doi:[10.1016/j.berh.2013.10.007](https://doi.org/10.1016/j.berh.2013.10.007).
- Carroll, C., Booth, A., Cooper, K., 2011. A worked example of "best-fit" framework synthesis: a systematic review of views concerning the taking of potential chemopreventive agents. *BMC Med. Res. Method.* 11. doi:[10.1186/1471-2288-11-29](https://doi.org/10.1186/1471-2288-11-29).
- Carroll, C., Booth, A., Leaviss, J., Rick, J., 2013. Best fit" framework synthesis: refining the method. *BMC Med. Res. Method.* 13 (1), 37. doi:[10.1186/1471-2288-13-37](https://doi.org/10.1186/1471-2288-13-37).
- Clarkson, C.E., Adams, N., 2018. A qualitative exploration of the views and experiences of women with Pregnancy related Pelvic Girdle Pain. *Physiotherapy* 104 (3), 338–346. doi:[10.1016/j.physio.2018.05.001](https://doi.org/10.1016/j.physio.2018.05.001).
- Close, C., Sinclair, M., Liddle, D., Mc Cullough, J., Hughes, C., 2016. Women's experience of low back and/or pelvic pain (LBPP) during pregnancy. *Midwifery* 37, 1–8. doi:<https://doi.org/10.1016/j.midw.2016.03.013>
- Crichton, M.A., Wellock, V.K., 2008. Pain, disability and symphysis pubis dysfunction: women talking. *Evidence Based Midwifery* 6 (1), 9–18.
- Critical Appraisal Skills Programme, 2017. CASP Qualitative Checklist [online] Retrieved from [http://docs.wixstatic.com/ugd/dded87\\_25658615020e427da194a325e7773d42.pdf](http://docs.wixstatic.com/ugd/dded87_25658615020e427da194a325e7773d42.pdf).
- Elden, H., Lundgren, I., Robertson, E., 2013. Life's pregnant pause of pain: pregnant women's experiences of pelvic girdle pain related to daily life: A Swedish interview study. *Sexual Reprod. Healthc.* 4 (1), 29–34. doi:[10.1016/j.srhc.2012.11.003](https://doi.org/10.1016/j.srhc.2012.11.003).
- Elden, H., Lundgren, I., Robertson, E., 2014. The pelvic ring of pain: pregnant women's experiences of severe pelvic girdle pain: An interview study. *Clin. Nursing Stud.* 2 (2), 30–41.
- Elkins-Bushnell, R., Boyle, P., 2019. The experience of women living with pelvic girdle pain and participation in activity after childbirth. *International Journal of Therapy and Rehabilitation* 26 (1), 1–10. doi:[10.12968/ijtr.2017.0158](https://doi.org/10.12968/ijtr.2017.0158).
- Engeset, J., Stuge, B., Fegran, L., 2014. Pelvic girdle pain affects the whole life—a qualitative interview study in Norway on women's experiences with pelvic girdle pain after delivery. *BMC Res. Notes* 7 (1), 686. doi:[10.1186/1756-0500-7-686](https://doi.org/10.1186/1756-0500-7-686).
- Fredriksen, E.H., Harris, J., Moland, K.M., Sundby, J., 2010. A Defeat not to Be Ultra-Fit": expectations and experiences related to pregnancy and employment in contemporary Norway. *NORA - Nordic J. Feminist Gender Res.* 18 (3), 167–184. doi:[10.1080/08038740.2010.498766](https://doi.org/10.1080/08038740.2010.498766).
- Fredriksen, E.H., Moland, K.M., Sundby, J., 2008. Listen to your body": A qualitative text analysis of internet discussions related to pregnancy health and pelvic girdle pain in pregnancy. *Patient Educ. Couns.* 73 (2), 294–299. doi:[10.1016/j.pec.2008.02.002](https://doi.org/10.1016/j.pec.2008.02.002).
- George, A., Luz, R.F., De Tychey, C., Thilly, N., Spitz, E., 2013. Anxiety symptoms and coping strategies in the perinatal period. *BMC Pregnancy Childbirth* 13 (1), 233. doi:[10.1186/1471-2393-13-233](https://doi.org/10.1186/1471-2393-13-233).
- Gutke, A., Boissonnault, J., Brook, G., Stuge, B., 2018. The Severity and Impact of Pelvic Girdle Pain and Low-Back Pain in Pregnancy: A Multinational Study. *J. Women's Health* 27 (4), 510–517. doi:[10.1089/jwh.2017.6342](https://doi.org/10.1089/jwh.2017.6342).
- Gutke, A., Bullington, J., Lund, M., Lundberg, M., 2017. Adaptation to a changed body. Experiences of living with long-term pelvic girdle pain after childbirth. *Disabil. Rehabil.* 40 (25), 3054–3060. doi:[10.1080/09638288.2017.1368724](https://doi.org/10.1080/09638288.2017.1368724).
- Gutke, A., Lundberg, M., Østgaard, H.C., Öberg, B., 2011. Impact of postpartum lumbopelvic pain on disability, pain intensity, health-related quality of life, activity level, kinesophobia, and depressive symptoms. *Eur. Spine J.* 20 (3), 440–448. doi:[10.1007/s00586-010-1487-6](https://doi.org/10.1007/s00586-010-1487-6).
- Hartvigsen, J., Hancock, M.J., Kongsted, A., Louw, Q., Ferreira, M.L., Genevay, S., ... Woolf, A., 2018. What low back pain is and why we need to pay attention. *Lancet North Am. Ed.* 391 (10137), 2356–2367. doi:[10.1016/S0140-6736\(18\)30480-X](https://doi.org/10.1016/S0140-6736(18)30480-X).
- Henshaw, E.J., Fried, R., Teeters, J.B., Siskind, E.E., 2014. Maternal expectations and postpartum emotional adjustment in first-time mothers: results of a questionnaire survey. *J. Psychosomatic Obstetr. Gynecol.* 35 (3), 69–75. doi:[10.3109/0167482X.2014.937802](https://doi.org/10.3109/0167482X.2014.937802).
- Jones, C., Jomeen, J., Hayter, M., 2014. The impact of peer support in the context of perinatal mental illness: a meta-ethnography. *Midwifery* 30 (5), 491–498. doi:[10.1016/j.midw.2013.08.003](https://doi.org/10.1016/j.midw.2013.08.003).
- Lafarge, C., Mitchell, K., Fox, P., 2013. Perinatal grief following a termination of pregnancy for foetal abnormality: the impact of coping strategies. *Prenat. Diagn.* 33 (12), 1173–1182. doi:[10.1002/pd.4218](https://doi.org/10.1002/pd.4218).
- Mackenzie, J., Murray, E., Lusher, J., 2018. Women's experiences of pregnancy related pelvic girdle pain: a systematic review. *Midwifery* 56, 102–111. doi:[10.1016/j.midw.2017.10.011](https://doi.org/10.1016/j.midw.2017.10.011).
- Miller, T., 2007. Is this what motherhood is all about?": Weaving experiences and discourse through transition to first-time motherhood. *Gender Soc.* 21 (3), 337–358. doi:[10.1177/0891243207300561](https://doi.org/10.1177/0891243207300561).
- Newham, J.J., Martin, C.R., 2013. Measuring fluctuations in maternal well-being and mood across pregnancy. *J. Reprod. Infant Psychol.* 31 (5), 531–540. doi:[10.1080/02646838.2013.834040](https://doi.org/10.1080/02646838.2013.834040).
- Persson, M., Winkvist, A., Dahlgren, L., Mogren, I., 2013. Struggling with daily life and enduring pain": a qualitative study of the experiences of pregnant women living with pelvic girdle pain. *BMC Pregnancy Childbirth* 13 (1), 111. doi:[10.1186/1471-2393-13-111](https://doi.org/10.1186/1471-2393-13-111).
- Shepherd, J., 2005. Symphysis pubis dysfunction: a hidden cause of morbidity. *Br. J. Midwifery* 13 (5), 301–307. doi:[10.12968/bjom.2005.13.5.18092](https://doi.org/10.12968/bjom.2005.13.5.18092).
- Tsai, M.-C., Tai, T.-o., 2018. How are mothers faring across the globe? Constructing a new mothers' well-being index and assessing its validity. *Appl. Res. Qual. Life* 13, 647–670. doi:[10.1007/s11482-017-9550-7](https://doi.org/10.1007/s11482-017-9550-7).
- Varley, M., Hunter, L., 2019. How does pelvic girdle pain impact on a woman's experience of her pregnancy and the puerperium? *Evidence Based Midwifery* 17 (2), 60–70.
- Vleeming, A., Albert, H., Østgaard, H., Sturesson, B., Stuge, B., 2008. European guidelines for the diagnosis and treatment of pelvic girdle pain. *Eur. Spine J.* 17 (6), 794–819. doi:[10.1007/s00586-008-0602-4](https://doi.org/10.1007/s00586-008-0602-4).
- Wadephul, Franziska, Glover, Lesley, Jomeen, Julie, 2020. Conceptualising women's perinatal well-being: A systematic review of theoretical discussions. *Midwifery* 81, 102629. doi:[10.1016/j.midw.2019.102598](https://doi.org/10.1016/j.midw.2019.102598).
- Wellock, V.K., Crichton, M.A., 2007a. Understanding pregnant women's experiences of symphysis pubis dysfunction: the effect of pain. *Evidence-Based Midwifery* 5 (2), 40–46.
- Wellock, V.K., Crichton, M.A., 2007b. Symphysis pubis dysfunction: women's experiences of care. *Br. J. Midwifery* 15 (8), 494–499. doi:[10.12968/bjom.2007.15.8.24390](https://doi.org/10.12968/bjom.2007.15.8.24390).
- Wu, W.H., Meijer, O.G., Uegaki, K., Mens, J.M.A., van Dieën, J.H., Wuisman, P.I.J.M., Østgaard, H.C., 2004. Pregnancy-related pelvic girdle pain (PPP), I: Terminology, clinical presentation, and prevalence. *Eur. Spine J.* 13 (7), 575–589. doi:[10.1007/s00586-003-0615-y](https://doi.org/10.1007/s00586-003-0615-y).
- Wuytack, F., Curtis, E., Begley, C., 2015a. Experiences of first-time mothers with persistent pelvic girdle pain after childbirth: descriptive qualitative study. *Phys. Ther.* 95 (10), 1354–1364. doi:[10.2522/ptj.20150088](https://doi.org/10.2522/ptj.20150088).
- Wuytack, F., Curtis, E., Begley, C., 2015b. The health-seeking behaviours of first-time mothers with persistent pelvic girdle pain after childbirth in Ireland: A descriptive qualitative study. *Midwifery* 31 (11), 1104–1109. doi:[10.1016/j.midw.2015.07.009](https://doi.org/10.1016/j.midw.2015.07.009).