



Metal complexes of 1,10-phenanthroline-5,6-dione alter the susceptibility of the yeast *Candida albicans* to Amphotericin B and Miconazole

Ahmed Eshwika¹, Barry Coyle², Michael Devereux³, Malachy McCann² & Kevin Kavanagh^{1,*}

¹Medical Mycology Unit, NICB, Department of Biology, National University of Ireland Maynooth, Co. Kildare, Ireland; ²Department of Chemistry, National University of Ireland Maynooth, Co. Kildare, Ireland; ³Dublin Institute for Technology, Cathal Brugha St., Dublin 1; *Author for correspondence (Tel: 353-1-708 3859, Fax: 353-1-708 3845, E-mail: kevin.kavanagh@may.ie.)

Received 15 October 2003; Accepted 28 November 2003

Abstract

Growth of the pathogenic yeast *Candida albicans* in sub-MIC (minimum inhibitory concentration) levels of $\text{Cu}(\text{ClO}_4)_2 \cdot 6\text{H}_2\text{O}$ and $[\text{Cu}(\text{phen})_3](\text{ClO}_4)_2 \cdot 4\text{H}_2\text{O}$ (phen = 1,10-phenanthroline-5,6-dione) increased the concentration of miconazole and amphotericin B required to achieve the MIC_{90} whereas pre-growth in AgClO_4 and $[\text{Ag}(\text{phen})_2]\text{ClO}_4$ resulted in a small decrease in the relevant MIC_{90} values. The copper complexes reduce the oxygen consumption of *C. albicans* while the silver complexes increase oxygen consumption. In addition, pre-growth of cells in the copper complexes resulted in a lower ergosterol content while the silver complexes induced an elevation in ergosterol synthesis.

The ability of copper and silver complexes to alter the susceptibility of *C. albicans* to miconazole and amphotericin B may be influenced by their action on respiration, since reduced respiration rates correlate with reduced cellular ergosterol which is the target for amphotericin B. Lower levels of ergosterol have previously been associated with elevated tolerance to this drug. In the case of reduced sensitivity to miconazole, tolerance may be mediated by lower ergosterol synthesis giving rise to fewer toxic side products once biosynthesis is inhibited by miconazole.

Abbreviations: phen = 1,10-phenanthroline-5,6-dione. phen = 1,10-phenanthroline

Introduction

The yeast *Candida albicans* is responsible for a range of superficial and systemic diseases in the immunocompromised patient. Conventional therapies for the control of these diseases rely upon the use of azole and polyene drugs which target ergosterol biosynthesis and alter membrane permeability, respectively (White *et al.*, 1998, Abu Salah, 1996). In recent years, the appearance of fungal strains manifesting resistance to conventional drugs (Canuto & Rodero, 2002) has prompted the search for novel anti-fungals with modes of action distinct to the existing range of anti-fungals.

Metal-based drugs represent a novel group of anti-fungal agents with potential applications for the control of fungal infections. Previous work in our laboratories has demonstrated that in RPMI medium at

37°C the metal-based drugs $[\text{Cu}(\text{phen})_2(\text{mal})] \cdot 2\text{H}_2\text{O}$, $[\text{Mn}(\text{phen})_2(\text{mal})] \cdot 2\text{H}_2\text{O}$ and $[\text{Ag}(\text{phen})_2]\text{ClO}_4$ (phen = 1,10-phenanthroline) inhibit the growth of *C. albicans* by around 95% at a concentration of 5 µg/ml (McCann *et al.* 2000; Coyle *et al.* 2003a). It was established that both metal-free phen and the metal-phen complexes affect mitochondrial function, retard the synthesis of cytochromes b and c and uncouple respiration. Treatment of fungal cells with the Cu(II) and Ag(I) complexes resulted in a reduced amount of ergosterol in the cell membrane and subsequent increase in its permeability. Cells exposed to metal-free phen and the Cu(II) and Mn(II) complexes (but not the Ag(I) complex) demonstrated an elevation in oxygen uptake. The general conclusion was that the drugs damage mitochondrial function and uncouple respiration. Fur-

thermore, the fact that the drugs were not uniformly active suggested that their bioactivity had a degree of metal-ion dependency.

More recently, metal-free phenidio (phenidio = 1,10-phenanthroline-5,6-dione) and the Ag(I) complex $[\text{Ag}(\text{phenidio})_2]\text{ClO}_4$ have been shown to cause extensive, non-specific DNA cleavage to *C. albicans*, disrupt cell division and induce gross distortions in fungal cell morphology (Coyle *et al.*, 2003b). Preliminary experiments on cultured human cancer cells produced IC_{50} values of $0.008 \mu\text{g/ml}$ ($0.04 \mu\text{M}$) and $0.025 \mu\text{g/ml}$ ($0.40 \mu\text{M}$) for metal-free phenidio and $[\text{Ag}(\text{phenidio})_2]\text{ClO}_4$, respectively (Coyle *et al.* 2003b). Studies by Igdaloff *et al.* (1983) revealed that both phenidio and its isomer, 1,7-phenanthroline-5,6-dione, inhibit the growth of S49 mouse lymphoma cells and S110 mouse cells, and although it was postulated that inhibition of DNA and RNA syntheses were major components of the cytotoxic effects the phenanthrolines were presumed to have more than one mode of action.

Metal-based drugs have well established fungistatic and fungicidal effects and the aim of the work presented here was to evaluate the possibility of using Cu (II) and Ag (I) phenidio complexes in combination with azole and polyene drugs for the control of *C. albicans*.

Materials and methods

Fungal isolate and culture conditions

C. albicans MEN (a kind gift from Dr. David Kerridge, Cambridge, UK) was grown to the stationary phase (approximately 1.5×10^8 cells/ml) at 30°C overnight in YEPD broth (2% (w/v) glucose (Sigma Aldrich Chemical Co., Dorset, UK), 2% (w/v) bacteriological peptone (Sigma Aldrich) and 1% (w/v) yeast extract (Sigma Aldrich)) in an orbital incubator at 200 rpm.

Drugs

Chemicals were obtained from commercial sources and used without further purification. $\text{Cu}(\text{ClO}_4)_2 \cdot 6\text{H}_2\text{O}$ was purchased from Sigma Aldrich and used without further purification. $[\text{Cu}(\text{phenidio})_3](\text{ClO}_4)_2 \cdot 4\text{H}_2\text{O}$ and $[\text{Ag}(\text{phenidio})_2]\text{ClO}_4$ were synthesized in accordance with the procedures outlined in McCann *et al.* (2003).

Anti-Candida susceptibility testing of metal-based drugs

Solutions of water-soluble copper and silver complexes were prepared by dissolving 0.02 g of the solid in sterile distilled water to yield a stock solution of $200 \mu\text{g/ml}$. The solutions were filter sterilised using a Millipore membrane filter ($0.45 \mu\text{m}$). Stationary phase cultures of *C. albicans* were harvested by centrifugation ($2220 \times g$ for 5 min in a Beckmann GS-6 centrifuge), washed twice with PBS and resuspended at a final density of 1×10^6 cells/ml. Cell suspension ($100 \mu\text{l}$) was added to each well of a 96-well microtitre plate except the first column (control medium). Serial dilutions of metal-based drugs were added to rows of wells to construct a concentration gradient from 50 – $0.78 \mu\text{g/ml}$ and the plates were incubated at 30°C for 24 h. The absorbance at 450 nm was determined using a MRX spectrophotometer (Dynax Technology, Chantilly, VA, USA) and the concentration that was capable of inhibiting growth by 90% (MIC_{90}) relative to the control was calculated.

Amphotericin B and miconazole susceptibility testing

Yeast cultures were grown to the stationary phase in antibiotic medium 3 (AB 3, Oxoid) overnight at 30°C and 200 rpm, harvested by centrifugation and diluted to 1×10^6 /ml. Cells (1×10^5 in $100 \mu\text{l}$) were added to each row of a 96-well plate containing amphotericin B in serial dilutions (Sigma-Aldrich) from 2.5 – $0.0048 \mu\text{g/ml}$ in AB3 medium.

Miconazole susceptibility testing was performed using stationary phase *C. albicans* that had been grown in RPMI medium overnight (Sigma Aldrich). Cells ($100 \mu\text{l}$ of 1×10^6 /ml) were added to each row of a 96-well plate containing miconazole, serially diluted in RPMI medium from 20 – $0.19 \mu\text{g/ml}$.

All plates were incubated at 30°C for 24 h and the optical density was read at 450 nm using a MRX spectrophotometer (Dynax Technology, Chantilly, VA, USA).

Oxygen consumption

Cells were grown in YEPD broth supplemented with sub-MIC levels of silver or copper complexes at 30°C for 24 h, harvested by centrifugation and re-suspended at a density of 1×10^8 /ml in phosphate buffered saline (PBS, pH 7.2). A Clark Type oxygen electrode (Ranks Brothers, Cambridge, UK) was employed to determine the respiration rate of cells. The rate of oxygen




Fig 1. Effect of metal-based drugs on the growth of *C. albicans*. Cells of *C. albicans* were exposed to increasing concentrations of MBD and the effect on growth after 24 h incubation was determined.

consumption is expressed as the number of μmoles of oxygen consumed per thirty sec per 1×10^8 cells.

Sterol extraction and analysis

Sterols were extracted according to the method of Arthington-Skaggs *et al.* (1999). Stationary phase cells (1.3 g wet weight) were harvested and washed with PBS (pH 7.2). Cells were re-suspended in 20% (w/v) KOH and 60% (v/v) ethanol and placed in a shaking water bath (80–90°C) for 1.5 h. Heptane was added to the solution which was then agitated for 10 sec and the aqueous layer removed. The sterol content of the hexane layer was quantified using a dual beam spectrophotometer over the range 250–300 nm.

Statistical analysis

The Kruskal–Wallis test was performed, where appropriate, on all results using the SigmaStat Statistical Analysis System (Version 1.00). Values are presented as \pm SE of the mean of data from three independent experiments.

Results

The effect of selected Cu(II) and Ag(I) phendio complexes on the growth of *C. albicans* was assessed. The results indicate (Figure 1) that AgClO_4 and $[\text{Ag}(\text{phendio})_2]\text{ClO}_4$ display MIC_{90} values of approximately 6.0 $\mu\text{g/ml}$ while $\text{Cu}(\text{ClO}_4)_2 \cdot 6\text{H}_2\text{O}$ and $[\text{Cu}(\text{phendio})_3](\text{ClO}_4)_2 \cdot 4\text{H}_2\text{O}$ display MIC_{90} values

of >50 and 46 $\mu\text{g/ml}$, respectively. In all subsequent assays, concentrations corresponding to 1/4 MIC_{90} values of these complexes were employed i.e. AgClO_4 and $[\text{Ag}(\text{phendio})_2]\text{ClO}_4$ at a concentration of 1.5 $\mu\text{g/ml}$ and $\text{Cu}(\text{ClO}_4)_2 \cdot 6\text{H}_2\text{O}$ and $[\text{Cu}(\text{phendio})_3](\text{ClO}_4)_2 \cdot 4\text{H}_2\text{O}$ at concentrations of 12.5 $\mu\text{g/ml}$ and 11.5 $\mu\text{g/ml}$, respectively. Sub- MIC values were employed in all subsequent assays so that any effect on drug susceptibility would not be due to the fungistatic effects of these compounds

Experiments were performed to determine whether pre-growth of *C. albicans* in the presence of different metal complexes affected the subsequent susceptibility of cells to azole and/or polyene drugs. Cells were pre-grown in medium supplemented with 1/4 MIC_{90} of the Cu(II) and Ag(I) phendio complexes for 24 h at 30°C and harvested by centrifugation. Anti-fungal susceptibility assays were performed as described. The results (Figure 2) indicate that pre-growth of *C. albicans* in medium supplemented with 12.5 $\mu\text{g/ml}$ $\text{Cu}(\text{ClO}_4)_2 \cdot 6\text{H}_2\text{O}$ or 11.5 $\mu\text{g/ml}$ $[\text{Cu}(\text{phendio})_3](\text{ClO}_4)_2 \cdot 4\text{H}_2\text{O}$ significantly decreases the susceptibility of *C. albicans* to miconazole ($p < 0.001$) (control $\text{MIC}_{90} = 7.94 \pm 1.2 \mu\text{g/ml}$, cells pre-grown in 1/4 MIC $\text{Cu}(\text{ClO}_4)_2 \cdot 6\text{H}_2\text{O}$ $\text{MIC}_{90} = 10.9 \pm 0.2 \mu\text{g/ml}$, cells pre-grown in 1/4 MIC $[\text{Cu}(\text{phendio})_3](\text{ClO}_4)_2 \cdot 4\text{H}_2\text{O}$ $\text{MIC}_{90} = 11.4 \pm 0.3 \mu\text{g/ml}$). In the case of the silver complexes, pre-growth of *C. albicans* in 1.5 $\mu\text{g/ml}$ AgClO_4 or $[\text{Ag}(\text{phendio})_2]\text{ClO}_4$ increases the susceptibility of the cells to miconazole. Cells pre-grown in 1/4 MIC_{90} AgClO_4 show a miconazole MIC_{90} value of $3.24 \pm 0.09 \mu\text{g/ml}$ while those pre-grown in




Fig 2. Susceptibilities to Miconazole of *C. albicans* pre-grown in the presence of sub-MIC₉₀ levels of metal complexes. The susceptibility of *C. albicans* to miconazole was determined after cells had been first pre-grown in sub-MIC₉₀ levels of metal-based drug.

1/4 MIC₉₀ [Ag(phendio)₂]ClO₄ demonstrate a MIC₉₀ value of 4.95 ± 0.27 µg/ml.

The susceptibility of *C. albicans* to the polyene drug amphotericin B was also affected following pre-growth of the cells in medium supplemented with copper or silver complexes at 1/4 MIC₉₀ values (Figure 3). In the case of pre-growth in 1/4 MIC₉₀ Cu(ClO₄)₂·6H₂O the amount of amphotericin B required to achieve the MIC₉₀ increases to 0.029 ± 0.007 µg/ml from 0.024 ± 0.003 µg/ml (control) and following pre-growth in a similar concentration of [Cu(phendio)₃](ClO₄)₂·4H₂O increases to 0.03 ± 0.003 µg/ml ($p = 0.238$). In the case of pre-exposure to the silver complexes there is a decrease in the amount of amphotericin B required to achieve the MIC₉₀. Pre-growth of *C. albicans* in either of the silver complexes reduces the amount of amphotericin B required to achieve the MIC₉₀ from 0.024 ± 0.003 µg/ml to 0.019 ± 0.001 µg/ml ($p = 0.238$) (Figure 3).

Azoles, such as miconazole, inhibit the action of lanosterol 14 α -demethylase which controls an essential intermediate step in the biosynthesis of ergosterol (White *et al.*, 1998, Daum *et al.*, 1998). In contrast, polyenes, such as amphotericin B, bind to ergosterol in the fungal cell membrane creating pores through which cell constituents may escape (Abu Salah, 1996). Ergosterol biosynthesis is an oxygen dependent process and also requires NADPH

which is synthesised in the aerobically respiring mitochondrion. Consequently, experiments were performed to establish whether the altered susceptibility of *C. albicans* to miconazole and amphotericin B was due to the ability of the copper or silver complexes to alter respiration. The results (Figure 4) demonstrate that pre-growth of cells in 1/4 MIC₉₀ of the copper complexes (Cu(ClO₄)₂·6H₂O and [Cu(phendio)₃](ClO₄)₂·4H₂O) results in a reduction in the cellular respiration rate ($p < 0.001$) whereas growth of cells in the presence of the silver complexes (AgClO₄ and [Ag(phendio)₂]ClO₄) increases the respiration rate of *C. albicans* ($p < 0.001$).

The ergosterol biosynthetic pathway is the target of the azole drugs (White *et al.* 1998) whilst polyenes bind ergosterol in the cell membrane and create pores (Abu Salah, 1996). Quantification of ergosterol in cells grown in medium supplemented with sub MIC₉₀ concentrations of the metal-based drugs demonstrated that growth in the presence of copper complexes resulted in a reduced ergosterol content, while growth in medium supplemented with the silver complexes resulted in an increase in cellular ergosterol (Figure 5). Interestingly, those compounds that increase the respiration rate (AgClO₄ and [Ag(phendio)₂]ClO₄) also cause an increase in ergosterol content while those that depress respiration (Cu(ClO₄)₂·6H₂O and [Cu(phendio)₃](ClO₄)₂·4H₂O) lead to a reduction in ergosterol content.




Fig 3. Susceptibilities to Amphotericin B of *C. albicans* pre-grown in the presence of sub-MIC₉₀ levels of metal complexes. The susceptibility of *C. albicans* to amphotericin B was determined after cells had been first pre-grown in sub-MIC₉₀ levels of metal-based drug.




Fig 4. Respiration rates of *C. albicans* cells pre-grown in sub-MIC₉₀ levels of metal complexes. The oxygen consumption rates of *C. albicans* cell pre-grown in sub-MIC₉₀ levels of metal-based drug were determined using a Rank Oxygen electrode and expressed as μmoles of oxygen consumed per 10^8 cells per 30 sec.

NEW ARTWORK PLEASE!

Fig 5. Ergosterol profiles of *C. albicans* cells pre-grown in sub-MIC₉₀ levels of metal complexes. (Representative figure). The relative amounts of ergosterol in cells pre-treated with sub-MIC₉₀ concentrations of metal-based drug were ascertained spectrophotometrically over the range 250–300 nm.

Discussion

Metal-based drugs represent a novel group of antimicrobial agents with potential therapeutic applications (Coyle *et al.* 2003a; McCann *et al.* 2000). With the advent of fungal isolates manifesting resistance to azole and polyene drugs (Canuto & Rodero, 2002) there is a requirement for new drugs with alternative modes of action or with the ability to increase the efficacy of existing prescription drugs. Many metal-based drugs display modes of action distinct to those of the prescription anti-fungals (Coyle *et al.* 2003a) possibly allowing their use where resistance to conventional drugs has emerged (White *et al.* 1998). In addition, their different mode(s) of action may be utilised by employing such drugs in conjunction with existing drugs in order to target two (or more) sites within the fungal cell and thus raising the possibility of achieving the same therapeutic effect by reducing the required amount of an azole or polyene drug. Conventional anti-fungal drugs such as polyenes or azoles target ergosterol in the cell membrane or the ergosterol biosynthetic pathway, respectively. In the work presented here we have evaluated the ability of Cu(II) and Ag(I) phendio complexes to alter the susceptibility of *C. albicans* to conventional azole and polyene drugs.

The copper and silver complexes evaluated here demonstrated fungistatic properties but for this work sub-MIC₉₀ values were employed to ascertain their effect on the sensitivity of *C. albicans* to the conventional antifungal drugs. Sub-MIC levels were chosen

to ensure that the observed effect was not due to the fungistatic actions of the metal complex. When cells are pre-grown in sub-MIC₉₀ levels of the silver or copper complexes and then assessed for their response to miconazole or amphotericin B there is evidence of alterations in susceptibility. Pre-growth in 1/4 MIC₉₀ Cu(ClO₄)₂·6H₂O or [Cu(phendio)₃](ClO₄)₂·4H₂O increased the tolerance of *C. albicans* to miconazole and amphotericin B, whereas pre-growth in equivalent concentrations of AgClO₄ and [Ag(phendio)₂ClO₄] lowers the susceptibility of *C. albicans* to miconazole and amphotericin B.

The alteration in susceptibility to miconazole and amphotericin B may occur due to the fact that the metal complexes which inhibit respiration (i.e. Cu(ClO₄)₂·6H₂O and [Cu(phendio)₃](ClO₄)₂·4H₂O) also cause a reduction in the cellular ergosterol content. Azoles, such as miconazole, target lanosterol 14 α -demethylase which regulates an intermediate step in ergosterol biosynthesis (Morschhauser, 2002). The azoles can kill cells by inhibiting this enzyme which subsequently leads to a decrease in ergosterol content of the fungal cell membrane. In addition, the inhibition of the action of $\Delta^{5,6}$ -sterol desaturase by fluconazole leads to the accumulation of toxic intermediates (such as 14 α -methylfecosterol) which may prove fatal to the fungal cell (Geber *et al.* 1995). The altered susceptibility to miconazole may be due to reduced ergosterol biosynthesis resulting in fewer toxic side products being formed once the action of the demethylase has been inhibited by miconazole. Enhanced ergosterol biosynthesis may inadvertently lead to greater production of toxic side-products once the azole-mediated inhibition of sterol biosynthesis occurs.

Reduced levels of ergosterol in the fungal cell membrane provide fewer binding sites for amphotericin B. Consequently a higher concentration of amphotericin B is required to retard the growth of the cell. Reductions in sterol levels in *C. albicans* have been identified previously as a mechanism for increased growth in the presence of amphotericin B (Kelly *et al.* 1997; White *et al.* 1998). Disruptions of the genes in the ergosterol biosynthesis pathway cause a decrease in ergosterol in *C. glabrata* and an increase in drug tolerance, particularly to amphotericin B (Geber *et al.* 1995; Vazquez *et al.* 1996). Inhibition of respiration in *C. albicans* by erythromycin leads to a drop in ergosterol and a concomitant increase in tolerance to amphotericin B (Geraghty & Kavanagh, 2003a). In addition, disruption of mitochondrial function leads to

tolerance to amphotericin B due to depleted ergosterol synthesis (Geraghty & Kavanagh, 2003b).

The requirement for a functional mitochondrion in ergosterol biosynthesis is well characterised and arises from the provision of NADPH for squalene dimerisation (Parks & Casey, 1995). In addition, *Erg1* encodes squalene epoxidase, which converts squalene to 2,3-oxidosqualene. This is an oxygen-dependent step, and in a cell with reduced respiration there would be a consequent reduction in the synthesis of ergosterol (Daum *et al.* 1998), thus leading to the reduced ergosterol content evident in cells exposed to the copper complexes.

An increase in respiration, as evident when cells are pre-grown in the presence of AgClO_4 and $[\text{Ag}(\text{phendio})_2]\text{ClO}_4$ stimulates ergosterol production. Thus, there is more ergosterol for amphotericin B to bind and, as a consequence, lowering the amount of the polyene necessary for inhibition of cell growth. In the case of the altered sensitivity to miconazole, the silver complexes stimulate respiration and ergosterol biosynthesis and consequently may indirectly lead to the generation of more toxic side-products once the cells are exposed to miconazole.

The work presented here demonstrates that using sub-MIC levels of copper or silver dione complexes it is possible to alter the amount of an azole or polyene drug required to inhibit the growth of *C. albicans*. This effect appears to be mediated through alterations in the respiration rate of the cell, thereby influencing the amount of ergosterol synthesis. This opens the possibility of utilising, non-toxic levels of silver complexes to stimulate respiration and ergosterol production with a concomitant reduction in the amount of miconazole or amphotericin B required to achieve the relevant MIC_{90} .

Acknowledgement

This work was supported by a grant from the Department of Education, Libya.

References

Abu-Salah KM. 1996 Amphotericin B: an update. *Brit J Biomed Sci* **53**, 122–133.

- Arthington-Skaggs BA, Warnock DW, Morrison CJ. 1999 Quantitation of ergosterol content: Novel methods for determination of fluconazole susceptibility of *Candida albicans*. *J Clin Microbiol* **37**, 3332–3337.
- Canuto M, Rodero FG. 2002 Antifungal drug resistance to azole and polyenes. *Lancet (Infect Dis)* **2**, 550–562.
- Coyle B, Kavanagh K, McCann M, Devereux M, Geraghty M. 2003a Mode of anti-fungal activity of 1,10-phenanthroline and its Cu(II), Mn(II) and Ag(I) complexes. *Biometals* **16**, 321–329.
- Coyle B, Kinsella P, McCann M, Devereux M, O'Connor R, Clynes M, Kavanagh K. 2003b Induction of apoptosis in yeast and mammalian cells by exposure to 1,10-phenanthroline metal complexes. *Toxicology in Vitro* (in press).
- Daum G, Lees ND, Bard M, Dickson R. 1998 Biochemistry, cell biology and molecular biology of lipids of *Saccharomyces cerevisiae*. *Yeast* **14**, 1471–1510.
- Geber A, Hitchcock CA, Swartz JE, Pullen FS, Marsden KE, Kwon-Chung KJ, Bennett JE. 1995 Deletion of the *Candida glabrata* *ERG3* and *ERG11* genes: Effect on cell viability, cell growth, sterol composition and antifungal susceptibility. *Antimicrob Agen Chemother* **39**, 2708–2717.
- Geraghty P, Kavanagh K. 2003a Erythromycin, an inhibitor of mitochondrial protein biosynthesis, alters the Amphotericin B susceptibility of *Candida albicans*. *J Pharm Pharmacol* **55**, 179–184.
- Geraghty P, Kavanagh K. 2003b Disruption of mitochondrial function in *Candida albicans* leads to reduced cellular ergosterol levels and elevated growth in the presence of Amphotericin B. *Arch Microbiol* **179**, 295–300.
- Igdaloff D, Santi DV, Eckert TS, Bruice TC. 1983. Effects of 1,7- and 1,10-phenanthroline dione on tissue culture cells. *Biochemical Pharmacol* **32**, 172–174.
- Kelly SI, Lamb DC, Kelly DE, Manning NJ, Loeffler J, Herbart H, Schumacher U, Einsele H. 1997 Resistance to fluconazole and cross resistance to amphotericin B in *Candida albicans* from AIDS patients caused by defective sterol $\Delta^{5,6}$ desaturation. *FEBS Letts* **400**, 80–82.
- McCann M, Geraghty M, Devereux M *et al.* 2000 Insights into the mode of action of the anti-*Candida* activity of 1,10-phenanthroline and its metal chelates. *Metal-Based Drugs* **7**, 185–193.
- McCann M, Coyle B, McKay S, McCormack P, Kavanagh K, Devereux M, McKee V, Kinsella P, O'Connor R, Clynes M. Synthesis and X-ray crystal structure of $[\text{Ag}(\text{phendio})_2]\text{ClO}_4$ (phendio = 1,10-phenanthroline-5,6-dione) and its effects on fungal and mammalian cells. *Biometals* (submitted).
- Morschhauser J. 2002 The genetic basis of fluconazole resistance development in *Candida albicans*. *Biochem Biophys Acta* **1587**, 240–248.
- Parks LW, Casey WM. 1995 Physiological implications of sterol biosynthesis in yeast. *Ann Rev Microbiol* **49**, 95–116.
- Vazquez JA, Arganoza MT, Vaishampayan JK, Akins RA. 1996 *In vitro* interaction between Amphotericin B and azoles in *Candida albicans*. *Antimicrob Agen Chemother* **40**: 2511–2516.
- White TC, Marr KA, Bowden RA. 1998 Clinical, cellular and molecular factors that contribute to anti-fungal drug resistance. *Clin Microbiol Rev* **11**: 382–402.