

DEUTSCHES ELEKTRONEN – SYNCHROTRON

DESY 93-066
ITP-UH-05/93
May 1993



**A Geometrical Interpretation of
Renormalisation Group Flow**

B. P. Dolan

Institut für Theoretische Physik, Universität Hannover

ISSN 0418-9833

NOTKESTRASSE 85 · D - 2000 HAMBURG 52

DESY behält sich alle Rechte für den Fall der Schutzrechtserteilung und für die wirtschaftliche Verwertung der in diesem Bericht enthaltenen Informationen vor.

DESY reserves all rights for commercial use of information included in this report, especially in case of filing application for or grant of patents.

To be sure that your preprints are promptly included in the
HIGH ENERGY PHYSICS INDEX,
send them to (if possible by air mail):

DESY
Bibliothek
Notkestraße 85
W-2000 Hamburg 52
Germany

DESY-IfH
Bibliothek
Platanenallee 6
0-1615 Zeuthen
Germany

§1 Introduction

Geometry and physics seem inextricably linked in our attempts to understand the fundamental forces of nature, not only in Einstein's geometrical interpretation of the gravitational force but also in the geometrical approach to the gauge interactions of particle physics which has introduced concepts such as fibre bundles and connections into elementary particle theory, together with their associated structures of instantons and monopoles. Many aspects of modern physics have a geometrical interpretation and can be described in a co-ordinate free manner. In conjunction with this emphasis on geometry is the importance of symmetry, which has played an ever increasing rôle in twentieth century physics. Symmetries in particle physics can be classified into two types - internal and space symmetries. If the dynamics of a physical system are determined by an action which has a symmetry then one can immediately deduce the existence of quantities that remain conserved under all possible circumstances. This concept, however, is laced with subtleties as it sometimes happens that a symmetry which is present in the classical action is violated by quantum processes and an anomaly ensues.

The renormalisation group (RG) equation was first introduced into particle physics with the observation that quantum effects can violate an important space-time symmetry of classical electromagnetism, that of conformal invariance. This is not a symmetry of the Minkowski metric, it only preserves angles not lengths, but it is a symmetry of the classical action for electromagnetism, in the energy regime $E > 0.5 MeV$ where the electron mass can be ignored. It was observed in [1] and [2] that the electromagnetic charge picks up a scale dependence through quantum effects and that the derivative of the electromagnetic coupling with respect to the scale should be an analytic function of the coupling itself, the β -function. More important than the couplings, however, are the physical amplitudes in a quantum field theory. Amplitudes depend not only on the couplings but also on a number of points in space (Minkowski or Euclidean) which must be labelled by some co-ordinates, x_i , (usually Cartesian in flat space). It was pointed out by Callan and Symanzik [3] that the variation in the couplings under a change in the scale at which they are defined can always be compensated for by a change in the separation of the points, or alternatively a rescaling of the co-ordinates x_i , so that the vacuum amplitudes remain invariant. This results in an inhomogeneous partial differential equation for the amplitudes - the RG equation. This concept was further extended to a homogeneous equation by 'tHooft and Weinberg [4]. Meanwhile, the RG equation was raised to central importance in statistical physics through the work of Wilson [5].

The idea that I wish to present in this paper is the following. A rescaling of points in Euclidean space, \mathbf{R}^D , is a diffeomorphism and is generated by the vector $\vec{D} = x^\mu \partial_\mu$. If the space of couplings is also considered to be a finite dimensional, differentiable manifold, \mathcal{G} , then the amplitudes can be interpreted as co-variant tensors on \mathcal{G} and one can interpret the rescaling of the couplings as a diffeomorphism of \mathcal{G} , and it is generated by a vector which is given by the β -functions of the theory. The renormalisation group equation then expresses the fact that the change in amplitudes under the diffeomorphism of \mathbf{R}^D generated by \vec{D} , keeping the couplings fixed, is exactly the same as the change effected by a diffeomorphism of \mathcal{G} , generated by the vector $\vec{\beta}$, keeping the spatial points x_i fixed. This will be proven in

A Geometrical Interpretation Of Renormalisation Group Flow *

Brian P. Dolan

*Institut für Theoretische Physik ****Universität Hannover
Hannover, Germany*

e-mail: bdolan@kantor.itp.uni-hannover.de

ABSTRACT

The renormalisation group (RG) equation in D -dimensional Euclidean space, \mathbf{R}^D , is analysed from a geometrical point of view. A general form of the RG equation is derived which is applicable to composite operators as well tensor operators (on \mathbf{R}^D) which may depend on the Euclidean metric. It is argued that physical N -point amplitudes should be interpreted as rank N co-variant tensors on the space of couplings, \mathcal{G} , and that the RG equation can be viewed as an equation for Lie transport on \mathcal{G} with respect to the vector field generated by the β -functions of the theory. In one sense it is nothing more than the definition of a Lie derivative. The source of the anomalous dimensions can be interpreted as being due to the change of the basis vectors on \mathcal{G} under Lie transport. The RG equation acts as a bridge between Euclidean space and coupling constant space in that the effect on amplitudes of a diffeomorphism of \mathbf{R}^D (that of dilations) is completely equivalent to a diffeomorphism of \mathcal{G} generated by the β -functions of the theory. A form of the RG equation for operators is also given. These ideas are developed in detail for the example of massive $\lambda\phi^4$ theory in 4 dimensions.

PACS Nos. 03.70.+k and 11.10.Gh

* Work carried out with an Alexander von Humboldt research stipendium.

** On leave of absence from: Department of Mathematical Physics, St. Patrick's College, Maynooth, Ireland.

§2 Conformal Transformations And The Renormalisation Group Equation

This section is based on the introductory part of [7]. Consider a field theory in flat D -dimensional Euclidean space. In principle there are an infinite number of operators that can be constructed out of the fields, each of which introduces a coupling, but the criterion of renormalisability requires that only a finite number of these couplings is independent, [5]. This means that, within the a priori infinite dimensional space of coupling constants, the theory can be formulated on a finite n -dimensional subspace which will be denoted by \mathcal{G} . It will be assumed that \mathcal{G} is, at least locally, a differentiable manifold and we will denote the renormalised couplings (co-ordinates on \mathcal{G}) by $g^a, a = 1, \dots, n$. Bare quantities will be represented by a subscript o . Thus the bare couplings are denoted by g^o . It will be assumed that a regularisation procedure is imposed which renders bare quantities very large but still finite. This is because bare quantities appear in some of the following formulae and we do not want to be manipulating infinite quantities. Ultimately, of course, all bare quantities disappear from physical amplitudes. The couplings g^a will be taken to be real and massless. If the theory contains any couplings which are massive these can always be made massless by multiplying by appropriate powers of the renormalisation mass scale, κ . Questions about the global structure of \mathcal{G} will not be addressed here.

The physics is given by the renormalised N -point vacuum amplitudes (or correlation functions in statistical physics)

$$A^{(N)}(x_1, \dots, x_N) = \langle \hat{A}_1(x_1) \dots \hat{A}_N(x_N) \rangle, \quad (1)$$

where $x_i, i = 1, \dots, N$ are positions in \mathbf{R}^D with $x_i \neq x_j, \forall i \neq j$ and \hat{A}_i are physical operators. In many of the following expressions we shall freely interchange between operator and functional integral formalism. The former will be represented as \hat{A}_i , whereas the corresponding quantity appearing in functional integrals, which can be thought of as classical functions of the fields, will be denoted by A_i without a circumflex. The amplitudes (1) can be expressed in terms of the couplings g^a , but their form depends on the choice of couplings and the renormalisation scheme chosen and does not necessarily bear any simple relation to the g^a . For example in scalar $\lambda\phi^4$ theory in four space dimensions the relation between the coupling, $\lambda(\kappa)$, at the renormalisation point, κ , and the four point amplitude, in momentum space at the symmetric point $p_1 p_2 p_3 p_4 = \frac{\kappa^2}{4}(\delta_{\mu\nu} - 1)$; depends on the renormalisation scheme. Using momentum subtraction they are the same (because they are defined to be, for $p^2 = \kappa^2$ but not otherwise) but using dimensional regularisation and minimal subtraction they are not the same. Even at $p^2 = \kappa^2$ the four point amplitude $\langle \hat{\phi}(p_1) \hat{\phi}(p_2) \hat{\phi}(p_3) \hat{\phi}(p_4) \rangle$ is a complicated, though analytic, function of $\lambda(\kappa)$ which is only approximately equal to λ for small λ . From this it is clear that the specific choice of co-ordinates on \mathcal{G} is not physically important (though, in practice, a clever choice makes calculations much simpler).

The theory can be probed by distorting it slightly to see how it responds. Let \bar{X} be an arbitrary vector field on \mathbf{R}^D (generating a diffeomorphism of space). For general \bar{X} the amplitudes will change under this diffeomorphism, and the response is described by the insertion of the stress operator, $\hat{T}_{\mu\nu}(x)$, into the amplitudes (in Lorentzian signature this would, of course, be called the energy-momentum tensor). Let us see how this comes

§3. Alternatively, if the RG equation is written as a differential equation involving $\kappa \frac{\partial}{\partial \kappa} |g$ rather than $x^a \frac{\partial}{\partial x^a} |g$, using standard naive scaling arguments, it becomes nothing more than the definition of a Lie derivative on \mathcal{G} with respect to the vector field $\bar{\beta}$. The terms in the RG equation involving anomalous dimensions are interpreted as coming from the change in the basis for co-vectors, dg^a , as we move along the RG trajectory.

Of course quantum field theory is famous for being plagued by "infinities" which, at least for renormalised theories, can be "tamed" by a regularisation procedure. This requires the introduction of "bare" couplings, $g^o(g, \epsilon)$, which are analytic functions of the renormalised couplings, g^a , and a regularisation parameter or parameter, ϵ , e.g. for a cut-off, $\Lambda, \epsilon = \kappa/\Lambda$ where κ is a renormalisation point and for dimensional regularisation $\epsilon = 4 - D$ where D is the dimension of space or space-time. It will further be argued in §3 that g^o and g^a can be thought of as different co-ordinate systems on \mathcal{G} . The matrix $\frac{\partial g^o}{\partial g^a}$ tells us how to transform tensors (amplitudes) between co-ordinate systems. g^o enter on a different footing from g^a however in that they depend on the regularisation parameter whereas g^a do not. The "infinities" of quantum field theory are then viewed as being due to the fact that the co-ordinate transformation between g^o and g^a is singular when the regularisation parameter is removed. This is not a disaster, singularities in co-ordinate transformations are common in differential geometry. For example the event horizon of a black hole in the Schwarzschild metric was for a long time viewed as being "singular" because, in polar co-ordinates, the metric appears to be degenerate there. Then a co-ordinate system was discovered, Kruskal co-ordinates, in which the metric is perfectly regular at the event horizon. The transformation between polar co-ordinates and Kruskal co-ordinates in the Schwarzschild metric is singular at the event horizon but the metric, and physics, is perfectly regular there. The apparent singularity in polars is just due to a bad choice of co-ordinates. Of course polar co-ordinates are fine for describing space outside of the event horizon. The singularity at the origin, however, is a real physical singularity of space-time, indicating that some new physics must enter here. One must be very careful to distinguish between genuine singularities and singularities that are merely due to the choice of co-ordinates.

The basic idea presented here, that the RG equation should be viewed in terms of a co-ordinate transformation on the space of couplings, was inspired by O'Connor and Stephens [6].

The main results will be described in §3 where it is shown that the RG equation for vacuum amplitudes can be interpreted as a Lie derivative on the space of couplings and an expression for the RG equation for operators is also presented. As a preliminary, §2 gives a discussion of the rôle of the stress tensor in the RG equation and a version of the RG equation is derived by demanding invariance of functional integrals under a specific change in integration variable, associated with dilations of Euclidean space. §4 illustrates the ideas with a detailed analysis of massive $\lambda\phi^4$ with a constant source, where the space \mathcal{G} is three dimensional. Finally §5 gives a summary and outlook.

about. Express the amplitudes in the form of functional integrals over the bare fields of the theory, which will be denoted by $\varphi_o(x)$ but need not be scalars,

$$A^{(N)}(x_1, \dots, x_N) = \int \rho[\varphi_o] A_1(x_1) \dots A_N(x_N) e^{-S_o[g_o, \varphi_o, \partial\varphi_o, \gamma]} / Z_o[g_o, \gamma]. \quad (2)$$

Here $\gamma_{\mu\nu}$ is the Euclidean metric on \mathbf{R}^D and $A_i(x_i)$ depend on the renormalised fields $\varphi(x_i)$ and possibly also on $\gamma_{\mu\nu}$, e.g. if $A_i(x_i)$ involves contractions of $\partial_\mu\varphi$ or if φ is a vector field and $A_i(x_i)$ are composite operators. $\rho[\varphi_o]$ is the measure. The notation $\rho[\varphi_o]$ is used purely for convenience, to simplify subsequent formulae, and is not intended to imply that the measure is anything other than the usual one, more conventionally written as $[d\varphi_o]$ or $\mathcal{D}\varphi_o$. $Z_o[g_o, \gamma]$ is the partition function,

$$Z_o[g_o, \epsilon, \gamma] = Z_o[g_o, \gamma] = \int \rho[\varphi_o] e^{-S_o[g_o, \varphi_o, \partial\varphi_o, \gamma]}, \quad (3)$$

and the action is the Hamiltonian density integrated over \mathbf{R}^D ,

$$S_o[g_o, \epsilon, \varphi, \partial\varphi, \gamma] = S_o[g_o, \varphi_o, \partial\varphi_o, \gamma] = \int d^D\tilde{y} H_o(g_o, \varphi_o(y), \partial\varphi_o(y), \gamma(y)), \quad (4)$$

where $d^D\tilde{y} = \sqrt{\gamma} d^Dy$ is the measure in D -dimensional Euclidean space. The subscript o here is to emphasise that it is the bare action that appears exponentiated in functional integrals. When doing perturbation theory this is split into a renormalised action plus counterterms and is then considered to be a function of renormalised fields and couplings but this split has no physical significance, just as the renormalised couplings have no intrinsic physical meaning. The action can then be thought as a function of the renormalised couplings, as indicated in equation (4), and the regularisation parameter(s) - the latter dependence being purely through the counterterms. The bare subscript on the partition function (3) could be removed by using a measure defined in terms of the renormalised fields rather than the bare fields,

$$Z[g_o, \epsilon, \gamma] = \int \rho[\varphi] e^{-S_o[g_o, \epsilon, \varphi, \partial\varphi, \gamma]}, \quad (5)$$

thus introducing a factor which cancels in all amplitudes.

Now consider the response of the amplitudes to a deformation generated by a vector field, \vec{X} , on \mathbf{R}^D

$$\mathcal{L}_{\vec{X}} A^{(N)}(x_1, \dots, x_N) = \sum_{i=1}^N \int \rho[\varphi_o] A_1(x_1) \dots \{ \mathcal{L}_{\vec{X}} A_i(x_i) \} \dots A_N(x_N) e^{-S_o} / Z_o, \quad (6)$$

where $\mathcal{L}_{\vec{X}}$ represents Lie differentiation with respect to \vec{X} . The partition function itself is unchanged by this deformation. This is because $\mathcal{L}_{\vec{X}} S_o$ vanishes since the action is integrated over \mathbf{R}^D and the fields are assumed to vanish at infinity.

We shall use two identities to re-express the right hand side of equation (6). Firstly

$$\mathcal{L}_{\vec{X}} A_i = \frac{\delta A_i}{\delta\varphi} \mathcal{L}_{\vec{X}}\varphi + \frac{\delta A_i}{\delta(\partial_\mu\varphi)} \mathcal{L}_{\vec{X}}(\partial_\mu\varphi) + \frac{\delta A_i}{\delta\gamma_{\mu\nu}} (\mathcal{L}_{\vec{X}}\gamma)^{\mu\nu}, \quad (7)$$

which is just Leibniz rule for differentiation allowing for the possibility that $A_i(x_i)$ might depend on $\partial_\mu\varphi(x_i)$ as well as $\varphi(x_i)$ and may have explicit metric dependence. The Lie derivative of the inverse metric is given by the usual formula,

$$(\mathcal{L}_{\vec{X}}\gamma)^{\mu\nu} = X^\tau \partial_\tau \gamma^{\mu\nu} - \gamma^{\mu\tau} \partial_\tau X^\nu - \gamma^{\tau\nu} \partial_\tau X^\mu. \quad (8)$$

In Cartesian co-ordinates, with $\gamma_{\mu\nu} = \delta_{\mu\nu}$, this reduces to the familiar deformation

$$(\mathcal{L}_{\vec{X}}\gamma)^{\mu\nu} = -\partial^\mu X^\nu - \partial^\nu X^\mu. \quad (9)$$

The second identity follows from the fact that functional integrals should be invariant under reparameterisations of the integration variables, φ_o , so that

$$\begin{aligned} \int \rho[\varphi_o] \int d^D\tilde{y} A_1(x_1) \dots A_N(x_N) \left\{ \frac{\delta S_o}{\delta\varphi_o(y)} \delta\varphi_o(y) + \frac{\delta S_o}{\delta(\partial_\mu\varphi_o(y))} \delta(\partial_\mu\varphi_o(y)) \right\} \\ = \sum_{i=1}^N \int \rho[\varphi_o] \int d^D\tilde{y} A_1(x_1) \dots \left\{ \frac{\delta A_i(x_i)}{\delta\varphi_o(y)} \delta\varphi_o(y) + \frac{\delta A_i(x_i)}{\delta(\partial_\mu\varphi_o(y))} \delta(\partial_\mu\varphi_o(y)) \right\} \dots A_N(x_N) e^{-S_o}, \end{aligned} \quad (10)$$

where we have allowed for the possibility that the measure might not be invariant. Clearly the bare field φ_o in the curly brackets on the right hand side of this equation can be replaced by the renormalised field φ since the renormalisation co-efficients just cancel. Applying this identity to the variation $\delta\varphi_o = \epsilon \mathcal{L}_{\vec{X}}\varphi_o$, with ϵ independent of position, and using the Leibniz rule in the form of equation (7) to replace all functional variations with respect to fields with Lie derivatives and metric variations gives

$$\begin{aligned} \int \rho[\varphi_o] \int d^D\tilde{y} A_1(x_1) \dots A_N(x_N) \left\{ \mathcal{L}_{\vec{X}} H_o - \left(\frac{\delta H_o}{\delta\gamma^{\mu\nu}} - \frac{1}{2} M_{\mu\nu} \right) (\mathcal{L}_{\vec{X}}\gamma)^{\mu\nu} \right\} \Big|_y e^{-S_o} \\ = \sum_{i=1}^N \int \rho[\varphi_o] A_1(x_1) \dots \left\{ \mathcal{L}_{\vec{X}} A_i - \frac{\delta A_i}{\delta\gamma^{\mu\nu}} (\mathcal{L}_{\vec{X}}\gamma)^{\mu\nu} \right\} \Big|_{x_i} \dots A_N(x_N) e^{-S_o}, \end{aligned} \quad (11)$$

where we have defined

$$\frac{1}{2} M_{\mu\nu} (\mathcal{L}_{\vec{X}}\gamma)^{\mu\nu} = -\rho^{-1} \frac{\delta\rho}{\delta\varphi_o} \mathcal{L}_{\vec{X}}\varphi_o(y).$$

Since $\delta\varphi_o = \epsilon \mathcal{L}_{\vec{X}}\varphi_o$ is linear in φ_o and $\frac{1}{2}(M_{\mu\nu} \mathcal{L}_{\vec{X}}\gamma)^{\mu\nu}$ is just the Jacobian of this transformation, one can expect that $M_{\mu\nu}$ is independent of φ_o , i.e. $M_{\mu\nu}$ is a multiple of the identity.

One may worry here about what happens when φ is a spinor field because it is difficult to find a consistent definition of the Lie derivative of a spinor field for an arbitrary diffeomorphism, \bar{X} . However in this section we will only really be interested in the case where \bar{X} is a conformal Killing vector for the Euclidean metric, $\gamma_{\mu\nu}$, and the Lie derivative of a spinor with respect to a conformal Killing vector can be well defined*.

The classical stress tensor (density) is defined as

$$T_{\mu\nu} = 2 \frac{\delta H_0}{\delta \gamma^{\mu\nu}} \quad (12)$$

and this is just the second term on the left hand side of equation (11). The reason why there is no bare subscript on $T_{\mu\nu}$ will be explained shortly. The first term in (11) can be dropped because it is a total derivative. To see this note that since H_0 is a D -form (a density) we have

$$\mathcal{L}_{\bar{X}} H_0 = i_{\bar{X}} dH_0 + d i_{\bar{X}} H_0 = d i_{\bar{X}} H_0 \quad (13)$$

and hence is a total derivative. Now dividing (11) by $Z_0[g, \epsilon, \gamma]$ in order to get genuine amplitudes gives

$$\begin{aligned} \mathcal{L}_{\bar{X}} A^{(N)}(x_1, \dots, x_N) &= -\frac{1}{2} \int d^D \tilde{y} (\mathcal{L}_{\bar{X}} \gamma)^{\mu\nu}(y) \langle (\hat{T}_{\mu\nu}(y) - \hat{M}_{\mu\nu}(y)) \hat{A}_1(x_1) \dots \hat{A}_N(x_N) \rangle \\ &+ \sum_{i=1}^N \langle \hat{A}_1(x_1) \dots \left[\frac{\delta \hat{A}_i}{\delta \gamma^{\mu\nu}} (\mathcal{L}_{\bar{X}} \gamma)^{\mu\nu} \right]_{x_i} \dots \hat{A}_N(x_N) \rangle. \end{aligned} \quad (14)$$

When $N = 0$ this gives

$$\int d^D \tilde{y} (\mathcal{L}_{\bar{X}} \gamma)^{\mu\nu}(y) \langle \hat{T}_{\mu\nu}(y) \rangle = -M_{\mu\nu}(y) = 0, \quad (15)$$

i.e. up to total derivatives we have that $\hat{M}_{\mu\nu}$ is just the expectation value of $\langle \hat{T}_{\mu\nu} \rangle$. An integration by parts can be performed in this equation and using equation (9) with an arbitrary diffeomorphism \bar{X} we deduce that, in Cartesians,

$$\langle \partial_\mu \hat{T}^{\mu\nu}(y) \rangle = 0, \quad (16)$$

where we have defined $\hat{T}^{\mu\nu} = \hat{T}_{\mu\nu} - \hat{M}_{\mu\nu}$. Then (14) can be written

$$\begin{aligned} \mathcal{L}_{\bar{X}} A^{(N)}(x_1, \dots, x_N) &= -\frac{1}{2} \int d^D \tilde{y} (\mathcal{L}_{\bar{X}} \gamma)^{\mu\nu}(y) \langle \hat{T}^{\mu\nu}(y) \hat{A}_1(x_1) \dots \hat{A}_N(x_N) \rangle \\ &+ \sum_{i=1}^N \langle \hat{A}_1(x_1) \dots \left[\frac{\delta \hat{A}_i}{\delta \gamma^{\mu\nu}} (\mathcal{L}_{\bar{X}} \gamma)^{\mu\nu} \right]_{x_i} \dots \hat{A}_N(x_N) \rangle. \end{aligned} \quad (17)$$

* I am grateful to Norbert Dragon for an illuminating discussion on this point.

Note that if the \hat{A}_i are composite operators, e.g. $\hat{A}_1 = \partial_\mu \hat{\varphi} \partial_\nu \hat{\varphi} \gamma^{\mu\nu}$, the terms $\frac{\delta \hat{A}_i}{\delta \gamma^{\mu\nu}} |_{x_i}$ can give a non-zero contribution to (17).

Equation (17) can be interpreted as a quantum mechanical statement of translational invariance and is often called the Ward identity associated with this symmetry. It is one half of the renormalisation group equation and will be used extensively later on. For the moment a short digression will be made on the properties of the quantum stress operator, $\hat{T}^{\mu\nu}$.

Classically translation invariance of the Euclidean metric ensures, via Noether's theorem, the existence of D conserved currents, when the equations of motion are used. In Cartesian co-ordinates,

$$\partial_\mu T^{C\mu}_\nu = 0, \quad \nu = 1, \dots, D. \quad (18)$$

Quantum mechanically, we must take the existence of a stress operator obeying equation (17) as an *assumption*, [8]. The stress operator, $\hat{T}^{\mu\nu}$, appearing in (17) must be the regularised stress operator. It satisfies the quantum version of (18),

$$\partial_\mu \hat{T}^{\mu\nu} = 0 \quad (19)$$

(modulo the equations of motion). It is defined (in flat space) so that $\langle \hat{T}^{\mu\nu} \rangle = 0$.

In quantum field theory conserved currents do not get renormalised. This means that for any global symmetry of the action, which does not acquire an anomaly upon quantisation, the bare conserved current \hat{j}^μ is equal to the renormalised current \hat{J}^μ . In other words the renormalisation constant Z_J , which is defined by

$$\hat{J}^\mu = Z_J \hat{j}^\mu, \quad (20)$$

is unity (rotational invariance demands that Z_J be the same for every component of the vector J^μ). To see this in the case of translational invariance consider equation (17), which is the mathematical expression of this symmetry. Assuming translational invariance to hold at both the bare and the renormalised level, this equation must be true for both bare and renormalised operators. Consider equation (17) for bare operators, \hat{A}_{oi} , which are independent of the metric (including metric dependence does not affect the argument)

$$\begin{aligned} \sum_{i=1}^n \langle \hat{A}_{oi}(x_1) \dots (\mathcal{L}_{\bar{X}} \hat{A}_{oi}(x_i)) \dots \hat{A}_{oN}(x_N) \rangle \\ = -\frac{1}{2} \int d^D \tilde{y} (\mathcal{L}_{\bar{X}} \gamma)^{\mu\nu}(y) \langle \hat{T}^{\mu\nu}_{oi}(y) \hat{A}_{oi}(x_1) \dots \hat{A}_{oN}(x_n) \rangle \end{aligned} \quad (21)$$

\Leftrightarrow

$$\begin{aligned} \left(\prod_{i=1}^N Z_{A_i} \right) \sum_{i=1}^n \langle \hat{A}_1(x_1) \dots (\mathcal{L}_{\bar{X}} \hat{A}_i(x_i)) \dots \hat{A}_N(x_N) \rangle \\ = -\frac{1}{2} \left(\prod_{i=1}^N Z_{A_i} \right) Z_T \int d^D \tilde{y} (\mathcal{L}_{\bar{X}} \gamma)^{\mu\nu}(y) \langle \hat{T}^{\mu\nu}(y) \hat{A}_1(x_1) \dots \hat{A}_N(x_n) \rangle, \end{aligned} \quad (22)$$

where Z_T is the renormalisation constant for the stress tensor, $\hat{T}_{\mu\nu} = Z_T \hat{T}_{\mu\nu}$, and Z_{A_i} are the renormalisation constants for the operators \hat{A}_i , $\hat{A}_{\alpha i} = Z_{A_i} \hat{A}_i$ (for simplicity it is assumed that there is no operator mixing between the \hat{A}_i 's). Putting (17) for the renormalised operators into equation (22) gives the non-renormalisation theorem $Z_T = 1$. Thus we have, as a consequence of the assumption that translational invariance holds at both the bare and renormalised level,

$$\hat{T}_{\mu\nu} = \hat{T}_{\mu\nu} \quad (23)$$

(modulo the equations of motion). This is why there is no bare subscript on the left hand side of equation (12).

Equation (23) is rather subtle in practice for (at least) two reasons. Firstly, for example in scalar $\lambda\varphi^4$ theory in four dimensions, the "naive" stress tensor is not a finite operator and one must construct an "improved" stress tensor [9]. Equation (23) can be demonstrated explicitly for scalar fields, see §4. For gauge theories the question of the existence of a finite stress operator is addressed in refs. [10] and [11] where it is argued that no improvement terms, other than those already necessary for scalar fields, are required. This is connected with the classical conformal invariance of gauge field theories. A second subtlety associated with (23) is that it assumes translational invariance at the bare level,

$$\partial_\mu \hat{T}_{\nu\sigma} = 0, \quad (24)$$

i.e. bare quantities are translationally invariant. This means that whatever regularisation procedure is adopted it must respect translational invariance, [9]. In particular if we use (23) then we cannot use a cut-off! Fortunately there exist regularisation procedures that preserve translational invariance, e.g. Pauli-Villars or dimensional regularisation. Nevertheless this is a rather unsatisfactory aspect of the analysis presented here. The philosophy of this paper is to try to develop a description of the renormalisation group in a co-ordinate independent manner, i.e. in a way that is independent of the regularisation scheme chosen, yet here we find ourselves immediately restricting our choice of schemes. Of course this only an intermediate step and in the end all physical amplitudes are independent of bare quantities. However, the results of §3 do not rely on the stress tensor as such, so this is not an important restriction.

Now let us return to the Ward identity (17). The real power of this equation lies in choosing particular diffeomorphisms, \vec{X} , which are related to the symmetries of the Euclidean metric γ . In particular, we can obtain very powerful results by taking \vec{X} to be the generators of the conformal group, $SO(1, D+1)$, in D Euclidean dimensions. Thus, for isometries (translations or rotations) of γ , the right hand side of (17) vanishes, and this equation tells us that all physical amplitudes are translationally and rotationally invariant.

The response of the system to changes in scale is given by taking \vec{X} to be the vector which generates dilations. In Cartesian this is

$$\vec{X} = \vec{D} = x^\mu \partial_\mu \quad \text{with} \quad (\hat{\mathcal{L}}_{\vec{D}} \gamma)^{\mu\nu} = -2\gamma^{\mu\nu}. \quad (25)$$

In this case equation (17) reduces to

$$\hat{\mathcal{L}}_{\vec{D}} A^{(N)}(x_1, \dots, x_N) = \int d^D y < \hat{T}'(y) \hat{A}_1(x_1) \dots \hat{A}_N(x_N) > - \sum_{i=1}^N < \hat{A}_1(x_1) \dots \{\Delta_\gamma \hat{A}_i(x_i)\} \dots \hat{A}_N(x_N) >, \quad (26)$$

where $\hat{T}' = \hat{T}'^\mu{}_\mu$ is the trace of the stress operator and $\Delta_\gamma \hat{A}_i = 2\gamma^{\mu\nu} \left(\frac{\delta \hat{A}_i}{\delta \gamma^{\mu\nu}} \right)$.

Now we shall eliminate \hat{T}' to obtain the renormalisation group equation. Following ref. [7] we consider the operators

$$\hat{\Phi}_\alpha(x) = \frac{\partial \hat{H}_\alpha(x)}{\partial \varphi^\alpha}, \quad (27)$$

where $\hat{H}_\alpha(x)$ is the bare Hamiltonian density, to be a basis for all relevant or marginal operators of the theory, i.e. any relevant or marginal operator, which is a scalar (or more precisely a density) in \mathbf{R}^D , can be written as a linear combination of $\hat{\Phi}_\alpha(x)$. There is one other operator which must be included in order to have a complete basis and this is an operator proportional to the equations of motion of the theory since this linear combination is obtained not by varying a coupling in the action but by varying the fields. Let

$$\hat{E}_\alpha(x) = \varphi_\alpha(x) \left\{ \frac{\delta S_0}{\delta \varphi_\alpha(x)} - \frac{\partial}{\partial x^\mu} \left(\frac{\delta S_0}{\delta (\partial_\mu \varphi_\alpha(x))} \right) \right\} = \varphi(x) \left\{ \frac{\delta S_0}{\delta \varphi(x)} - \frac{\partial}{\partial x^\mu} \left(\frac{\delta S_0}{\delta (\partial_\mu \varphi(x))} \right) \right\} \quad (28)$$

then the operator $\hat{E}_\alpha(x)$ has canonical mass dimension D . When inserted into amplitudes $A^{(N)}(x_1, \dots, x_N)$ it gives

$$< \hat{E}'_0(x) \hat{A}_1(x_1) \dots \hat{A}_N(x_N) > = \sum_{i=1}^N < \hat{A}_1(x_1) \dots \{\Delta_{\varphi(x)} \hat{A}_i(x_i)\} \dots \hat{A}_N(x_N) >, \quad (29)$$

where we have defined

$$\Delta_{\varphi(x)} \hat{A}_i(x_i) = \varphi(x) \left\{ \frac{\delta \hat{A}_i(x_i)}{\delta (\varphi(x))} - \partial_\mu \left(\frac{\delta \hat{A}_i(x_i)}{\delta (\partial_\mu \varphi(x))} \right) \right\}. \quad (30)$$

Here $\hat{E}'_0(x) = \hat{E}_0(x) - < \hat{E}_0(x) >$ where $< \hat{E}_0(x) >$ can be obtained by using (10) with $N = 0$,

$$< \hat{E}_0(x) > = < \rho^{-1} \frac{\delta \rho}{\delta \varphi_0(x)} \varphi_0(x) >. \quad (31)$$

For general N equation (29) is just equation (10) with $\delta \varphi_0(y) = \epsilon \delta^{(N)}(x-y) \varphi_0(x)$, ϵ a constant, and integrated by parts using Cartesian co-ordinates.

The right hand side of (29) vanishes unless $x = x_i$ for some i . In particular if $\hat{A}_i(x_i) = \hat{\varphi}(x_i)$ then

$$\langle \hat{E}'_i(x) \hat{\varphi}(x_1) \cdots \hat{\varphi}(x_N) \rangle = \sum_{i=1}^N \delta^{(N)}(x - x_i) \langle \hat{\varphi}(x_1) \cdots \hat{\varphi}(x_N) \rangle. \quad (32)$$

In the ensuing analysis equation (29) will only ever appear integrated over x so it will be convenient to define

$$\Delta_\varphi A_i(x_i) = \int d^D \tilde{x} \Delta_\varphi(x) A_i(x_i) = \int d^D \tilde{x} \varphi(x) \left\{ \frac{\delta A_i(x_i)}{\delta(\varphi(x))} - \partial_\mu \left(\frac{\delta A_i(x_i)}{\delta(\partial_\mu \varphi(x))} \right) \right\}. \quad (33)$$

Note that applying the same arguments to (29) as were used on the Ward identity (17) to derive the non-renormalisation theorem (23) shows that $\hat{E}'_i(x) = \hat{E}(x)$, i.e. the "equation of motion operator" does not get renormalised, [12]. This is analogous to a Ward identity in that the equations of motion themselves constitute a linear relation between operators which must hold at both the bare and renormalised level and thus prevent $\hat{E}(x)$ from getting renormalised. However this linear relation does not represent a symmetry of the action and it would therefore be an abuse of language to call it a Ward identity (unless one wants to promote reparameterisation invariance of functional integrals, equation (10), to a symmetry of the theory).

To incorporate tensor (or spinor) operators would require a larger basis than $\hat{\Phi}_a$ but this will not be necessary here. A consequence of the definition of $\hat{\Phi}_a(x)$ is

$$\partial_a \hat{\Phi}_b = \partial_b \hat{\Phi}_a. \quad (34)$$

The set of operators $\hat{\Phi}_a$ can be thought of as forming an operator valued one-form, $\hat{\Phi} = \hat{\Phi}_a d\theta^a = d\hat{H}_a$, in the co-tangent space, $T^*(\mathcal{G})$, of the space of couplings. Using (34) gives $d\hat{\Phi} = 0$ where d represents the exterior derivative on $T^*(\mathcal{G})$, i.e. the one-form $\hat{\Phi}$ is closed because it is, at least locally, exact. This is a direct consequence of the definition (27) and can be thought of as representing the fact that $\hat{\Phi}_a$ are an operational analogue of a co-ordinate basis for real valued one-forms, e.g. for dx^μ , $d^2 x^\mu = 0$. Of course $d\theta^a$ constitute a co-ordinate basis for ordinary real valued one-forms on $T^*(\mathcal{G})$ as usual.

Now consider what happens to the amplitude (1) under a variation of the renormalised couplings. In general the renormalised operators \hat{A}_i , as well as the action itself, will depend on the couplings so

$$\begin{aligned} \partial_b \langle \hat{A}_1(x_1) \cdots \hat{A}_N(x_N) \rangle &= \sum_{i=1}^N \langle \partial_b \hat{A}_i(x_i) \rangle \cdots \hat{A}_N(x_N) \rangle \\ &\quad - \int d^D \tilde{y} \langle \hat{\Phi}'_b(y) \hat{A}_1(x_1) \cdots \hat{A}_N(x_N) \rangle, \end{aligned} \quad (35)$$

where $\hat{\Phi}'_b = \hat{\Phi}_b - \langle \hat{\Phi}_b \rangle$. The term involving $\langle \hat{\Phi}_b \rangle$ appears in here because the variation of the couplings in the normalisation factor $Z_0[g_0, \epsilon, \gamma]^{-1}$, which is necessary for

the definition of physical amplitudes, introduces factors of $\langle \frac{\partial \hat{E}_0(y)}{\partial g^a} \rangle$ into the amplitudes on the right hand side.

Now the trace of the stress operator, \hat{T} , can be expanded as a linear combination $\hat{T}(z) = \beta^b(g) \hat{\Phi}_b(z) - e(g) \hat{E}(z)$ for some real functions, $\beta^b(g)$ and $e(g)$, of g^a . It will be convenient to define $\hat{\Theta}(z) = \beta^b(g) \hat{\Phi}_b(z)$ so that $\hat{T}(z) = \hat{\Theta}(z) - e(g) \hat{E}(z)$. Contracting (35) with the functions β^b yields [B]

$$\begin{aligned} \beta^b \partial_b \langle \hat{A}_1(x_1) \cdots \hat{A}_N(x_N) \rangle &= \sum_{i=1}^N \langle \hat{A}_1(x_1) \cdots \{ (\partial^b \beta^b - e \Delta_\varphi) \hat{A}_i(x_i) \} \cdots \hat{A}_N(x_N) \rangle \\ &\quad - \int d^D \tilde{y} \langle \hat{T}(y) \hat{A}_1(x_1) \cdots \hat{A}_N(x_N) \rangle, \end{aligned} \quad (36)$$

where equation (29) has been used. We now have two equations both involving the expectation value of \hat{T} with the \hat{A}_i , (26) and (36). Combining these equations to eliminate \hat{T} results in the following version of the renormalisation group equation

$$\begin{aligned} \mathcal{L}_\beta \langle \hat{A}_1(x_1) \cdots \hat{A}_N(x_N) \rangle &+ \sum_{i=1}^N \langle \hat{A}_1(x_1) \cdots \{ \Delta_\gamma \hat{A}_i(x_i) \} \cdots \hat{A}_N(x_N) \rangle \\ &= \langle (\beta^b \partial_b - e \Delta_\varphi) \{ \hat{A}_1(x_1) \cdots \hat{A}_N(x_N) \} \rangle > - \beta^b \partial_b \langle \hat{A}_1(x_1) \cdots \hat{A}_N(x_N) \rangle. \end{aligned} \quad (37)$$

This can be related to the more familiar form of the RG equation in the following manner. Consider a scalar field and let $\hat{A}_i(x_i) = \hat{\varphi}^{p_i}(x_i)$, with p_i a positive integer, be monomials of the fundamental fields, so that $\Delta_\varphi \hat{A}_i(x_i) = p_i \hat{A}_i(x_i)$ and $\Delta_\gamma \hat{A}_i(x_i) = 0$. Now, following [7], identify $e(g)$ with the canonical mass dimension, d_φ , of the operator $\hat{\varphi}$ and $\beta^b \partial_b \hat{A}_i(x_i)$ with minus the anomalous dimension of \hat{A}_i so that

$$\beta^b \partial_b \hat{A}_i(x_i) = -\gamma_{A_i} \hat{A}_i(x_i) \quad (38)$$

(γ_{A_i} may be a matrix if there is operator mixing). Finally note that, for scalar operators using Cartesian co-ordinates, $\mathcal{L}_\beta \hat{A}_i(x_i) = x_i^\mu \frac{\partial}{\partial x_i^\mu} \hat{A}_i(x_i)$. Then the renormalisation group equation emerges in the standard homogeneous form

$$\sum_{i=1}^N \left(x_i^\mu \frac{\partial}{\partial x_i^\mu} + d_{A_i} + \gamma_{A_i} \right) \langle \hat{A}_1(x_1) \cdots \hat{A}_N(x_N) \rangle + \beta^b \partial_b \langle \hat{A}_1(x_1) \cdots \hat{A}_N(x_N) \rangle = 0, \quad (39)$$

where $d_{A_i} = p_i d_\varphi$ is the canonical dimension of \hat{A}_i , $\hat{A}_i = \hat{\varphi}^{p_i}$. β^a are thus seen to be nothing other than the usual β functions of the theory, for the couplings g^a ,

$$\beta^a = \kappa \frac{dg^a}{d\kappa}. \quad (40)$$

Since the couplings, g^a , are defined to be massless, these β functions include the canonical dimensions for the couplings. For example, if the theory involves a mass, m^2 , then the

corresponding massless coupling is $\tilde{m}^2 = m^2 \kappa^{-2}$ and $\beta \tilde{m}^2 = (-2 + \delta) \tilde{m}^2$ where δ is the usual anomalous mass dimension, $\kappa \frac{d\tilde{m}^2}{d\kappa} = \delta \tilde{m}^2$. Note that equation (39) is the version of the RG equation for connected amplitudes (or Green's functions $G^{(N)}$) not for proper vertices (one particle irreducible functions, $\Gamma^{(N)}$) which would require the opposite sign for $d_{\hat{A}_i} + \gamma_{\hat{A}_i}$.

Returning to the general form of the RG equation, (37), identifying $e(g)$ with the canonical dimension of the field, d_ϕ , finally gives

$$\begin{aligned} \mathcal{L}_{\tilde{D}} < \hat{A}_1(x_1) \cdots \hat{A}_N(x_N) > + \sum_{i=1}^N < \hat{A}_1(x_1) \cdots \{ \Delta_\gamma \hat{A}_i(x_i) \} \cdots \hat{A}_N(x_N) > \\ &= < (\beta^b \partial_b - d_\phi \Delta_\phi) \{ \hat{A}_1(x_1) \cdots \hat{A}_N(x_N) \} > - \beta^b \partial_b < \hat{A}_1(x_1) \cdots \hat{A}_N(x_N) >, \end{aligned} \quad (41)$$

with $\Delta_\gamma \hat{A}_i$ defined in equation (26) and $\Delta_\phi \hat{A}_i$ in (33). This is the main result of this section. It is a slight generalisation of the usual RG equation for amplitudes. It includes the possibility of \hat{A}_i being Euclidean tensors or even spinors, since the Lie derivative with respect to the dilation generator, $\mathcal{L}_{\tilde{D}}$, automatically tells us how to handle tensor (and spinor) indices, as well as allowing for the possibility that \hat{A}_i might be composite operators involving the Euclidean metric $\gamma_{\mu\nu}$. Clearly this equation is applicable to theories with more than one field, φ , one merely sums $d_\phi \Delta_\phi$ over the fields.

§3 The Renormalisation Group Equation As A Lie Derivative

In this section it will be demonstrated that the RG equation, (41), can be interpreted as an equation for Lie derivatives of amplitudes. The Lie derivative of an amplitude with respect to the dilation generator, \tilde{D} , on Euclidean space is exactly the same as the Lie derivative with respect to the vector $\tilde{\beta} = \beta^a \partial_a$ on the space of couplings. We shall first give a simple proof and then relate the result to the analysis of the previous section.

Since $\hat{\Phi}_a$ are a basis for rotationally invariant operators of dimension D it suffices to consider amplitudes of these operators. All other amplitudes for Euclidean scalars (more precisely densities) which are relevant or marginal operators can be obtained from linear combinations of these basic operators. Consider, therefore, amplitudes of the form

$$\hat{\Phi}_{a_1 \cdots a_N}^{(N)}(x_1, \dots, x_N) = < \hat{\Phi}_{a_1}(x_1) \cdots \hat{\Phi}_{a_N}(x_N) >. \quad (42)$$

The usual derivation of the renormalisation group equation relies on the simple fact that all physical amplitudes should be independent of the renormalisation point κ . However $\hat{\Phi}_{a_1 \cdots a_N}^{(N)}(x_1, \dots, x_N)$ cannot be independent of κ in general. If we chose a different parameterisation of the renormalised couplings (a different co-ordinate system on \mathcal{G}), which will be denoted by primes g' , then $\hat{\Phi}_{a_1 \cdots a_N}^{(N)}(x_1, \dots, x_N)$ transforms as a rank N co-variant tensor,

$$\hat{\Phi}_{a_1 \cdots a_N}^{(N)}(x_1, \dots, x_N) = \left(\frac{\partial g'^1}{\partial g^1} \right) \cdots \left(\frac{\partial g'^N}{\partial g^N} \right) \hat{\Phi}_{a'_1 \cdots a'_N}^{(N)}(x_1, \dots, x_N), \quad (43)$$

and these cannot both be independent of κ , since $\frac{\partial g'^i}{\partial g^i}$ is not, in general. The object that ought to be independent of κ is the reparameterisation invariant amplitude

$$< \hat{\Phi}(x_1) \cdots \hat{\Phi}(x_N) > = < \hat{\Phi}_{a_1}(x_1) \cdots \hat{\Phi}_{a_N}(x_N) > dg^{a_1} \cdots dg^{a_N}. \quad (44)$$

The correct statement that amplitudes are independent of the renormalisation point is,

$$\kappa \frac{d}{d\kappa} < \hat{\Phi}(x_1) \cdots \hat{\Phi}(x_N) > = 0. \quad (45)$$

Allowing for the fact that $< \hat{\Phi}_{a_1}(x_1) \cdots \hat{\Phi}_{a_N}(x_N) >$ are also functions of the g^a gives

$$\begin{aligned} -\kappa \frac{\partial}{\partial \kappa} < \hat{\Phi}_{a_1}(x_1) \cdots \hat{\Phi}_{a_N}(x_N) > = \\ & \beta^b \partial_b < \hat{\Phi}_{a_1}(x_1) \cdots \hat{\Phi}_{a_N}(x_N) > + \sum_{i=1}^N (\partial_{a_i} \beta^b) < \hat{\Phi}_{a_1}(x_1) \cdots \hat{\Phi}_{a_N}(x_N) >, \end{aligned} \quad (46)$$

where we have used

$$\kappa \frac{d}{d\kappa} (dg^a) = d \left(\kappa \frac{dg^a}{d\kappa} \right) = d\beta^a = \frac{\partial \beta^a}{\partial g^b} dg^b. \quad (47)$$

In co-ordinate free notation this is

$$-\kappa \frac{\partial}{\partial \kappa} < \hat{\Phi}(x_1) \cdots \hat{\Phi}(x_N) > = \mathcal{L}_{\tilde{\beta}} < \hat{\Phi}(x_1) \cdots \hat{\Phi}(x_N) >, \quad (48)$$

where $\mathcal{L}_{\tilde{\beta}}$ represents Lie differentiation on \mathcal{G} with respect to the vector field $\tilde{\beta}$. Thus the renormalisation group equation is nothing other than the definition of a Lie derivative provided that it is appreciated that the amplitudes are tensors on $T^*(\mathcal{G})$, and not scalars (the minus sign in (48) is standard in the definition of a Lie derivative, see e.g. [13]). This analysis makes it clear that the anomalous dimensions have the geometrical interpretation of arising from the change in dg^a (which are a basis for real valued one-forms on $T^*(\mathcal{G})$) as we move along the vector field $\tilde{\beta}$.

Equation (48) can be expressed in terms of the dilation vector on Euclidean space, \tilde{D} , using the usual scaling arguments. To see this first note that $\hat{\Phi}_a$ are densities therefore in Cartesian co-ordinates

$$\mathcal{L}_{\tilde{D}} \hat{\Phi}_a(x_i) = \frac{\partial}{\partial x_i^\mu} (x_i^\mu \hat{\Phi}_a(x_i)) = \left(x_i^\mu \frac{\partial}{\partial x_i^\mu} + D \right) \hat{\Phi}_a(x_i). \quad (49)$$

Now by definition, equation (27), $\hat{\Phi}_a(x)$ has mass dimension D thus the usual naive scaling arguments give

$$\begin{aligned} \kappa \frac{\partial}{\partial \kappa} < \hat{\Phi}_{a_1}(x_1) \cdots \hat{\Phi}_{a_N}(x_N) > &= \sum_{i=1}^N \left(x_i^\mu \frac{\partial}{\partial x_i^\mu} + D \right) < \hat{\Phi}_{a_1}(x_1) \cdots \hat{\Phi}_{a_N}(x_N) > \\ &= \mathcal{L}_{\tilde{D}} < \hat{\Phi}_{a_1}(x_1) \cdots \hat{\Phi}_{a_N}(x_N) >, \end{aligned} \quad (50)$$

hence

$$\mathcal{L}_{\vec{D}} \langle \hat{\Phi}(x_1) \cdots \hat{\Phi}(x_N) \rangle = -\mathcal{L}_{\vec{\beta}} \langle \hat{\Phi}(x_1) \cdots \hat{\Phi}(x_N) \rangle. \quad (51)$$

This is the main result of this paper. The RG equation is seen as a bridge connecting a particular diffeomorphism (dilations generated by the conformal Killing vector; \vec{D}) of Euclidean space, \mathbf{R}^D , with a diffeomorphism (generated by the associated vector $\vec{\beta}$) of the space of couplings \mathcal{G} . (I apologise for the minus sign in equation (51). This is due to the usual field theory definition of β^a as being derivatives of the couplings with respect to the renormalisation mass κ . Had they been defined as derivatives with respect to a length then this equation would have had a plus sign on the right hand side.)

This can be related to the analysis of the previous section and equation (41) in the following manner. Let $\hat{A}_i(x_i) = \hat{\Phi}_{a_i}(x_i)$ in (41). Now consider the meaning of $\Delta_\gamma \hat{\Phi}_a$. Using the definition of $\hat{\Phi}_a$ in equation (27) and assuming that metric variations commute with variations of the couplings leads to

$$\Delta_\gamma \hat{\Phi}_a = 2\gamma^{\mu\nu} \frac{\delta}{\delta\gamma^{\mu\nu}} \left(\frac{\partial \hat{H}_0}{\partial g^a} \right) = \frac{\partial}{\partial g^a} \left(2\gamma^{\mu\nu} \frac{\delta \hat{H}_0}{\delta\gamma^{\mu\nu}} \right) = \partial_a \hat{T}. \quad (52)$$

Note that (34) implies that

$$\partial_a \hat{T} = (\partial_a \beta^b) \hat{\Phi}_b + \beta^b \partial_b \hat{\Phi}_a - d_\varphi \partial_a \hat{E} = \partial_a \hat{\Theta} - d_\varphi \partial_a \hat{E} \quad (53),$$

where the previous identification of $e(g)$ with d_φ has been made. Now assuming that field variations commute with coupling variations gives

$$\Delta_\varphi \hat{\Phi}_a = \Delta_\varphi \partial_a \hat{H}_0 = \partial_a \Delta_\varphi \hat{H}_0 = \partial_a \hat{E}, \quad (54)$$

where (29) and (33) have been used. Putting all this together (41) becomes

$$\begin{aligned} \mathcal{L}_{\vec{D}} \langle \hat{\Phi}_{a_1}(x_1) \cdots \hat{\Phi}_{a_N}(x_N) \rangle &= \sum_{i=1}^N \langle \hat{\Phi}_{a_1}(x_1) \cdots \{ \beta^b \partial_b \hat{\Phi}_{a_i}(x_i) \} \cdots \hat{\Phi}_{a_N}(x_N) \rangle > -\beta^b \partial_b \langle \hat{\Phi}_{a_1}(x_1) \cdots \hat{\Phi}_{a_N}(x_N) \rangle > \\ &= \sum_{i=1}^N \langle \hat{\Phi}_{a_1}(x_1) \cdots \hat{\Phi}_{a_N}(x_N) \rangle > - \sum_{i=1}^N \langle \hat{\Phi}_{a_1}(x_1) \cdots \{ \partial_{a_i} \hat{\Theta}(x_i) \} \cdots \hat{\Phi}_{a_N}(x_N) \rangle > \\ &= -\beta^b \partial_b \langle \hat{\Phi}_{a_1}(x_1) \cdots \hat{\Phi}_{a_N}(x_N) \rangle > - \sum_{i=1}^N \langle \partial_{a_i} \beta^b \rangle \langle \hat{\Phi}_{a_1}(x_1) \cdots \hat{\Phi}_b(x_i) \cdots \hat{\Phi}_{a_N}(x_N) \rangle >. \end{aligned} \quad (55)$$

which is just equation (51). From this analysis it is clear that the source of the anomalous dimensions in the RG equation is the term involving $\partial_{a_i} \beta^b$ in this equation, which comes from Lie transport of the basis dg^a (this term actually contains both the canonical dimensions and the anomalous dimensions because the couplings are defined to be dimensionless).

Another way of expressing this idea is to define the matrix of renormalisation coefficients, Z_a^b , by

$$\hat{\Phi}_{oa} = Z_a^b(g) \hat{\Phi}_b, \quad (56)$$

where $\hat{\Phi}_{oa}$ constitute the bare basis

$$\hat{\Phi}_{oa} = \frac{\partial \hat{H}_0}{\partial g^a}. \quad (57)$$

The matrix Z_a^b depends on the renormalisation scheme, of course. For example in dimensional regularisation it is a function not only of the renormalised couplings, g^a , but also of the dimension $D = 4 - \epsilon$ and can be expanded as a series of poles in ϵ . Clearly the definition of the renormalised basis (27) also implies that

$$Z_a^b(g) = \frac{\partial g^b}{\partial g^a} \Leftrightarrow d g^a Z_a^b(g) = d g^b \quad (58)$$

so we can write

$$\hat{\Phi}(x) = \hat{\Phi}_{oa}(x) d g^a = \hat{\Phi}_a(x) d g^a. \quad (59)$$

We now demand that the bare operators be independent of the renormalisation point. There is a slight subtlety here, though, in that the bare couplings are defined to be massless. This means that they change, under changes in κ , by their canonical dimensions, e.g. for a mass the massless bare coupling is $\tilde{m}_0^2 = m_0^2 \kappa^{-2}$ and $\kappa \frac{d\tilde{m}_0^2}{d\kappa} = 0$ requires $\kappa \frac{d\tilde{m}_0^2}{d\kappa} = -2\tilde{m}_0^2$. We shall denote the canonical dimension of the coupling g^a by d_a , thus $\kappa \frac{d g^a}{d\kappa} = -d_a g^a$ where there is no sum over a . As in the previous argument for amplitudes, it is the operator-valued one form $\hat{\Phi}_a = \hat{\Phi}_{oa} d g^a$ which is independent of κ . Thus

$$\kappa \frac{d \hat{\Phi}_{oa}}{d\kappa} - d_a \hat{\Phi}_{oa} = 0, \quad \Leftrightarrow \quad \kappa \frac{d \hat{\Phi}_a}{d\kappa} + \Gamma_a^b \hat{\Phi}_b = 0 \quad (60)$$

or

$$\left(\kappa \frac{\partial}{\partial \kappa} + \beta^b \partial_b \right) \hat{\Phi}_a + \Gamma_a^b \hat{\Phi}_b = 0, \quad (61)$$

where the matrix, Γ_a^b , is defined by

$$\Gamma_a^b = \frac{\partial \beta^b}{\partial g^a}. \quad (62)$$

This should be distinguished from the matrix of anomalous dimensions,

$$\gamma_a^b = (Z^{-1})^a_c \left(\frac{d Z_c^b}{d\kappa} \right). \quad (63)$$

Equation (60) is a renormalisation group equation for the operators $\hat{\Phi}_a$. One must be careful in evaluating amplitudes involving this equation however since neither $\kappa \frac{d}{d\kappa}$ nor

$\beta^b \partial_b$ can be pulled outside of expectation values separately, though the combination can be since S_0 is independent of κ .

It is crucial to this interpretation that massless couplings are used. In particular this means that

$$\bar{\beta} = \beta^a \frac{\partial}{\partial g^a} = \beta_0^a \frac{\partial}{\partial g_0^a}, \quad (64)$$

where $\beta_0^a = \kappa \frac{d\beta^a}{d\kappa} = -d_a$ are just the canonical dimensions. Had massive couplings been used, then β_0^a would be zero since the bare massive couplings are defined to be independent of κ , but β^a are non-zero in general. One can hardly transform from a non-zero vector to one which vanishes using a co-ordinate transformation! (This has nothing to do with the "singularities" of the renormalisation program, the regulator is still in place so the co-ordinate transformation is still non-singular.) The important quantity is the total dimension, canonical plus anomalous, and to split it up spoils general co-ordinate invariance. Stated differently, the redefinition $m^2 \rightarrow \bar{m}^2 = m^2 \kappa^{-2}$ is *not* a co-ordinate transformation because κ is not a co-ordinate.

We end this section with some comments on the matrix of dimensions (canonical plus anomalous), $\Gamma_a^b = \frac{\partial \beta^b}{\partial g^a}$, which appears in (55) and (60). This matrix plays a very important rôle near a critical point in statistical mechanics - its eigenvalues determine the relevant and irrelevant directions in the space of interactions. A linear combination of operators corresponding to a positive eigenvalue (negative mass dimension of a coupling) tends to blow up for large κ and decrease for small κ - such an operator is termed irrelevant and corresponds to a non-renormalisable interaction in field theory. A negative or zero eigenvalue (relevant or marginal operator) corresponds to a renormalisable interaction in field theory. Thus for a renormalisable field theory near a fixed point, the matrix of dimensions had better have only non-positive eigenvalues but this restriction is not necessary away from the fixed point. This is related to the triviality problem for $\lambda\varphi^4$ in four dimensions. In $D = 4 - \epsilon$ dimensions the canonical dimension of the coupling, λ , is ϵ and in perturbation theory β starts of at order λ^2 , $\beta^\lambda = -\epsilon\lambda + \frac{3}{16\pi^2}\lambda^2 + \dots$ and $\partial_\lambda \beta^\lambda = -\epsilon + \frac{3}{8\pi^2}\lambda + \dots$. Provided $\epsilon > 0$ this is negative for small λ but when $\epsilon = 0$ it is positive. This does not of course mean that the theory is non-renormalisable in four dimensions because at the fixed point $\lambda = \lambda_* = 0$, $(\partial_\lambda \beta^\lambda)_* = 0$, i.e. λ is a marginal coupling at the trivial fixed point in four dimensions.

In a theory in which Γ has complex eigenvalues one would have operators with complex anomalous dimensions which would presumably correspond to a non-unitary theory. It is shown in [14] that this circumstance is automatically avoided if the β -functions are derivable from a potential function.

§4 An Example - $\lambda\varphi^4$ Theory

The ideas presented in the previous sections will now be exemplified in a concrete model - that of massive $\lambda\varphi^4$ in $D = 4 - \epsilon$ Euclidean dimensions with a constant source. The renormalisation procedure adopted will be that of dimensional regularisation and

minimal subtraction. This section is a generalisation of the concepts developed in [12]. The action is

$$S_0 = \int d^4-x \left(\frac{1}{2} (\partial\varphi_0(x))^2 + \frac{1}{2} m_0^2 \varphi_0^2(x) + j_0 \varphi_0(x) + \frac{1}{4!} \lambda_0 \varphi_0^4(x) \right). \quad (65)$$

The space \mathcal{G} is thus three dimensional and is parameterised by three renormalised couplings

$$g^1 = \lambda, \quad g^2 = \bar{j} = j \kappa^{-3+\epsilon/2}, \quad \text{and} \quad g^3 = \bar{m}^2 = m^2 \kappa^{-2}, \quad (66)$$

where, as usual in dimensional regularisation, λ is dimensionless. The massless bare couplings are,

$$g_0^1 = \bar{\lambda}_0 = \lambda_0 \kappa^{-\epsilon}, \quad g_0^2 = \bar{j}_0 = j_0 \kappa^{-3+\epsilon/2}, \quad \text{and} \quad g_0^3 = \bar{m}_0^2 = m_0^2 \kappa^{-2}. \quad (67)$$

The basis operators, $\hat{\Phi}_{0a}(x)$ of equation (27), can thus be represented as a three component vector (in Cartesian co-ordinates so that $\sqrt{7} = 1$),

$$\hat{\Phi}_0(x) = \begin{pmatrix} \hat{\Phi}_{\bar{\lambda}_0} \\ \hat{\Phi}_{\bar{j}_0} \\ \hat{\Phi}_{\bar{m}_0^2} \end{pmatrix} = \begin{pmatrix} \frac{1}{4!} \kappa^\epsilon \varphi_0^4(x) \\ \kappa^{3-\epsilon/2} \varphi_0(x) \\ \frac{1}{2} \kappa^{2-\epsilon/2} \varphi_0^2(x) \end{pmatrix}. \quad (68)$$

$\hat{\Phi}_0$, as defined here, differs, by factors of the coupling constants, from the Q_0 of reference [12]. This is because (68) is defined so as to have a natural interpretation as a co-vector in $T^*(\mathcal{G})$, which transforms co-variantly under general co-ordinate transformations on \mathcal{G} , whereas multiplying by couplings spoils this interpretation.

The operator $\partial^2 \varphi_0^2(x)$, which is also of dimension $4 - \epsilon$ and could in principle mix with $\hat{\Phi}_0(x)$, has been omitted. More will be said concerning this operator later but for the moment we merely note that it is a total derivative and does not contribute to the action. The only other operator with the same dimensions as $\hat{\Phi}_0(x)$ is $\varphi_0 \partial^2 \varphi_0(x)$ but rather than use this directly the equations of motion can be used as the remaining linearly independent operator,

$$E_0(x) = \varphi_0 \left(-\partial^2 \varphi_0 + m_0^2 \varphi_0 + j_0 + \frac{1}{6} \lambda_0 \varphi_0^3 \right) \Big|_x. \quad (69)$$

This is convenient because, as explained in §2, the equations of motion do not get renormalised, $\hat{E}_0 = \hat{E}$.

The renormalisation constants, Z_a^b , relate bare to renormalised operators

$$\hat{\Phi}_0(x) = Z \hat{\Phi}(x) = Z \begin{pmatrix} \hat{\Phi}_\lambda \\ \hat{\Phi}_j \\ \hat{\Phi}_{\bar{m}^2} \end{pmatrix}. \quad (70)$$

The matrix Z can be determined as follows. Relate the bare couplings g_0^a to their renormalised counterparts in the usual way,

$$\lambda_0 = z_\lambda^{-1}(\lambda, \epsilon) \kappa^\epsilon \lambda, \quad j_0 = z_j^{-1}(\lambda, \epsilon) j \quad \text{and} \quad m_0^2 = z_m^{-1}(\lambda, \epsilon) m^2. \quad (71)$$

Then demanding that λ_0 , j_0 and m_0^2 be independent of κ gives

$$\begin{aligned} \kappa \frac{d\lambda}{d\kappa} &= -\epsilon\lambda + \beta & \text{with} & \quad \frac{\beta}{\lambda} = z_\lambda^{-1} \left(\kappa \frac{dz_\lambda}{d\kappa} \right) \\ \kappa \frac{dj}{d\kappa} &= \left(-3 + \frac{\epsilon}{2} + \gamma \right) j & \text{with} & \quad \gamma = z_1^{-1} \left(\kappa \frac{dz_1}{d\kappa} \right) \\ \kappa \frac{d\tilde{m}^2}{d\kappa} &= (-2 + \delta) \tilde{m}^2 & \text{with} & \quad \delta = z_2^{-1} \left(\kappa \frac{dz_2}{d\kappa} \right), \end{aligned} \quad (72)$$

where β , γ and δ are functions of λ only, independent of ϵ . $\beta(\lambda)$ is the usual four dimensional β -function which starts with λ^2 in perturbation theory. The vector $\vec{\beta}$ of equation (40) is thus,

$$\vec{\beta} = \begin{pmatrix} -\epsilon\lambda + \beta \\ (-3 + \epsilon/2 + \gamma)j \\ (-2 + \delta)\tilde{m}^2 \end{pmatrix}. \quad (73)$$

Equations (71) and (72) now give

$$\begin{aligned} d\tilde{\lambda}_0 &= -\frac{\epsilon z_\lambda^{-1} \lambda}{(-\epsilon\lambda + \beta)} d\lambda \\ d\tilde{j}_0 &= z_1^{-1} dj - \frac{\gamma z_1^{-1} j}{(-\epsilon\lambda + \beta)} d\lambda \\ d\tilde{m}_0^2 &= z_2^{-1} d\tilde{m}^2 - \frac{\delta z_2^{-1} \tilde{m}^2}{(-\epsilon\lambda + \beta)} d\lambda, \end{aligned} \quad (74)$$

Using (58),

$$Z_a^b(g) = \frac{\partial g^b}{\partial g^a} \Leftrightarrow dg_a^b Z_a^b(g) = dg^b, \quad (58)$$

immediately leads to

$$(Z^{-1})_a^b = \frac{\partial g_a^b}{\partial g^a} = \begin{pmatrix} -\frac{\epsilon z_\lambda^{-1} \lambda}{(-\epsilon\lambda + \beta)} & -\frac{\gamma z_1^{-1} j}{(-\epsilon\lambda + \beta)} & -\frac{\delta z_2^{-1} \tilde{m}^2}{(-\epsilon\lambda + \beta)} \\ 0 & z_1^{-1} & 0 \\ 0 & 0 & z_2^{-1} \end{pmatrix} \quad (75)$$

or

$$Z = \begin{pmatrix} -\frac{\epsilon\lambda}{(-\epsilon\lambda + \beta)z_\lambda} & -\frac{\gamma j}{z_1} & -\frac{\delta \tilde{m}^2}{z_2} \\ 0 & z_1 & 0 \\ 0 & 0 & z_2 \end{pmatrix}. \quad (76)$$

The matrix of dimensions, Γ , follows easily from (73),

$$\Gamma_a^b = \frac{\partial \beta^b}{\partial g^a} = \begin{pmatrix} -\epsilon + \beta & \gamma & \delta \\ 0 & -3 + \epsilon/2 + \gamma & 0 \\ 0 & 0 & -2 + \delta \end{pmatrix}, \quad (77)$$

where a dot denotes differentiation with respect to λ . This should be compared with the matrix of anomalous dimensions

$$\gamma^a_b = (Z^{-1})^a_c \kappa \left(\frac{dZ^c_b}{d\kappa} \right) = \begin{pmatrix} \beta & \left[\dot{\gamma} + \frac{3(-1+\epsilon/2)\dot{\gamma}}{(-\epsilon\lambda + \beta)} \right] \dot{j} & \left[\dot{\delta} + \frac{2(-1+\epsilon/2)\dot{\delta}}{(-\epsilon\lambda + \beta)} \right] \dot{\tilde{m}}^2 \\ 0 & \gamma & \delta \\ 0 & 0 & \delta \end{pmatrix}. \quad (78)$$

It is worth noting that $\hat{\Phi}_\lambda$ is not simply $\frac{1}{4}\dot{\varphi}^4$. It can be shown, using the techniques of reference [12], that it mixes with the equation of motion operator, \hat{E} . In fact

$$\begin{aligned} \hat{\Phi}_\lambda &= \frac{\kappa^4}{4!} \dot{\varphi}^4 - \frac{\gamma}{(-\epsilon\lambda + \beta)} \hat{E} \\ \hat{\Phi}_j &= \kappa^3 - \epsilon/2 \dot{\varphi} \\ \hat{\Phi}_{m^2} &= \frac{1}{2} \kappa^2 - \epsilon/2 \dot{\varphi}^2, \end{aligned} \quad (79)$$

where $\dot{\varphi}^4$, $\dot{\varphi}^2$ and $\dot{\varphi}$ are the renormalised operators. We will not prove these relations here, since they will not be needed, but merely refer to Brown's paper [12].

Let us now turn to the consideration of the stress tensor. Variation of the metric in (65) produces the classical stress tensor

$$T_{\mu\nu}^C = \partial_\mu \varphi_a \partial_\nu \varphi_a - \delta_{\mu\nu} \left\{ \frac{1}{2} (\partial\varphi_a)^2 + \frac{1}{2} m_0^2 \varphi_a^2 + j_0 \varphi_0 + \frac{1}{4!} \lambda_0 \varphi_0^4 \right\}, \quad (80)$$

where $\gamma_{\mu\nu}$ has been set to $\delta_{\mu\nu}$ after the variation. As is well known this gives rise to an infinite operator, [9], but it can be made finite by adding the "improvement" term

$$T_{\mu\nu} = \partial_\mu \varphi_a \partial_\nu \varphi_a - \delta_{\mu\nu} \left\{ \frac{1}{2} (\partial\varphi_a)^2 + \frac{1}{2} m_0^2 \varphi_a^2 + j_0 \varphi_0 + \frac{1}{4!} \lambda_0 \varphi_0^4 \right\} - \frac{1}{4} \left(\frac{2-\epsilon}{3-\epsilon} \right) (\partial_\mu \partial_\nu - \delta_{\mu\nu} \partial^2) \varphi_0^2. \quad (81)$$

The improvement term can be obtained by introducing the Riemann curvature, \mathcal{R} , adding a coupling to the curvature, $\frac{1}{2} \mathcal{R} \varphi_0^2$, with co-efficient $\frac{1}{4} \left(\frac{2-\epsilon}{3-\epsilon} \right)$ to (65) and then varying the metric before setting $\mathcal{R} = 0$ to give Euclidean space. Of course this new coupling should also run, but it is omitted here for simplicity. It can be treated quite consistently and is done so in [15], but it is sufficient for our purposes to know that it can be done.

Now $T_{\mu\nu}$ is conserved (modulo the equations of motion) and finite, i.e. $T_{\sigma\mu\nu} = T_{\mu\nu}$. Note that an arbitrary finite multiple of $(\partial_\mu \partial_\nu - \delta_{\mu\nu} \partial^2) z_2^{-1} \varphi_0^2$ can be added to (81) without affecting either of these properties, but we choose not to do this because (81) as it stands is manifestly independent of κ and any further finite additions would spoil this feature (see below).

From (81) the trace of the stress tensor is, using the equations of motion, (69), to eliminate $\varphi_0 \partial^2 \varphi_0$,

$$T = (-3 + \epsilon/2) j_0 \varphi_0 - 2 \left(\frac{1}{2} m_0^2 \varphi_0^2 \right) - \epsilon \left(\frac{1}{4!} \lambda_0 \varphi_0^4 \right) + (-1 + \epsilon/2) E_0. \quad (82)$$

The co-efficients in this equation are just given by minus the classical dimensions of the couplings and minus the canonical dimension of φ for E_0 . It is a simple matter to use the matrix Z^a_b to convert this to an expression involving only renormalised quantities,

$$\begin{aligned} \hat{T} &= (-3 + \epsilon/2 + \gamma)j\hat{\Phi}_j + (-2 + \delta)\frac{m^2}{2}\hat{\Phi}_{m^2} + (-\epsilon\lambda + \beta)\hat{\Phi}_\lambda + (-1 + \epsilon/2)E \\ &= (-3 + \epsilon/2 + \gamma)j\hat{\varphi} + (-2 + \delta)\frac{m^2}{2}\hat{\varphi}^2 + (-\epsilon\lambda + \beta)\frac{1}{4!}\kappa^4\hat{\varphi}^4 + (-1 + \epsilon/2 - \gamma)\hat{E}, \end{aligned} \quad (83)$$

which is the same as the expression in [12]. Thus

$$\hat{T}_0 = \hat{T} = \beta^a \hat{\Phi}_a - (1 - \epsilon/2)\hat{E} = \hat{\Theta} - (1 - \epsilon/2)\hat{E}, \quad (84)$$

which is the expansion of \hat{T} used in §2, with the co-efficient of \hat{E} being the canonical dimension of the field, as was shown in that section. Since the equations of motion do not get renormalised it immediately follows that $\hat{\Theta}$ is also not renormalised $\hat{\Theta}_0 = \hat{\Theta}$.

It is straightforward to show, using (74), that

$$\beta^a \partial_a g^b = -d_a g^a \quad (\text{no sum over } a). \quad (85)$$

This is an explicit example of equation (64) of the previous section, the statement of the fact that

$$\tilde{\beta} = \beta^a \frac{\partial}{\partial g^a} = \beta^a \frac{\partial}{\partial g^a} \quad \text{where} \quad \beta^a = \beta^b \left(\frac{\partial g^a}{\partial g^b} \right) = -d_a, \quad (86)$$

i.e. the change from bare to renormalised couplings is just a co-ordinate transformation in this language (it is a singular co-ordinate transformation when $\epsilon = 0$, but this need not worry us!).

Note that \hat{T}_0 and \hat{T} are the *same* operator. The bare stress operator is equal to the renormalised stress operator, once the improvement term (which corresponds to the purely classical conformal coupling to the curvature) is added. This is an explicit example of the non-renormalisation theorem discussed in §2.

Since $T_{\mu\nu}$ should be independent of the renormalisation point, κ , we can immediately deduce, from (23) that

$$\kappa \frac{d\hat{T}_{\mu\nu}}{d\kappa} = 0 \quad (87)$$

and, in particular,

$$\kappa \frac{d\hat{T}}{d\kappa} = 0 \quad \Leftrightarrow \quad \kappa \frac{d\hat{\Theta}}{d\kappa} = 0. \quad (88)$$

It is worthwhile making a comment about the operator $\partial^2 \hat{\varphi}^2$ at this point. In reference [12] this operator was included in the basis set and a renormalisation co-efficient introduced for it. This co-efficient has the effect of modifying the factor $\frac{1}{4} \left(\frac{2-\epsilon}{3-\epsilon} \right)$ in the stress tensor, (81), but does not affect the form of the trace, (83). However it does affect the κ dependence and results in a stress operator for which $\kappa \frac{d\hat{T}_{\mu\nu}}{d\kappa} \neq 0$ but instead depends (in a finite way)

on the new renormalisation constant. This is the usual ambiguity in the definition of the stress operator, but here we turn this ambiguity into a virtue and use the freedom it gives to *choose* a stress operator satisfying (88). This requires omitting the operator $\partial^2 \hat{\varphi}^2$ from the basis altogether, in flat space, and considering $\partial^2 \hat{\varphi}^2$, to be not fundamental, but rather derivable from the more fundamental operator $\hat{\varphi}^2$. This is somewhat similar to the idea of "primary" and "descendant" fields in two dimensional conformal field theory, [8]. In this language $\hat{\varphi}^2$ would be considered to be a primary operator and $\partial^2 \hat{\varphi}^2$ to be a descendant, though of course there is as yet no general theory of the classification of such objects in anything other than two dimensions.

In a curved space the story becomes much more complicated and more terms, involving invariants of the Riemann tensor, must be introduced for a consistent description of the theory, [15]. One would not expect (88) to hold in a curved space since the curvature introduces another length scale into the theory and the right hand side would involve tensor operators which incorporate the Riemann tensor.

It is not difficult to verify that

$$\beta^a \partial_a \hat{\Phi}_j = -\gamma \hat{\Phi}_j \quad \text{and} \quad \beta^a \partial_a \hat{\Phi}_{m^2} = -\delta \hat{\Phi}_{m^2}, \quad (89)$$

as claimed in §2. The anomalous dimension of $\hat{\Phi}_\lambda$ is, however, more complicated

$$\beta^a \partial_a \hat{\Phi}_\lambda = - \left\{ \hat{\beta} \hat{\Phi}_\lambda + \left(\gamma - \frac{3(1-\epsilon/2)}{(-\epsilon\lambda + \beta)} \right) j \hat{\Phi}_j - \left(\delta + \frac{2(1-\epsilon/2)}{(-\epsilon\lambda + \beta)} \delta \right) m^2 \hat{\Phi}_{m^2} \right\}. \quad (90)$$

We note here also the following interesting relation,

$$\beta^a \partial_a \hat{T} = -(4 - \epsilon)\hat{\Theta} = -(4 - \epsilon)\beta^a \hat{\Phi}_a = -(4 - \epsilon)(\hat{T} + d_\varphi \hat{E}). \quad (91)$$

This equation is perhaps most easily obtained by expressing \hat{T} in terms of bare quantities and using equation (86).

The co-efficient $4 - \epsilon$ is the canonical dimension of \hat{T} . The operator \hat{T} is, of course, an operator valued scalar on \mathcal{G} , not a tensor, so there is no contribution from $d g^a$ to interpret as anomalous. Equation (91) can be thought of as giving the *total* dimension of \hat{T} , canonical plus anomalous. This is yet another statement of the non-renormalisation of the stress tensor - its anomalous dimension, and therefore the anomalous dimension of the trace, vanishes, modulo the equations of motion.

Since $\hat{\Phi}_a$ are a linearly independent basis of operators one can decompose (91) and deduce that

$$\partial_a \hat{T} = -(4 - \epsilon)\hat{\Phi}_a, \quad (92)$$

a formula that can also be checked explicitly for each $\hat{\Phi}_a$ independently. This formula shows that the basis $\hat{\Phi}_a$ can be determined purely from the knowledge of the trace of the stress operator, \hat{T} , *without* the Hamiltonian, H_0 . Thus \hat{T} contains all the information of the theory since all physical amplitudes can be calculated from a knowledge of this operator.

§5 Conclusions

It has been argued that the renormalisation group equation for physical amplitudes can be given a geometrical interpretation in the sense that it may be viewed as an equation for Lie transport. In terms of Lie transport on the space of couplings it reduces to no more than the definition of a Lie derivative, but its real significance lies in the way that it ties a particular diffeomorphism of Euclidean space, that of dilations, to the diffeomorphism of the space of couplings generated by the vector field $\vec{\beta}$ through equation (51)

$$\mathcal{L}_{\vec{\beta}} < \hat{\Phi}(x_1) \cdots \hat{\Phi}(x_N) > = -\mathcal{L}_{\vec{\beta}} < \hat{\Phi}(x_1) \cdots \hat{\Phi}(x_N) >. \quad (51)$$

It is crucial to this interpretation that N -point amplitudes be viewed as rank N co-variant tensors on the space of couplings, and equation (51) states that the two diffeomorphisms are completely equivalent. From this point of view the anomalous dimension terms in the RG equation are seen as coming from the change in the basis dg^a under Lie transport along the trajectories of $\vec{\beta}$. Since Lie derivatives are the natural geometric way in which to describe symmetries, this picture gives a clearer insight into the structure of the RG equation and the manner in which conformal symmetry, and perhaps also other space symmetries, may be broken. For instance one might postulate the existence of a β -function, $\mathcal{L}_{\vec{\beta}x}$ associated with a more general diffeomorphism, \vec{X} , of space and consider the equation

$$\mathcal{L}_{\vec{X}} < \hat{\Phi}(x_1) \cdots \hat{\Phi}(x_N) > = -\mathcal{L}_{\vec{\beta}x} < \hat{\Phi}(x_1) \cdots \hat{\Phi}(x_N) >. \quad (93)$$

This would presumably necessitate the introduction of position dependent couplings, a concept which has already proved to be of some use in understanding the RG equation [16].

An alternative version of the RG equation, for operators rather than amplitudes, is given by equation (61),

$$\left(\kappa \frac{\partial}{\partial \kappa} \right) + \beta^b \partial_b \hat{\Phi}_a + \Gamma_a^b \hat{\Phi}_b = 0, \quad (61)$$

but it must be stressed that neither $\kappa \frac{\partial}{\partial \kappa} |g$ nor $\beta^b \partial_b$ can be pulled out of expectation values separately since the action, S_0 , is not invariant under either separately.

The emphasis here is on a co-variant description of the RG equation and the notion that one is free to choose any set of renormalised co-ordinates that one wishes in a description of physical amplitudes.

A generalised form of the RG equation, for arbitrary operators rather than just the basis $\hat{\Phi}_a$, is given by equation (41) which allows for the possibility of composite operators, which may depend on the Euclidean metric, as well as operators which are Euclidean tensors rather than scalars.

The example of massive $\lambda\phi^4$ with a constant source has been analysed in detail where the rôle of the stress tensor is clearly displayed. Equation (92) seems very important in this respect. It states that complete knowledge of the basis operators $\hat{\Phi}_a$, and therefore of the entire theory, is in principle obtainable just from the trace of the stress tensor without any knowledge of the underlying Hamiltonian. This seems to be a general property, not

specific to this theory, and it would be interesting to examine this in other examples. In particular the subtleties associated with gauge theories have not been addressed here. The definition of a renormalised stress tensor for a gauge theory is fraught with difficulties but it is to be hoped that, since gauge theories are one of the jewels in the crown of the geometric picture of physics and the analysis presented here is fundamentally geometric in approach, that gauge theories should fit very naturally into this picture but this is not yet clear.

It is a pleasure to thank Denjoe O'Connor for many discussions on the geometric nature of the RG.

References

- [1] E.C.G. Stueckelberg and A. Petermann, *Helv. Phys. Acta.* **26**, (1953), 499
- [2] M. Gell-Mann and F. Low, *Phys. Rev.* **98**, (1954), 1300
- [3] C.G. Callan, *Phys. Rev. D2*, (1970), 1541
K. Symanzik, *Comm. Math. Phys.* **18**, (1970), 227
- [4] S. Weinberg, *Phys. Rev. D8*, (1973), 3497
G. 'tHooft, *Nuc. Phys.* **B61**, (1973), 455
- [5] K.G. Wilson and J. Kogut, *Phys. Rep.* **12C**, (1974), 75
- [6] D. O'Connor and C.R. Stephens, *Geometry The Renormalisation Group And Gravity* D.I.A.S. Preprint (1992)
- [7] A.B. Zamolodchikov, *Rev. Math. Phys.* **1**, (1990), 197
- [8] A.A. Belavin, A.M. Polyakov and A.B. Zamolodchikov, *Nuc. Phys.* **B241**, (1984), 333
- [9] C.G. Callan, S. Coleman and R. Jackiw, *Ann. Phys.* **59**, (1970), 42
- [10] D.Z. Freedman, I.J. Muzinich and E.J. Weinberg, *Ann. Phys.* **87**, (1974), 95
- [11] S.D. Joglekar, *Ann. Phys.* **100**, (1976), 395
- [12] L.S. Brown, *Ann. Phys.* **126**, (1980), 135
- [13] S.W. Hawking and G.F.R. Ellis, *The Large Scale Structure Of Space-Time* C.U.P. (1973)
- [14] D.J. Wallace and R.K.P. Zia, *Phys. Lett.* **48A**, (1974), 325
Ann. Phys. **92**, (1975), 142
- [15] L.S. Brown and J.C. Collins, *Ann. Phys.* **130**, (1980), 215
- [16] I.T. Drummond and G.M. Shore, *Phys. Rev. D19*, (1979), 1134
H. Osborn, *Phys. Lett.* **B222**, (1989), 97
I. Jack and H. Osborn, *Nucl. Phys.* **343**, (1990), 647
G.M. Shore, *Nucl. Phys.* **B362**, (1991), 85