

# The latent structure of ICD-11 posttraumatic stress disorder (PTSD) and complex PTSD in a general population sample from USA: A factor mixture modelling approach<sup>☆</sup>

Enya Redican<sup>a</sup>, Marylene Cloitre<sup>b</sup>, Philip Hyland<sup>c</sup>, Orla McBride<sup>a</sup>, Thanos Karatzias<sup>d</sup>,  
Jamie Murphy<sup>a</sup>, Mark Shevlin<sup>a,\*</sup>

<sup>a</sup> Ulster University, School of Psychology, Coleraine, UK

<sup>b</sup> National Center for PTSD, Veterans Affairs Palo Alto Health Care System, Palo Alto, CA, USA

<sup>c</sup> Department of Psychology, Maynooth University, Ireland

<sup>d</sup> Edinburgh Napier University, School of Health and Social Care, Edinburgh, UK

## ARTICLE INFO

### Keywords:

ICD-11  
Posttraumatic Stress Disorder  
Complex Posttraumatic Stress Disorder  
Factor mixture model

## ABSTRACT

The validity of ICD-11 Posttraumatic Stress Disorder (PTSD) and Complex PTSD (CPTSD), as measured by the International Trauma Questionnaire (ITQ; Cloitre et al., 2018) has been supported in many factor analytic and mixture modelling studies. There is, however, a paucity of research investigating the latent structure of the ITQ using factor mixture modelling (FMM). FMM was applied to data collected from a nationally representative sample of U.S. adults ( $N = 1834$ ). FMM results demonstrated strong support for a two-factor second-order model with four qualitatively distinct latent classes: a 'PTSD class', a 'CPTSD class', a 'DSO' (Disturbances in Self-Organisation) class and a 'low symptoms class'. Sexual abuse increased likelihood of membership to the 'CPTSD' ( $OR = 3.22$ ) and physical abuse decreased likelihood of membership to the 'PTSD' ( $OR = 0.51$ ). Trauma exposure in adulthood predicted 'PTSD' and 'CPTSD' class membership. The 'CPTSD class' was characterised by higher levels of psychopathological co-morbidities and poorer psychological wellbeing compared to all other classes. Results provide additional support for the validity of PTSD and CPTSD as measured by the ITQ.

## 1. Introduction

In the 11th version of the International Classification of Diseases (ICD-11; World Health Organization, 2018) Posttraumatic Stress Disorder (PTSD) and Complex PTSD (CPTSD) are included under the parent category of 'disorders specifically associated with stress'. Exposure to a trauma, defined as an extremely threatening or horrific event or series of events is a prerequisite for consideration of either disorder. The diagnostic criteria for PTSD consist of three symptom clusters which relate specifically to the traumatic event and were selected based on investigations indicating that these clusters best discriminated PTSD from other disorders (Brewin, Lanius, Novac, Schnyder, & Galea, 2009). These symptoms include 1) re-experiencing of the trauma in the here and now (Re), 2) avoidance of traumatic reminders (Av), and 3) heightened sense of current threat (Th). The diagnosis of CPTSD is an

adaptation of the ICD-10 "Enduring Personality Change after Catastrophic Experiences" (EPCACE) which, following from the theoretical proposal of Herman (1992), was included to recognize changes in affect, identity and relational capacities that can occur following prolonged or chronic exposure to trauma (WHO, 1999). CPTSD consists of six symptom clusters: the three PTSD clusters as well as three symptom clusters representing disturbances in self-organization (DSO), namely (1) affect dysregulation (AD), (2) negative self-concept (NSC), and (3) disturbances in relationships (DR). Significant functional impairment associated with the PTSD and DSO symptoms are required for diagnosis (WHO, 1992). The selection of the DSO symptom clusters and items was based on those symptoms identified as most frequently occurring in the DSM-IV field trials of a version of Complex PTSD "disorders of extreme Stress, not otherwise specified" (DESNOS) (see Roth, Newman, Pelcovitz, Van Der Kolk, & Mandel, 1997; Van der Kolk, Roth, Pelcovitz,

<sup>☆</sup> This work was supported by the Department for the Economy PhD studentship. The Department for the Economy had no role in the study design, the collection, analysis and interpretation of data, in writing the report and in the decision to submit the article for publication.

\* Corresponding author.

E-mail address: [m.shevlin@ulster.ac.uk](mailto:m.shevlin@ulster.ac.uk) (M. Shevlin).

<https://doi.org/10.1016/j.janxdis.2021.102497>

Received 4 June 2021; Received in revised form 13 October 2021; Accepted 1 November 2021

Available online 6 November 2021

0887-6185/© 2021 Elsevier Ltd. All rights reserved.

Sunday, & Spinazzola, 2005) as well those identified as most frequent and most impairing by expert clinicians in a consensus survey on Complex PTSD (Cloitre et al., 2011). International field trials have found substantial support among clinicians for the distinction between PTSD and CPTSD (Keeley et al., 2016) and that the addition of the CPTSD diagnosis has improved differential diagnosis across the stress related disorders relative to ICD-10 (Gaebel et al., 2020; Reed et al., 2018).

The characterisation of PTSD and CPTSD as distinct, albeit related, disorders has been supported by factor analytic research whereby symptom structures consistent with the ICD-11 distinction between PTSD and DSO have been identified and in mixture modelling research whereby distinct trauma groups representing PTSD and CPTSD have been identified (Brewin et al., 2017; Redican et al., 2021). This research has been facilitated by the development of the International Trauma Questionnaire (ITQ; Cloitre et al., 2018). Although the ITQ accommodates diagnostic and dimensional scoring, and many studies have applied categorical (mixture modelling) and dimensional (factor analysis) latent variable modelling to investigate the latent structure of PTSD and CPTSD, few studies have sought to simultaneously assess whether the latent structure of the ITQ is best represented by a categorical or dimensional model, or a combination of both. Factor analysis is a dimensional approach where continuous latent variable(s) are modelled to explain the variance and covariance of multiple observed indicators (Lubke & Muthén, 2005). Conversely, mixture modelling is a categorical approach where a categorical latent variable is modelled to separate individuals into discrete groups based on similar response patterns to observed indicators (Nylund, Asparouhov, & Muthén, 2007). Factor mixture modelling (FMM) is a hybrid model that integrates features of factor analysis and mixture modelling, and has been developed as an alternative method by which to examine the latent structure of psychological constructs (Miettunen, Nordström, Kaakinen, & Ahmed, 2016). FMM assigns individuals into discrete subgroups via a mixture modelling approach and models heterogeneity within these groups through incorporating continuous latent factors, allowing the underlying structure of a construct to be assessed at the categorical and dimensional level simultaneously (Clark et al., 2013). FMM facilitates direct comparisons of model fit across categorical, dimensional, and hybrid models thereby allowing for informed decisions to be made regarding which approach best captures the latent structure of the construct under investigation (Whalen, 2017).

Clark et al. (2013) have previously demonstrated the superiority of FMM over both factor analytic and mixture modelling in relation to conduct disorder (CD) symptomology. These researchers first identified the best fitting dimensional structure of their CD symptom data using factor analysis, identifying a single unidimensional construct of CD. They then identified the best fitting mixture model of CD symptoms using latent class analysis (LCA), identifying a three-class solution (i.e. three distinct groups characterised by the same pattern of variation across the CD symptoms). Next, the researchers tested a series of FMM's, utilising the number of classes from the optimal LCA solution (i.e. three-class model) as the upper-limit for extracting classes in the FMM, and incorporating the best-fitting FA model (i.e., a one-factor model) as the dimensional component of the FMM. Utilising standard fit indices to compare model fit across the best dimensional (i.e. FA), categorical (i.e. LCA) and hybrid (i.e. FMM) model, Clark et al. (2013) found that the FMM best captured the latent structure of CD, highlighting the utility of FMM in understanding the latent structure of psychological constructs.

Only two studies have investigated the latent structure of ICD-11 PTSD and CPTSD using FMM. Wolf et al. (2015) were the first to apply this method using two samples including a nationally representative community sample of adults living in the United States and a small veteran sample. A FMM with two latent dimensional variables and four latent classes provided the best fit to the data in both samples. It was determined that classes were quantitatively rather than qualitatively distinct, a finding inconsistent with the proposed distinction between PTSD and CPTSD in ICD-11. A second study by Frost et al. (2019)

applied FMM to investigate the latent structure of PTSD and CPTSD among a refugee sample. A correlated six-factor model with five qualitatively different latent classes was identified as the best fitting model. In contrast to Wolf et al. (2015), distinct classes reflecting the distinction between PTSD and CPTSD were found, supporting the ICD-11 model. An important limitation of these studies was that they attempted to measure the ICD-11 symptoms of PTSD and CPTSD using proxy items. Further work is therefore required that uses data collected via a reliable and valid measure of ICD-11 PTSD and CPTSD (i.e., the ITQ).

In this study, a secondary analysis of data from a nationally representative sample of adults living in the U.S. was used to test whether responses to the ITQ are best represented by a categorical, dimensional, or 'hybrid' FMM model. We proposed that if the ICD-11's distinction between PTSD and CPTSD is meaningful and valid, a hybrid model which captures the distinction between PTSD and DSO symptoms at a dimensional level, and which identifies groups of adults with symptom profiles consistent with PTSD and CPTSD at a categorical level should provide the best fit to the data, similar to Frost et al. (2019). Alternatively, if the ICD-11 model is invalid, results would be similar to those reported by Wolf et al. (2015) whereby either (a) there would be no evidence of a distinction between PTSD and DSO symptoms at the dimensional level or (b) there would be no evidence of a qualitative distinction between PTSD and CPTSD at the categorical level.

Furthermore, we aimed to investigate the role of various demographic and trauma-related risk factors in predicting latent class membership. Although the effects of ACEs on adult psychopathology are well-documented (e.g. Kalmakis & Chandler, 2015; Kessler et al., 2010; McLaughlin et al., 2017) and previous research has highlighted the influential role of adverse childhood experiences (ACEs) in the development of PTSD and CPTSD (Cloitre et al., 2019; Frewen, Zhu, & Lanius, 2019; G.W. Ho et al., 2020b,a; G.W.K. Ho et al., 2019; Jowett, Karatzias, Shevlin, & Albert, 2020; Karatzias et al., 2017a,b; Karatzias et al., 2020), the contribution of specific ACEs in predicting both disorders has been largely unexplored. Thus, we sought to examine the role of specific ACEs in predicting membership to each symptom profile. Childhood sexual and physical abuse are considered as traumatic stressors and forms of adversity (McLaughlin et al., 2017). Thus, in line with Cloitre et al. (2019) who analysed the same dataset, we categorized both events as traumatic stressors. We anticipated similar results to Cloitre et al. (2019) where childhood physical and sexual abuse significantly predicted CPTSD. We hypothesized that the CPTSD class would be characterised by the highest levels of adulthood traumatic exposure (Cloitre et al., 2019; Hyland et al., 2017; Palic et al., 2016) and higher levels of psychological comorbidities and overall poorer wellbeing (Cloitre et al., 2019; Gilbar, 2020; Hyland & Shevlin, 2018; Hyland, Ceannt et al., 2018; Karatzias & Shevlin, 2017; Karatzias et al., 2019; Karatzias, Cloitre et al., 2017).

## 2. Methods

### 2.1. Participants

The sample was a nationally representative household sample of non-institutionalised adults living in the U.S. Data were collected in 2017, using an existing online research panel for a project investigating the validity of ICD-11 PTSD and CPTSD symptom proposals (Cloitre et al., 2019). Participants were randomly recruited using a probability-based sampling strategy. Eligible participants were aged between 18 and 70 years at the time of the survey and had experienced at least one traumatic event in their lifetime. Of the 3953 individuals screened for participation in the survey, 1839 met the inclusion criteria and were invited to complete the survey (eligibility rate=46.3%). The survey design oversampled women and minority populations, each at a 2:1 ratio. To adjust for this oversampling, the data were weighted according to age/ethnic distribution of the US population (Cloitre et al., 2019; Shevlin et al., 2018). Ethical approval was obtained from the

research ethics committee at Ulster University.

## 2.2. Findings from Cloitre et al. (2019)

The present study involves the re-analysis of an existing dataset (see Cloitre et al., 2019 for a detailed overview) and much is already known about the sample characteristics. In brief, the sample had a mean age of 44.56 years (SD= 14.89) and 52% were female (n = 956). More than half of the sample were married (55.3%; n = 1016), 62.2% (n = 1143) attained a college education, and the majority (71.1%; n = 1306) were employed. Table 1 presents weighted sociodemographic characteristics of the sample. Cloitre et al. (2019) reported 7.2% (n = 128) of the sample as meeting the requirements for a diagnosis of either PTSD or CPTSD (PTSD prevalence=3.4%, CPTSD prevalence= 3.8%). Females were more likely to meet the criteria for PTSD (OR=2.53) and CPTSD (OR=1.82) compared to males. In terms of childhood traumatic stressors, 17.5% (n = 321) of the sample reported experiencing sexual abuse and 15.7% (n = 289) reported experiencing physical abuse. The average number of ACEs reported was 1.36 (SD=1.72, Range= 0–8, Median = 1.00), with the most frequently reported ACEs being parental separation or divorce (33.5%; n = 613), substance abuse in the home (24.5%; n = 450) and verbal abuse (21.3%; n = 388). The average number of traumas reported in adulthood was 2.55 (SD=2.17, Range= 0–14, Median = 2.00), with the most commonly endorsed traumas being natural disaster (39.9%, n = 725), transportation accident (55.8%, n = 1021) and the sudden or unexpected death of a loved one (51.5%, n = 943).

**Table 1**  
Sociodemographic (Weighted) Characteristics of the Sample (N = 1839).

|                                       | % (n)                 |
|---------------------------------------|-----------------------|
| <b>Sex</b>                            |                       |
| Male                                  | 48.0 (883)            |
| Female                                | 52.0 (956)            |
| <b>Age in years</b>                   |                       |
| 18–24                                 | 10.0 (184)            |
| 25–34                                 | 20.7 (382)            |
| 35–44                                 | 19.0 (350)            |
| 45–54                                 | 18.5 (339)            |
| 55–64                                 | 21.6 (398)            |
| 65 +                                  | 10.2 (187)            |
| Age                                   | M = 44.55, SD = 14.89 |
| <b>Region</b>                         |                       |
| Northeast                             | 18.1 (333)            |
| Midwest                               | 20.9 (385)            |
| South                                 | 38.2 (702)            |
| West                                  | 22.8 (420)            |
| <b>Highest educational attainment</b> |                       |
| Less than high school                 | 9.1 (168)             |
| High school                           | 28.7 (528)            |
| Some college                          | 30.3 (558)            |
| Bachelor's degree or higher           | 31.8 (585)            |
| <b>Current relationship status</b>    |                       |
| In a committed relationship           | 63.4 (1165)           |
| Not in a committed relationship       | 36.6 (674)            |
| <b>Current employment status</b>      |                       |
| Employed                              | 71.1 (1307)           |
| Unemployed                            | 28.9 (532)            |
| <b>Ethnicity</b>                      |                       |
| Non-white                             | 36.2 (666)            |
| White                                 | 63.8 (1173)           |
| <b>Income level (\$)</b>              |                       |
| 0–19,999                              | 10.8 (198)            |
| 20,000–34,999                         | 11.0 (202)            |
| 35,000–49,999                         | 12.2 (225)            |
| 50,000–74,999                         | 17.6 (322)            |
| 75,000–99,999                         | 14.4 (264)            |
| 100,000 or more                       | 34.1 (628)            |

## 2.3. Measures

### 2.3.1. ICD-11 PTSD and CPTSD

The International Trauma Questionnaire (ITQ; Cloitre et al., 2018) is a self-report measure designed to capture all elements of an ICD-11 PTSD and CPTSD diagnosis. The ITQ first assesses a person's most distressing traumatic event, and how long ago the event occurred. There are six items measuring the three PTSD symptom clusters (Re, Av and Th) and six items measuring the three DSO symptom clusters (AD, NSC, DR). Three questions enquire about the extent to which the PTSD and DSO symptoms, respectively, have affected relationships, work and other important domains of functioning. Participants rate how often they have been bothered by each of the symptoms in the past month using a five-point Likert scale ranging from 'Not at all' (0) to 'Extremely' (4). To satisfy the criteria for a diagnosis of PTSD, one of two items from each PTSD symptom cluster and at least one functional impairment item must be endorsed (endorsement is based on a Likert score of  $\geq 2$  (i.e., 'Moderately'). To qualify for a diagnosis of CPTSD, criteria for PTSD must be satisfied in addition to the endorsement of one of two items from each DSO symptom cluster and at least one functional impairment item (endorsement is Likert score  $\geq 2$ ). An individual can receive a diagnosis of PTSD or CPTSD, but not both. Cronbach's alpha for the PTSD ( $\alpha = .89$ ) and DSO ( $\alpha = .89$ ) sub-scales in the current study were excellent.

### 2.3.2. Childhood Traumatic Events, Adversities and Adulthood trauma

The Adverse Childhood Experiences Questionnaire (ACE; Felitti et al., 1998) was used to measure exposure to traumatic stressors and adversities in childhood. The ACE comprises ten questions measuring exposure to adverse experiences in the first eighteen years of life, and uses a yes (1) or no (0) response format. Two ACE items (childhood sexual and physical abuse) were categorized as traumatic stressors and the remaining items were categorized as childhood adversities. A revised version of the Life Events Checklist (LEC-R; Gray, Litz, Hsu, & Lombardo, 2004) was used to measure trauma exposure in adulthood. Participants were asked to indicate whether they had been exposed to 14 different traumatic events after the age of eighteen years, using yes (1) or no (0) responses. Responses were summed to create a total adulthood trauma score. Cronbach's alpha for the ACE items ( $\alpha = .77$ ) and adult LEC-R items ( $\alpha = .69$ ) were satisfactory.

### 2.3.3. Major Depressive Disorder and Generalized Anxiety Disorder

The Patient Health Questionnaire (PHQ-8; Kroenke et al., 2009) is an eight-item self-report measure of depression symptoms and the Generalized Anxiety Disorder Scale (GAD-7; Spitzer, Kroenke, Williams, & Löwe, 2006) is a seven-item scale used to assess anxiety symptoms. For the PHQ-8, respondents indicate the number of the days over the past two weeks that they have been bothered by a particular depressive symptom using a four-point Likert scale ranging from 0 (not at all) to 3 (nearly every-day). Total scores range between 0 and 24, with higher scores indicative of greater symptom severity. A PHQ-8 score  $\geq 10$  indicates clinically significant depression (Kroenke & Spitzer, 2002). For the GAD-7, participants indicate on a 4-point Likert scale the degree to which they have been bothered by symptoms over the last two weeks, with responses ranging from 0 (not at all) to 3 (nearly every day). A GAD-7 score  $\geq 10$  can be used as a cut-off point for identifying cases of GAD (Spitzer et al., 2006). The internal reliability of the PHQ-8 ( $\alpha = .93$ ) and GAD-7 ( $\alpha = .94$ ) in this study were excellent.

### 2.3.4. Psychological Wellbeing

The World Health Organisation Well-being Index (WHO-5, (WHO, 1998) is a five-item scale measuring subjective wellbeing, and comprises five positively phrased items including "cheerful and in good spirits" and "felt calm and relaxed". Participants indicate the extent to which each of the five statements has applied to them over the past two weeks. The WHO-5 uses a six-point Likert scale, ranging from 0 (at no time) to 5 (all of the time). Raw scores range from 0 to 25, with higher scores

indicating a higher level of psychological wellbeing. Cronbach's alpha of the WHO-5 in the current study was excellent ( $\alpha = .93$ ).

### 2.3.5. Demographic variables

Demographic variables assessed included age (years), gender (0 = male, 1 = female), educational status (0 = no college, 1 = college), relationship status (0 = not in committed relationship, 1 = in a committed relationship), employment status (0 = unemployed, 1 = employed), ethnicity (0 = non-white, 1 = white) and household income (measured in categories ranging from 0 = less than \$5000 to 21 = \$250,000 or more).

## 2.4. Statistical Analysis

FMM was conducted in three consecutive steps as per Clark et al. (2013). First, three alternative CFA models were tested: **Model 1** was a one-factor model where all PTSD (Re, Av, Th) and DSO (AD, NSC, DR) symptoms loaded onto a first-order CPTSD factor; **Model 2** was a correlated six-factor model (Re, Av, Th, AD, NSC, DR); and **Model 3** was a two-factor second-order model where Av, Re and Th loaded onto the second-order 'PTSD' factor, and, AD, NSC and DR load onto the second-order 'DSO' factor. To determine the best-fitting model, a number of fit indices were assessed: the chi-square statistic, the comparative fit index (CFI; Bentler, 1990), the Tucker-Lewis index (TLI; Tucker & Lewis, 1973), the root mean square error of approximation (RMSEA; Browne & Cudeck, 1992; Steiger, 1990) and the Standardized Root Mean Square Residual (SRMR; Jöreskog & Sörbom, 1981). Standard cut-off criteria were used to determine model fit (Hu & Bentler, 1999) with a non-significant  $\chi^2$  value ( $p > .05$ ) indicating good fit; CFI and TLI values  $\geq .90$  and  $\geq .95$  considered as good and excellent model fit, respectively; SRMR values  $\leq 0.8$  indicating good fit; RMSEA values  $< .05$  indicating close fit and  $< .08$  indicating adequate fit (Steiger, 1990). In addition, Bayesian Information Criterion (BIC; Sclove, 1987), sample size adjusted BIC (ssaBIC; Sclove, 1987) and Akaike Information Criterion (AIC) were used to assess relative fit with lower values indicative of superior model fit. The model with the lowest BIC was considered to be the best model, with differences  $\geq 10$  being considered strong evidence for the selection of the lower BIC model (Raftery, 1995).

Second, a latent profile analysis (LPA) on the item level data was conducted, testing models with two to six latent classes. The relative fit of the solutions were assessed using information criterion statistics and the Lo-Mendell-Rubin adjusted likelihood ratio test (LMR-A; Lo, Mendell, & Rubin, 2001). A non-significant LMR-A indicates that there is no statistically significant improvement in fit with the inclusion of an additional class, and thus the more parsimonious model should be chosen (Nylund et al., 2007). It is often the case for LPA that as the number of latent classes increases, the information theory-based statistics fail to reach a single smallest value (Masyn, 2013, p. 572). Thus, diminishing gains in model fit were explored using 'elbow plots' to identify the point where changes in fit values begin to plateau, which in itself is indicative of minimal and non-significant gains in information (Masyn, 2013; Nylund et al., 2007). Entropy values, a measure of classification certainty, were also inspected with higher values indicative of greater classification accuracy (Lubke & Muthén, 2007).

The third step involved fitting a series of FMMs to the data. The number of classes from the best-fitting LPA model were used as the upper-limit for extracting classes in the FMM (Clark et al., 2013). There are several variations of the FMM, each with different restrictions to identify the model (Clark et al., 2013; Lubke & Muthén, 2007). For the current study, a variation of 'type-1' FMMs were employed which are typically characterised by class-varying factor means, class-invariant item intercepts, class-invariant factor loadings and a factor covariance matrix fixed at zero (Clark et al., 2013). However, rather than allowing factor means to vary, the current study estimated item-level intercepts to assess the performance of the individual ITQ symptom indicators across the latent classes. Factor means were fixed at zero to achieve model

identification. To avoid solutions based on local maxima, 500 random sets of starting values were used in the initial stage, and 100 optimizations were used in the final stage of convergence. These values were increased for more complex models to ensure replication of the best log-likelihood value. In addition to using the same model fit criteria as used for LPA, profile plots were inspected to assess the substantive interpretability of the classes comprising each model (Nylund et al., 2007). A chi-square test was conducted to determine the degree to which the variables representing most likely class membership corresponded with probable diagnostic status (criteria not met, PTSD criteria met, CPTSD criteria met). Observed and expected counts for each latent class with respect to diagnostic status were examined and the magnitude of the differences between observed and expected values were substantiated by assessing adjusted standardized residuals with values  $> 1.96$  indicating a statistically significant difference.

The final stage of the analyses involved investigating predictors of the latent classes. The demographic variables, childhood sexual and physical abuse, eight ACEs, and total adult LEC were entered into the model in two stages using the R3step auxiliary command in Mplus (Asparouhov & Muthén, 2014a, 2014b). The R3step method is similar to multinomial logistic regression model (Vermunt, 2010) and ensures that covariates do not influence latent class formation whilst also recognising classification uncertainty (Asparouhov & Muthén, 2014a, 2014b). The first model included all covariates separately to examine the bivariate associations between covariates and latent class membership. In the second model, all covariates were added to the model simultaneously to assess whether any covariate was uniquely associated with latent class membership, over and beyond the effects of the other demographic and trauma-related predictors included in the model. Following this, differences across the latent classes with regards to mean scores on the GAD-7, PHQ-8 and WHO-5 were examined using the Bolck-Croon-Hagenaars Method (BCH method; Bolck, Croon, & Hagenaars, 2004). This method (analogous to four separate one-way ANOVAs) has been shown to be a robust method for investigating the relationship between class membership, covariates and distal outcomes (Bakk and Vermunt, 2016). For each distal outcome a Wald chi-square test was used to test the null hypothesis that the means were equal across all classes (akin to post-hoc tests following ANOVA); if this null hypothesis was rejected, pairwise comparisons were conducted to determine where the significant differences occurred.

All analyses were conducted using Mplus version 8.2 (Muthén & Muthén, 2017), with initial descriptive statistics computed using SPSS version 27 (IBM Corp. Released, 2020). All models were estimated using robust maximum likelihood estimation (MLR; Yuan & Bentler, 2000). For the covariate analyses, listwise deletion was used which is the default when using the auxiliary procedure in Mplus.

## 3. Results

Goodness of fit statistics for the CFA, LPA and FMM models are reported in Table 2.

### 3.1. CFA Results

Model 1 was rejected due to poor model fit. Model 2 ( $\chi^2(39) = 83.104, p < .001, RMSEA = .025, CFI = .993, TLI = .987$ ) and Model 3 ( $\chi^2(66) = 101.624, p < .001, RMSEA = .025, CFI = .991, TLI = .987$ ) demonstrated excellent fit. The chi-squared statistic was significant for both models; however this should not lead to rejection of these models as the power of chi-square tests is positively related to sample size (Tanaka, 1987). The difference in BIC values between Model 2 (correlated six-factor) and Model 3 (two-factor second-order) exceeded ten points ( $\Delta BIC = 19.25$ ), and thus the second-order model was selected as the optimal model. All items loaded significantly ( $p < .001$ ) and strongly ( $> .74$ ) onto their respective first-order PTSD and DSO

**Table 2**  
Fit Statistics for the CFA, LCA and FMM of ICD-11 PTSD and CPTSD.

| Model                      | Log-likelihood    | AIC              | BIC              | ssaBIC           | Entropy      | LMR-A (p)    |
|----------------------------|-------------------|------------------|------------------|------------------|--------------|--------------|
| CFA                        |                   |                  |                  |                  |              |              |
| Model 1                    | -26497.891        | 53067.782        | 53266.295        | 53151.924        | –            | –            |
| Model 2                    | -24307.505        | 48717.010        | 48998.237        | 48836.212        | –            | –            |
| <b>Model 3</b>             | <b>-24327.939</b> | <b>48741.877</b> | <b>48978.990</b> | <b>48842.381</b> | –            | –            |
| LPA                        |                   |                  |                  |                  |              |              |
| 2 classes                  | -27193.751        | 54461.501        | 54665.529        | 54547.981        | .974         | .0000        |
| 3 classes                  | -26117.401        | 52334.802        | 52610.515        | 52451.667        | .946         | .2366        |
| <b>4 classes</b>           | <b>-25274.137</b> | <b>50674.274</b> | <b>51021.672</b> | <b>50821.523</b> | <b>.940</b>  | <b>.5142</b> |
| 5 classes                  | -24832.832        | 49817.663        | 50236.747        | 49995.297        | .938         | .7761        |
| 6 classes                  | -24384.670        | 48947.340        | 49438.109        | 49155.359        | .938         | .7588        |
| FMM                        |                   |                  |                  |                  |              |              |
| 2 factors 2 classes        | -23599.779        | 47311.558        | 47620.356        | 47442.446        | 0.990        | .0000        |
| 2 factors 3 classes        | -23358.178        | 46854.355        | 47234.839        | 47015.628        | 0.979        | .7047        |
| <b>2 factors 4 classes</b> | <b>-22966.634</b> | <b>46097.268</b> | <b>46549.437</b> | <b>46288.926</b> | <b>0.986</b> | <b>.7398</b> |

Note: AIC = Akaike Information Criterion, BIC = Bayesian Information Criterion, ssaBIC = sample size adjusted Bayesian Information Criterion, LMR-A Lo-Mendell-Rubin adjusted likelihood ratio test. Best-fitting models for each approach (CFA, LPA, FMM) shown in bold.

factors, with the exception of one affective dysregulation item ‘takes a long time to calm down’, which had a relatively weaker factor loading (.55). All first-order PTSD (Re, Av, Th) and DSO (AD, NSC, DR) factors loaded strongly onto their respective second-order factors (all >.81). The correlation between the PTSD and DSO latent factors was .70.

### 3.2. LPA Results

The LMR-A was significant for the two-class solution only, however, the loglikelihood and BIC values continued to decrease with each additional class. ‘Elbow’ plots demonstrated how increases in loglikelihood values were minimal after the three-class solution whilst increases in BIC were minimal after the four-class solution. Given that BIC is superior to other fit indices in identifying the correct number of latent classes with the use of continuous indicators (Nylund et al., 2007) and because the size of each class was adequate with the smallest class comprising 9% of the sample (Nylund-Gibson & Choi, 2018), the four-class solution was selected as the best model.

### 3.3. FMM Results

BIC values were lower for the FMMs compared to the CFA and LPA models, which indicated the superior fit of the FMMs. LMR-A was significant for the two-class model only, however, the log-likelihood was highest and information criterion statistics (i.e. BIC, ssaBIC and AIC) were substantially lower for the four-class model. Entropy increased from the three-class to the four-class solution, indicating improved classification certainty from the three to four-class solution. Inspection of profile plots and class compositions for each solution demonstrated the classes comprising the four-class solution to be most interpretable and thus, the four-class second-order FMM was selected as the best-fitting model.

The profile plot (Fig. 1) presents the ITQ item endorsement patterns as influenced by the continuous latent factors derived from the CFA model across the four latent classes. Class 1 (78.9%,  $n = 1447$ ) was characterised by low symptom severity across all symptom indicators of the PTSD and DSO factors, and was thus labelled ‘low symptoms’. Class 2 (7.1%,  $n = 131$ ) was labelled ‘DSO class’, as individuals in this class reported lower severity on all PTSD factor indicators but greater symptom severity on all DSO indicators, particularly the NSC item “I feel worthless”. Class 3 (6.0%,  $n = 109$ ) was characterised by high symptom severity on all PTSD and DSO item indicators, particularly the Th item ‘jumpy or easily startled’ and both NSC items, and was therefore labelled ‘CPTSD class’. Finally, class 4 (8.0%,  $n = 147$ ) was characterised by lower overall symptom severity across all symptom indicators compared to class 3, but higher symptom severity on the PTSD items compared to class 1 and class 2, and lower symptom severity on the DSO indicators

compared to class 2. This class was also characterised by high endorsement of the Th item ‘jumpy or easily startled’ and as a result, this class was labelled the ‘PTSD class’.

### 3.4. Agreement between ITQ classification and FMM classes

A chi-square test between variables representing most likely class membership and probable diagnostic status was conducted, with results demonstrating a significant association, ( $6, N = 1735$ ) = 132.419,  $p < .001$ . There was a high degree of correspondence between the two variables. The adjusted standardized residuals with respect to probable diagnostic status revealed that those in the ‘no diagnosis group’ were most likely to be in the ‘low symptoms’ class (adjusted standardised residual = 6.6), individuals in the ‘CPTSD diagnosis’ group were most likely to be in the ‘CPTSD class’ (adjusted standardised residual = 10.0) and individuals in the ‘PTSD diagnosis’ group were most likely to be in the ‘PTSD class’ (adjusted standardised residual = 4.8).

### 3.5. R3step (covariate) analyses

A series of bivariate analyses were conducted using the R3step procedure. The reference class for all analyses was ‘low symptoms’. Sexual abuse was found to be a significant predictor of all classes compared to the reference class and both the ‘PTSD class’ and ‘CPTSD class’ but not the ‘DSO class’ were significantly predicted by physical abuse, with these effects being strongest for the ‘CPTSD class’. In terms of ACEs, all classes were predicted by physical neglect, emotional neglect and verbal abuse, with these effects being strongest for the ‘CPTSD class’. Compared to the reference class, all classes were significantly predicted by substance abuse and mental illness in the home whilst the ‘PTSD’ and ‘CPTSD’ classes were predicted by household violence and incarceration of a family member. Table 3.

As demonstrated in Table 4, the second step involved calculating adjusted odds ratios for demographic and trauma-related predictors of class membership. Compared to the ‘low symptoms’, individuals in the ‘DSO class’ (OR=0.973), ‘CPTSD class’ (OR=0.971) and ‘PTSD class’ (OR=0.964) were likelier to be younger. Unique demographic predictors of the ‘DSO class’ included not being in a committed relationship (OR=0.421) and having no college education (OR=0.509). Membership of the ‘PTSD class’ (OR=1.161) and ‘CPTSD class’ (OR=1.190) was predicted by higher adulthood LEC scores. Individuals in the ‘CPTSD class’ were approximately three times more likely to report sexual abuse (OR=3.22) and surprisingly, those in the ‘PTSD class’ were less likely to report physical abuse (OR= 0.505) relative to the reference class. In terms of ACEs, individuals in the ‘DSO’ class were approximately three times more likely than the reference class to report emotional neglect (OR=3.70) whilst conversely, individuals in the ‘CPTSD class’ were less




Fig. 1. Profile Plot of Estimated Means for 4-class Factor Mixture Model Solution.

Table 3  
Unadjusted Odds Ratios for Demographic and Trauma-Related Predictors of Latent Class Membership.

| Predictor   | Class 2: DSO<br>OR (95% CI) |                | Class 3: CPTSD<br>OR (95% CI) |                 | Class 4: PTSD<br>OR (95% CI) |                 |
|---|-----------------------------|----------------|-------------------------------|-----------------|------------------------------|-----------------|
| Childhood Sexual Abuse                            | 2.223*                      | (1.369, 3.611) | 7.028*                        | (4.401, 11.223) | 2.067*                       | (1.318, 3.240)  |
| Childhood Physical Abuse                          | 1.978                       | (1.176, 3.327) | 6.961*                        | (4.332, 11.185) | 2.191*                       | (1.332, 3.603)  |
| Physical neglect (ACE 3)                          | 2.876**                     | (1.305, 6.337) | 3.888*                        | (1.927, 7.841)  | 6.531**                      | (3.543, 12.041) |
| Verbal abuse (ACE 4)                              | 3.784**                     | (2.399, 5.968) | 7.614*                        | (4.770, 12.152) | 4.069**                      | (2.584, 6.406)  |
| Emotional neglect (ACE 5)                         | 5.974**                     | (3.777, 9.449) | 8.853*                        | (5.490, 14.276) | 4.824**                      | (3.007, 7.738)  |
| Violence in the home (ACE 6)                      | 1.575                       | (0.829, 2.992) | 4.914*                        | (2.914, 8.286)  | 4.014*                       | (2.354, 6.842)  |
| Parental separation or divorce(ACE 7)             | 1.487                       | (0.954, 2.319) | 1.253                         | (0.792, 1.983)  | 1.872                        | (1.212, 2.893)  |
| Substance abuse in the home (ACE 8)               | 1.866*                      | (1.183, 2.941) | 5.551**                       | (3.495, 8.816)  | 2.700*                       | (1.733, 4.208)  |
| Mental illness in the home (ACE 9)                | 2.634*                      | (1.600, 4.337) | 4.898*                        | (3.034, 7.906)  | 2.413**                      | (1.446, 4.027)  |
| Household member incarnated(ACE10)                | 1.525                       | (0.756, 3.075) | 2.893*                        | (1.651, 5.071)  | 3.488*                       | (1.837, 6.473)  |
| Gender  | 1.591                       | (1.024, 2.472) | 1.342                         | (0.899, 2.005)  | 2.760                        | (1.322, 5.717)  |
| LEC Adult Total Score                             | 1.055                       | (0.959, 1.162) | 1.333*                        | (1.219, 1.458)  | 1.186*                       | (1.091, 1.289)  |
| Age   | 0.973**                     | (0.959, 0.989) | 0.989                         | (0.974, 1.004)  | 0.968**                      | (0.955, 0.982)  |
| Ethnicity   | 0.922                       | (0.592, 1.437) | 0.675*                        | (0.434, 1.050)  | 0.682*                       | (0.442, 1.052)  |
| Relationship Status (in a committed relationship) | 0.363**                     | (0.233, 0.565) | 0.710                         | (0.447, 1.128)  | 0.709                        | (0.457, 1.098)  |
| Employment status (employed)                      | 0.810                       | (0.521, 1.259) | 0.577**                       | (0.366, 0.911)  | 1.467                        | (0.881, 2.441)  |
| Educational Status (college education)            | 0.616*                      | (0.396, 0.958) | 0.675                         | (0.428, 1.065)* | 0.722                        | (0.463, 1.126)  |
| Household income                                  | 0.936*                      | (0.894, 0.980) | 0.905                         | (0.863, 0.949)  | 0.947                        | (0.910, 0.986)  |

Note: Class 1 (Baseline) is the reference category,

\* - significant at  $p < 0.05$ ,

\*\* - significant at  $p < 0.01$ .

likely to report physical neglect (OR=0.417) and parental separation or divorce (OR=0.448) relative to those in 'low symptoms'.

### 3.6. BCH (distal outcomes) Results

Sum scores for the PHQ-8, GAD-7 and WHO-5 were added into the model as distal outcomes using the BCH method (see Table 5). For the overall sample, the average GAD-7 score was 3.61 (SD= 4.84) and the average PHQ-8 scores 4.08 (SD= 5.40) with 13.8% (n = 253) and 11.2% (n = 199) of the sample exceeding the cut-off scores indicative of depression and generalised anxiety respectively. The mean WHO-5 score was 14.99 (SD=6.35). With regards to GAD-7 scores, pairwise comparisons between classes using  $\chi^2$  showed that GAD-7 scores were significantly higher for the 'CPTSD class' compared to the 'PTSD class' whilst there was no significant difference in mean scores between the 'DSO class' and the 'PTSD class'. With regards to PHQ-8 scores, the mean

score was highest for the 'CPTSD class' followed by the 'DSO class' and 'PTSD class'. Results from the  $\chi^2$  analysis demonstrated how the average level of depression symptomology was significantly lower for the reference class compared to all other classes. Additionally, the average level of depression symptomology for the 'CPTSD class' was significantly higher than the 'DSO class' which in turn was identified as being significantly higher than the 'PTSD class'. The mean WHO-5 score was highest for the 'low symptoms' class, followed by the 'PTSD class' and 'DSO class' whilst the mean level of psychological wellbeing was lowest for the 'CPTSD class'. Results from the  $\chi^2$  analysis demonstrated how the average WHO-5 score was significantly lower for the 'CPTSD class' compared to all other classes.

## 4. Discussion

This was the first study to investigate the latent structure of ICD-11

**Table 4**  
Demographic and Trauma-Related Predictors and Latent Class Membership (adjusted odds ratios).

| Predict                               | Class 2: DSO<br>OR (95% CI) |                | Class 3: CPTSD<br>OR (95% CI) |                | Class 4: PTSD<br>OR (95% CI) |                |
|---------------------------------------|-----------------------------|----------------|-------------------------------|----------------|------------------------------|----------------|
| Age                                   | 0.973**                     | (0.956, 0.990) | 0.971**                       | (0.951, 0.991) | 0.964**                      | (0.945, 0.982) |
| Gender                                | 1.560                       | (0.908, 2.680) | 1.324                         | (0.762, 2.300) | 1.439                        | (0.868, 2.386) |
| Household Income                      | 0.987                       | (0.932, 1.045) | 0.965                         | (0.912, 1.022) | 0.972                        | (0.925, 1.021) |
| Education Status                      | 0.509**                     | (0.316, 0.822) | 0.754                         | (0.399, 1.423) | 0.831                        | (0.503, 1.372) |
| Relationship Status                   | 0.421**                     | (0.254, 0.698) | 1.120                         | (0.611, 2.053) | 0.941                        | (0.554, 1.598) |
| Employment Status                     | 1.141                       | (0.690, 1.885) | 0.732                         | (0.402, 1.332) | 1.937                        | (1.176, 3.189) |
| Ethnicity                             | 1.255                       | (0.747, 2.107) | 0.888                         | (0.515, 1.531) | 0.938                        | (0.581, 1.515) |
| Childhood Sexual Abuse                | 1.504                       | (0.847, 2.688) | 3.221*                        | (1.847, 5.617) | 1.271                        | (0.728, 2.216) |
| Childhood Physical Abuse              | 0.891                       | (0.466, 1.703) | 1.779                         | (0.849, 3.731) | 0.505*                       | (0.224, 1.138) |
| Physical neglect (ACE 3)              | 0.950                       | (0.331, 2.728) | 0.417**                       | (0.174, 0.999) | 2.179                        | (0.962, 4.937) |
| Verbal abuse (ACE 4)                  | 1.971                       | (1.041, 3.731) | 1.867                         | (0.875, 3.982) | 2.121                        | (1.058, 4.254) |
| Emotional neglect (ACE 5)             | 3.700*                      | (1.873, 7.310) | 2.322                         | (1.151, 4.684) | 2.156                        | (1.172, 3.967) |
| Violence in the home (ACE 6)          | 0.593                       | (0.286, 1.228) | 1.211                         | (0.555, 2.642) | 1.849                        | (0.848, 4.031) |
| Parental separation or divorce(ACE 7) | 0.909                       | (0.543, 1.523) | 0.448**                       | (0.251, 0.799) | 0.883                        | (0.544, 1.433) |
| Substance abuse in the home (ACE 8)   | 1.210                       | (0.713, 2.053) | 2.321                         | (1.248, 4.317) | 1.240                        | (0.712, 2.158) |
| Mental illness in the home (ACE 9)    | 1.743                       | (0.975, 3.118) | 2.295                         | (1.242, 4.243) | 1.038                        | (0.573, 1.880) |
| Household member incarnated(ACE10)    | 0.767                       | (0.312, 1.886) | 0.880                         | (0.397, 1.952) | 1.547                        | (0.657, 3.640) |
| Total LEC Adult Score                 | 1.032                       | (0.925, 1.150) | 1.190**                       | (1.058, 1.339) | 1.161*                       | (1.037, 1.301) |

Note: Class 1 (Baseline) is the reference category,

\* - significant at p < 0.05,

\*\* - significant at p < 0.01.

**Table 5**  
Equality test of means of PHQ-8, GAD-7 and WHO-5 scores across the latent classes.

|                                 | Class 1: Low symptoms |         | Class 2: DSO |         | Class 3: CPSTD |         | Class 4: PTSD |         | Overall Chi-Square test<br>(p < 0.01) | Pairwise comparison<br>(p < .05) |
|---------------------------------|-----------------------|---------|--------------|---------|----------------|---------|---------------|---------|---------------------------------------|----------------------------------|
|                                 | Mean                  | (se)    | Mean         | (se)    | Mean           | (se)    | Mean          | (se)    |                                       |                                  |
| Depression (PHQ-8)              | 2.355                 | (0.102) | 10.158       | (0.795) | 14.106         | (0.681) | 7.840         | (0.709) | 444.287<br>(p < 0.01)                 | 2, 3, 4 > 1<br>3 > 2, 4<br>4 > 2 |
| Anxiety (GAD-7)                 | 2.098                 | (0.096) | 7.804        | (0.609) | 12.265         | (0.649) | 8.036         | (0.634) | 408.194<br>(p < 0.01)                 | 2, 3, 4 > 1<br>3 > 2, 4          |
| Psychological Wellbeing (WHO-5) | 16.504                | (0.176) | 8.902        | (0.636) | 6.958          | (0.522) | 11.315        | (0.651) | 445.262<br>(p < 0.01)                 | 2, 3, 4 < 1<br>2 > 3<br>4 > 2, 3 |

PTSD and CPTSD as measured by the ITQ in a general population sample of U.S. adults using FMM. Furthermore, we sought to explore demographic and trauma-related predictors of the identified symptom profiles as well as assessing differences across classes in terms of comorbid psychological disorders and wellbeing.

Consistent with prior research (Frost et al., 2019; Wolf et al., 2015), a hybrid model was found to best capture the latent structure of PTSD and CPTSD. Specifically, a two-factor second-order model with four qualitatively different latent classes was identified as the best-fitting model. The superior fit of the hierarchical CFA model corresponds with findings from much of the research conducted to date (Redican et al., 2021), and aligns with the ICD-11 conceptualisation of CPTSD as comprising of two distinct dimensions of PTSD and DSO (Brewin et al., 2017). Notably, similar to a small number of prior studies (Haselgruber et al., 2020; Vang et al., 2021), the affective dysregulation item representing hyperactivation loaded weakly onto the affective dysregulation factor in the present study. Although the hyperactivation and hypoactivation items capture alternative facets of affective dysregulation, the representation of both items as manifestations of a shared underlying construct (i.e. affective dysregulation) has been supported within the literature (Karatzias, Hyland, Ben-Ezra, & Shevlin, 2018). The emergence of four qualitatively different symptom profiles is consistent with findings from other general population studies where ‘PTSD’, ‘CPTSD’, ‘DSO’ and ‘low symptoms’ classes were also identified (Ben-Ezra et al., 2018; Kazlauskas et al., 2020; Tian et al., 2020; Rink and Lipinska, 2020). Altogether, our findings contradict those of Wolf et al. (2015) as our results demonstrate that even when differences between PTSD and DSO symptoms are acknowledged at the dimensional level, there is evidence

of qualitatively distinct trauma populations at the categorical level, and that these groups reflect the distinction between PTSD and CPTSD. These findings provide support for the ICD-11 predictions that there are distinct trauma groups in the population. The diagnostic status of some individuals did not accurately reflect their latent class membership (i.e., 17% (n = 273) of participants in the no diagnosis, PTSD diagnosis or CPTSD diagnosis groups), the results revealed that there was a significantly greater proportion of individuals in the ‘low symptoms’, ‘PTSD’ and ‘CPTSD’ classes with the correct corresponding diagnostic status than expected, which supports the diagnostic accuracy of the ‘PTSD’ and ‘CPTSD’ symptom profiles. These results add to the growing body of research establishing the ITQ as a valid measure of PTSD and CPTSD (Redican et al., 2021).

Similar to findings from existing research (e.g. Cloitre et al., 2019; Hyland et al., 2017), sexual abuse significantly predicted membership of the ‘CPTSD class’, concurring with the ICD-11 theorization that childhood sexual abuse increases likelihood of developing CPTSD (Maercker et al., 2013). Surprisingly, and in contrast to Cloitre et al. (2019), our results indicated that those in the ‘PTSD class’ were less likely to report physical abuse compared to the reference class. This divergent finding may have resulted from the different analytic procedure adopted in the current study such that both events were analysed in conjunction with other ACEs whereas Cloitre et al. (2019) analysed both events in the presence of other childhood traumatic events, not ACEs, using the revised version of the Life Events Checklist (LEC; Gray et al., 2004).

ACEs are recognised as significant risk-factors for all posttraumatic stress responses (Cloitre et al., 2019; Frewen et al., 2019; Ho et al., 2020; Karatzias et al., 2020). Indeed, bivariate results showed a strong

association between all ACEs measured in the current study, with the exception of parental separation or divorce, and the PTSD and CPTSD symptom profiles when compared to the non-symptomatic group. Notably, when considered within the multivariate context, only a small number of associations remained statistically significant. Given the established role of shared variance in explaining the effects of childhood maltreatment on mental health outcomes (Cecil, Viding, Fearon, Glaser, & McCrory, 2017), it is unsurprising that many of the associations became non-statistically significant when considered in the context of other traumas and adversities. Contradictory to existing research where physical neglect increased the risk of CPTSD (e.g. Choi, Lee, & Hyland, 2021; Gilbar, Hyland, Cloitre, & Dekel, 2018; Karatzias & Shevlin, 2017; Karatzias, Cloitre et al., 2017), multivariate results showed how individuals in the 'CPTSD' were less likely to report physical neglect despite there being a positive bivariate association. It is possible that the shared variance between physical neglect and CPTSD is fully accounted for by the other covariates included in the multivariate model. Alternatively, Knefel et al. (2019) highlighted how the PTSD criteria of re-experiencing and avoidance may not be particularly relevant to those who have experienced neglect and similarly, Sölva, Haselgruber, and Lueger-Schuster (2020) in their investigation of the different profiles of childhood adversities amongst foster children, reported how the 'high neglect' group experienced lowest CPTSD symptom severity. When considered in the presence of other more interpersonal traumas and adversities, it is possible that that physical neglect may decrease risk of membership to the 'CPTSD class'.

Similar to findings from Cloitre et al. (2019), adulthood traumatic exposure significantly predicted membership of both the 'PTSD class' and 'CPTSD class'. However, Cloitre et al. (2019) utilised an aggregate trauma variables representing different levels of trauma exposure (i.e. 0, 1, 2, 3 or  $\geq 4$ ), and found that the risk of CPTSD relative to PTSD only increased at the highest level of adulthood trauma exposure (i.e.  $\geq 4$  traumas). Conversely, the current study assessed trauma utilising a sum score reflecting the total number of traumas to which an individual was exposed, with the effect of cumulative adulthood trauma exposure being strongest for CPTSD. Collectively, these results demonstrate why the type of trauma is a risk factor for either condition, but not a prerequisite for the diagnosis. As per previous studies (Ben-Ezra et al., 2018; Ho, Bressington et al., 2020; Karatzias et al., 2019; Perkonig et al., 2016), age significantly predicted PTSD and CPTSD symptomology, thereby indicating how maladaptive posttraumatic stress responses tend to be endorsed predominantly by younger age groups.

Our results demonstrated that the 'CPTSD class' reported the highest levels of depression and anxiety symptomology, and the lowest levels of overall psychological wellbeing, which is consistent with previous research (Elklit, Hyland, & Shevlin, 2014; Hyland & Shevlin, 2018; Hyland & Shevlin, 2018; Hyland, Ceannt et al., 2018; Hyland, Ceannt et al., 2018; Karatzias et al., 2019). Similar to existing research (Gilbar, 2020; Hyland et al., 2017; Ho et al., 2020), those in the 'DSO class' reported higher levels of depression symptomology compared to the 'PTSD class' whilst those in the 'PTSD class' reported a higher mean anxiety score compared to those in the 'DSO class', although this difference was not statistically significant. The high degree of comorbidity between disorders may be explained by the fact that the symptoms constituting DSO are cross-diagnostic. For example, emotional dysregulation is also a central component of many anxiety and mood disorders (Dvir, Ford, Hill, & Frazier, 2014) and the symptoms reflecting negative self-concept, which were heavily endorsed by both the 'DSO class' and 'CPTSD class', are also features of major depressive disorder (Zahn et al., 2015). Conversely, the co-occurrence of depression and anxiety disorders with posttraumatic stress responses can also be considered in the context of the 'Hierarchical Taxonomy of Psychopathology' (HiTop; Kotov et al., 2017) in which PTSD falls under the 'internalising dimension' and thus should be most strongly correlated with other disorders within the same dimension such as depressive and anxiety disorders (Fox, Hyland, Power, & Coogan, 2020). There are a myriad of other potential

explanations for the high levels of comorbidity observed such as a shared vulnerability across disorders in terms of risk factors (Spinhoven, Penninx, Van Hemert, De Rooij, & Elzinga, 2014), genetic disposition as well as similarities in cognitive and memory processes (Angelakis & Nixon, 2015). Further research is required to unpack the patterns of comorbidity observed among PTSD/CPTSD and other forms of psychopathology.

With regard to the identification of the 'DSO class', various proposals have been made to elucidate what exactly this class may represent: (1) it may represent individuals with other psychological disorders (Cloitre et al., 2020; Knefel, Garvert, Cloitre, & Lueger-Schuster, 2015), (2) it may be a function of the larger sample sizes typical of studies analysing data from large samples of the general population, which tend to extract more classes or (3) it may reflect individuals with subthreshold CPTSD who possess a greater vulnerability to the symptoms constituting DSO but greater resilience to PTSD symptoms (Perkonig et al., 2016). Given the high levels of depression symptomology, poorer psychological wellbeing and high endorsement of NSC items in the 'DSO class' in the current study, it is plausible that this class may reflect individuals with other psychiatric disorders which can also occur post-trauma (Cloitre et al., 2020; Knefel et al., 2015; Perkonig et al., 2016). Individuals in the 'DSO class' were also approximately three times more likely to report emotional neglect, an adversity linked to internalising symptoms in adulthood (Cohen, Menon, Shorey, Le, & Temple, 2017). Further investigation of the 'DSO class' is necessary to understand what exactly this class represents, especially in general population samples (Cloitre et al., 2020).

Several study limitations are worth noting. Although the sample was a large nationally representative general-population sample of U.S. adults, the prevalence of PTSD and CPTSD was relatively low thereby limiting generalisability of findings to highly traumatised samples. Although clinically meaningful and distinct symptom profiles were identified, the relatively low endorsement of PTSD and DSO symptoms in the current study resulted in an uneven distribution of participants across the latent classes with the majority of participants belonging to the 'low symptoms' class. There is clearly a need for further research in traumatised clinical samples who are likely to present with higher rates of PTSD and CPTSD. Future research conducted on clinical samples is therefore recommended. There are also limitations associated with the ACE measure utilised in the current study such as the lack of consensus regarding which events can be considered as ACEs and the dichotomous yes/no nature of this scale prohibits examination of the frequency, intensity or chronicity of ACE exposure (Anda, Porter, & Brown, 2020). Thus, it is possible that the unanticipated findings surrounding the effects of physical abuse and neglect may be explained by the implementation of this particular measure. Finally, the cross-sectional nature of the study does not allow for inferences regarding causality to be made.

Overall, this study provides further support for the validity of PTSD and CPTSD, as defined by ICD-11, in a general population sample of US adults through the application of FMM. Establishing risk factors that distinguish these disorders from each other has important implications for the diagnosis, treatment and prevention of both disorders (Cloitre, 2020). Thus, the identification of risk factors specific to and shared across the various symptom profiles in the present study affords a comprehensive understanding of the aetiological risk factors associated with the various posttraumatic responses. In saying that, several unexpected findings were observed in terms of the role of physical abuse and neglect in predicting posttraumatic responses, thus determining whether such findings replicate across other samples is an important avenue of exploration for future studies.

#### Declarations of Interest

Dr Marylene Cloitre was a member of the World Health Organisation Working Group on the classification of disorders specifically associated



with stress, reporting to the International Advisory Group for the revision of ICD-10 Mental and Behavioural Disorders. Professor Mark Shevlin, Professor Philip Hyland and Professor Thanos Karatzias were part of the International Trauma Questionnaire development team and authors on the ITQ validation paper. The views provided in this article are the opinions of the authors and do not represent WHO policy.

## Acknowledgements

This work was supported by a Department for the Economy PhD studentship. The Department for the Economy had no role in the study design, the collection, analysis and interpretation of data, in writing the report and in the decision to submit the article for publication.

## References

- Anda, R. F., Porter, L. E., & Brown, D. W. (2020). Inside the adverse childhood experience score: Strengths, limitations, and misapplications. *American Journal of Preventive Medicine*, 59(2), 293–295.
- Angelakis, S., & Nixon, R. D. (2015). The comorbidity of PTSD and MDD: Implications for clinical practice and future research. *Behaviour Change*, 32(1), 1–25.
- Asparouhov, T., & Muthén, B. (2014aaa). Auxiliary variables in mixture modeling: Using the BCH method in Mplus to estimate a distal outcome model and an arbitrary secondary model. *Mplus Web Notes*, 21(2), 1–22.
- Asparouhov, T., & Muthén, B. (2014bbb). Auxiliary variables in mixture modelling: Three-step approaches using M plus. *Structural Equation Modelling: A Multidisciplinary Journal*, 21(3), 329–341.
- Bakk, Z., & Vermunt, J. K. (2016). Robustness of stepwise latent class modeling with continuous distal outcomes. *Structural Equation Modeling: A Multidisciplinary Journal*, 23(1), 20–31.
- Ben-Ezra, M., Karatzias, T., Hyland, P., Brewin, C. R., Cloitre, M., Bisson, J. I., & Shevlin, M. (2018). Posttraumatic stress disorder (PTSD) and complex PTSD (CPTSD) as per ICD-11 proposals: A population study in Israel. *Depression and Anxiety*, 35(3), 264–274.
- Bentler, P. M. (1990). Comparative fit indexes in structural models. *Psychological Bulletin*, 107(2), 238.
- Bolck, A., Croon, M., & Hagenaars, J. (2004). Estimating latent structure models with categorical variables: One-step versus three-step estimators. *Political Analysis*, 3–27.
- Brewin, C. R., Cloitre, M., Hyland, P., Shevlin, M., Maercker, A., Bryant, R. A., & Somasundaram, D. (2017). A review of current evidence regarding the ICD-11 proposals for diagnosing PTSD and complex PTSD. *Clinical Psychology Review*, 58, 1–15.
- Brewin, C. R., Lanius, R. A., Novac, A., Schnyder, U., & Galea, S. (2009). Reformulating PTSD for DSM-V: life after criterion A. *Journal of Traumatic Stress*, 22(5), 366–373.
- Browne, M. W., & Cudeck, R. (1992). Alternative ways of assessing model fit. *Sociological Methods & Research*, 21(2), 230–258.
- Cecil, C. A., Viding, E., Fearon, P., Glaser, D., & McCrory, E. J. (2017). Disentangling the mental health impact of childhood abuse and neglect. *Child Abuse & Neglect*, 63, 106–119.
- Choi, H., Lee, W., & Hyland, P. (2021). Factor structure and symptom classes of ICD-11 complex posttraumatic stress disorder in a South Korean general population sample with adverse childhood experiences. *Child Abuse & Neglect*, 114, Article 104982.
- Clark, S. L., Muthén, B., Kaprio, J., D'Onofrio, B. M., Viken, R., & Rose, R. J. (2013). Models and strategies for factor mixture analysis: An example concerning the structure underlying psychological disorders. *Structural Equation Modelling: a Multidisciplinary Journal*, 20(4), 681–703.
- Cloitre, M. (2020). ICD-11 complex post-traumatic stress disorder: simplifying diagnosis in trauma populations. *The British Journal of Psychiatry*, 216(3), 129–131.
- Cloitre, M., Brewin, C., Bisson, J., Hyland, P., Karatzias, T., Lueger-Schuster, B., et al. (2020). Evidence for the coherence and integrity of the complex PTSD (CPTSD) diagnosis: response to Achterhof et al., (2019) and Ford (2020). *European Journal Of Psychotraumatology*, 11(1), 1739873.
- Cloitre, M., Courtois, C. A., Charuvastra, A., Carapezza, R., Stolbach, B. C., & Green, B. L. (2011). Treatment of complex PTSD: Results of the ISTSS expert clinical survey on best practices. *Journal of Traumatic Stress*, 24(6), 615–627.
- Cloitre, M., Hyland, P., Bisson, J. I., Brewin, C. R., Roberts, N. P., Karatzias, T., & Shevlin, M. (2019). ICD-11 posttraumatic stress disorder and complex posttraumatic stress disorder in the United States: A population-based study. *Journal of Traumatic Stress*, 32(6), 833–842.
- Cloitre, M., Shevlin, M., Brewin, C. R., Bisson, J. I., Roberts, N. P., Maercker, A., & Hyland, P. (2018). The International Trauma Questionnaire: development of a self-report measure of ICD-11 PTSD and complex PTSD. *Acta Psychiatrica Scandinavica*, 138(6), 536–546.
- Cohen, J. R., Menon, S. V., Shorey, R. C., Le, V. D., & Temple, J. R. (2017). The distal consequences of physical and emotional neglect in emerging adults: A person-centered, multi-wave, longitudinal study. *Child Abuse & Neglect*, 63, 151–161.
- Dvir, Y., Ford, J. D., Hill, M., & Frazier, J. A. (2014). Childhood maltreatment, emotional dysregulation, and psychiatric comorbidities. *Harvard Review of Psychiatry*, 22(3), 149.
- Elklit, A., Hyland, P., & Shevlin, M. (2014). Evidence of symptom profiles consistent with posttraumatic stress disorder and complex posttraumatic stress disorder in different trauma samples. *European Journal of Psychotraumatology*, 5(1), 24221.
- Felitti, V. J., Anda, R. F., Nordenberg, D., Williamson, D. F., Spitz, A. M., Edwards, V., & Marks, J. S. (1998). Relationship of childhood abuse and household dysfunction to many of the leading causes of death in adults: The Adverse Childhood Experiences (ACE) Study. *American Journal of Preventive Medicine*, 14(4), 245–258.
- Fox, R., Hyland, P., Power, J. M., & Coogan, A. N. (2020). Patterns of comorbidity associated with ICD-11 PTSD among older adults in the United States. *Psychiatry Research*, 290, Article 113171.
- Frewen, P., Zhu, J., & Lanius, R. (2019). Lifetime traumatic stressors and adverse childhood experiences uniquely predict concurrent PTSD, complex PTSD, and dissociative subtype of PTSD symptoms whereas recent adult non-traumatic stressors do not: Results from an online survey study. *European Journal of Psychotraumatology*, 10(1), 1606625.
- Frost, R., Hyland, P., McCarthy, A., Halpin, R., Shevlin, M., & Murphy, J. (2019). The complexity of trauma exposure and response: Profiling PTSD and CPTSD among a refugee sample. *Psychological Trauma: Theory, Research, Practice, and Policy*, 11(2), 165.
- Gaebel, W., Stricker, J., Riesbeck, M., Zielasek, J., Kerst, A., Meisenzahl-Lechner, E., & Falkai, P. (2020). Accuracy of diagnostic classification and clinical utility assessment of ICD-11 compared to ICD-10 in 10 mental disorders: findings from a web-based field study. *European Archives of Psychiatry and Clinical Neuroscience*, 270(3), 281–289.
- Gilbar, O. (2020). Examining the boundaries between ICD-11 PTSD/CPTSD and depression and anxiety symptoms: A network analysis perspective. *Journal of Affective Disorders*, 262, 429–439.
- Gilbar, O., Hyland, P., Cloitre, M., & Dekel, R. (2018). ICD-11 complex PTSD among Israeli male perpetrators of intimate partner violence: Construct validity and risk factors. *Journal of Anxiety Disorders*, 54, 49–56.
- Gray, M. J., Litz, B. T., Hsu, J. L., & Lombardo, T. W. (2004). Psychometric properties of the life events checklist. *Assessment*, 11, 330–341. doi.org/10.1177/1073191104269954.
- Haselgruber, A., Sölvä, K., & Lueger-Schuster, B. (2020). Validation of ICD-11 PTSD and complex PTSD in foster children using the International Trauma Questionnaire. *Acta Psychiatrica Scandinavica*, 141(1), 60–73.
- Herman, J. L. (1992). Complex PTSD: A syndrome in survivors of prolonged and repeated trauma. *Journal of Traumatic Stress*, 5(3), 377–391.
- Ho, G. W., Bressington, D., Karatzias, T., Chien, W. T., Inoue, S., Yang, P. J., & Hyland, P. (2020b). Patterns of exposure to adverse childhood experiences and their associations with mental health: a survey of 1346 university students in East Asia. *Social Psychiatry and Psychiatric Epidemiology*, 55(3), 339–349.
- Ho, G. W., Hyland, P., Shevlin, M., Chien, W. T., Inoue, S., Yang, P. J., & Karatzias, T. (2020a). The validity of ICD-11 PTSD and Complex PTSD in East Asian cultures: findings with young adults from China, Hong Kong, Japan, and Taiwan. *European Journal of Psychotraumatology*, 11(1), 1717826.
- Ho, G. W. K., Chan, A. C. Y., Shevlin, M., Karatzias, T., Chan, P. S., & Leung, D. (2019). Childhood Adversity, Resilience, and Mental Health: A Sequential Mixed-Methods Study of Chinese Young Adults. *Journal of Interpersonal Violence*. https://doi.org/10.1177/0886260519876034
- Hu, L. T., & Bentler, P. M. (1999). Cutoff criteria for fit indexes in covariance structure analysis: Conventional criteria versus new alternatives. *Structural Equation Modeling: A Multidisciplinary Journal*, 6(1), 1–55.
- Hyland, P., Ceannat, R., Daccache, F., Abou Daher, R., Sleiman, J., Gilmore, B., & Vallières, F. (2018a). Are posttraumatic stress disorder (PTSD) and complex-PTSD distinguishable within a treatment-seeking sample of Syrian refugees living in Lebanon? *Global Mental Health*, 5.
- Hyland, P., Murphy, J., Shevlin, M., Vallières, F., McElroy, E., Elklit, A., & Cloitre, M. (2017). Variation in post-traumatic response: The role of trauma type in predicting ICD-11 PTSD and CPTSD symptoms. *Social Psychiatry and Psychiatric Epidemiology*, 52(6), 727–736.
- Hyland, P., Shevlin, M., Fyvie, C., & Karatzias, T. (2018b). Posttraumatic stress disorder and complex posttraumatic stress disorder in DSM-5 and ICD-11: clinical and behavioral correlates. *Journal of Traumatic Stress*, 31(2), 174–180.
- IBM Corp. Released. (2020). *IBM SPSS Statistics for Windows, Version 27.0*. Armonk, NY: IBM Corp.
- Jöreskog, K. G., & Sörbom, D. (1981). *LISREL V: Analysis of linear structural relationships by maximum likelihood and least squares methods*. University of Uppsala. Department of Statistics.
- Jowett, S., Karatzias, T., Shevlin, M., & Albert, I. (2020). Differentiating symptom profiles of ICD-11 PTSD, complex PTSD, and borderline personality disorder: A latent class analysis in a multiply traumatized sample. *Personality Disorders: Theory, Research, and Treatment*, 11(1), 36.
- Kalmakis, K. A., & Chandler, G. E. (2015). Health consequences of adverse childhood experiences: A systematic review. *Journal of the American Association of Nurse Practitioners*, 27(8), 457–465.
- Karatzias, T., Cloitre, M., Maercker, A., Kazlauskas, E., Shevlin, M., Hyland, P., & Brewin, C. R. (2017a). PTSD and Complex PTSD: ICD-11 updates on concept and measurement in the UK, USA, Germany and Lithuania. *European Journal of Psychotraumatology*, 8(sup7), 1418103.
- Karatzias, T., Hyland, P., Ben-Ezra, M., & Shevlin, M. (2018). Hyperactivation and hypoactivation affective dysregulation symptoms are integral in complex posttraumatic stress disorder: Results from a nonclinical Israeli sample. *International Journal of Methods in Psychiatric Research*, 27(4).
- Karatzias, T., Hyland, P., Bradley, A., Cloitre, M., Roberts, N. P., Bisson, J. I., & Shevlin, M. (2019). Risk factors and comorbidity of ICD-11 PTSD and complex PTSD:

- Findings from a trauma-exposed population-based sample of adults in the United Kingdom. *Depression and Anxiety*, 36(9), 887–894.
- Karatzias, T., Shevlin, M., Fyvie, C., Hyland, P., Efthymiadou, E., Wilson, D., & Cloitre, M. (2017b). Evidence of distinct profiles of posttraumatic stress disorder (PTSD) and complex posttraumatic stress disorder (CPTSD) based on the new ICD-11 trauma questionnaire (ICD-TQ). *Journal of Affective Disorders*, 207, 181–187.
- Karatzias, T., Shevlin, M., Hyland, P., Fyvie, C., Grandison, G., & Ben-Ezra, M. (2020). ICD-11 posttraumatic stress disorder, complex PTSD and adjustment disorder: the importance of stressors and traumatic life events. *Anxiety, Stress, & Coping*, 1–12.
- Kazlauskas, E., Zelviene, P., Daniunaite, I., Hyland, P., Kvedaraitė, M., Shevlin, M., & Cloitre, M. (2020). The structure of ICD-11 PTSD and Complex PTSD in adolescents exposed to potentially traumatic experiences. *Journal of Affective Disorders*, 265, 169–174.
- Keeley, J. W., Reed, G. M., Roberts, M. C., Evans, S. C., Robles, R., Matsumoto, C., & Maercker, A. (2016). Disorders specifically associated with stress: A case-controlled field study for ICD-11 mental and behavioural disorders. *International Journal of Clinical and Health Psychology*, 16(2), 109–127.
- Kessler, R. C., McLaughlin, K. A., Green, J. G., Gruber, M. J., Sampson, N. A., Zaslavsky, A. M., & Williams, D. R. (2010). Childhood adversities and adult psychopathology in the WHO World Mental Health Surveys. *The British Journal of Psychiatry*, 197(5), 378–385.
- Knefel, M., Garvert, D. W., Cloitre, M., & Lueger-Schuster, B. (2015). Update to an evaluation of ICD-11 PTSD and complex PTSD criteria in a sample of adult survivors of childhood institutional abuse by Knefel & Lueger-Schuster (2013): A latent profile analysis. *European Journal of Psychotraumatology*, 6(1), 25290.
- Van der Kolk, B. A., Roth, S., Pelcovitz, D., Sunday, S., & Spinazzola, J. (2005). Disorders of extreme stress: The empirical foundation of a complex adaptation to trauma. *Journal of Traumatic Stress: Official Publication of the International Society for Traumatic Stress Studies*, 18(5), 389–399.
- Knefel, M., Lueger-Schuster, B., Karatzias, T., Shevlin, M., & Hyland, P. (2019). From child maltreatment to ICD-11 complex post-traumatic stress symptoms: The role of emotion regulation and re-victimisation. *Journal of Clinical Psychology*, 75(3), 392–403.
- Kotov, R., Krueger, R. F., Watson, D., Achenbach, T. M., Althoff, R. R., Bagby, R. M., & Zimmerman, M. (2017). The Hierarchical Taxonomy of Psychopathology (HiTOP): A dimensional alternative to traditional nosologies. *Journal of Abnormal Psychology*, 126(4), 454.
- Kroenke, K., & Spitzer, R. L. (2002). The PHQ-9: a new depression diagnostic and severity measure. *Psychiatric Annals*, 32(9), 509–515.
- Kroenke, K., Strine, T. W., Spitzer, R. L., Williams, J. B., Berry, J. T., & Mokdad, A. H. (2009). The PHQ-8 as a measure of current depression in the general population. *Journal of Affective Disorders*, 114(1–3), 163–173.
- Lo, Y., Mendell, N. R., & Rubin, D. B. (2001). Testing the number of components in a normal mixture. *Biometrika*, 88(3), 767–778.
- Lubke, G., & Muthén, B. O. (2007). Performance of factor mixture models as a function of model size, covariate effects, and class-specific parameters. *Structural Equation Modeling*, 14(1), 26–47.
- Lubke, G. H., & Muthén, B. (2005). Investigating population heterogeneity with factor mixture models. *Psychological Methods*, 10(1), 21.
- Maercker, A., Brewin, C. R., Bryant, R. A., Cloitre, M., van Ommeren, M., Jones, L. M., & Somasundaram, D. J. (2013). Diagnosis and classification of disorders specifically associated with stress: proposals for ICD-11. *World Psychiatry*, 12(3), 198–206.
- Masyn, K. E. (2013). Latent class analysis and finite mixture modeling. In T. D. Little (Ed.), *The Oxford handbook of quantitative methods* (volume 2). Oxford: University Press.
- McLaughlin, K. A., Koenen, K. C., Bromet, E. J., Karam, E. G., Liu, H., Petukhova, M., & Kessler, R. C. (2017). Childhood adversities and post-traumatic stress disorder: evidence for stress sensitisation in the World Mental Health Surveys. *The British Journal of Psychiatry*, 211(5), 280–288.
- Miettunen, J., Nordström, T., Kaakinen, M., & Ahmed, A. (2016). Latent variable mixture modeling in psychiatric research – a review and application. *Psychological Medicine*, 46(3), 457–467. <https://doi.org/10.1017/S0033291715002305>
- Muthén, L. K., & Muthén, B. O. (2017). 1998–2017. Mplus User's Guide. Los Angeles, CA: Muthén & Muthén.
- Nylund, K. L., Asparouhov, T., & Muthén, B. O. (2007). Deciding on the number of classes in latent class analysis and growth mixture modeling: A Monte Carlo simulation study. *Structural Equation Modelling: A Multidisciplinary Journal*, 14(4), 535–569.
- Nylund-Gibson, K., & Choi, A. Y. (2018). Ten frequently asked questions about latent class analysis. *Translational Issues in Psychological Science*, 4(4), 440.
- Palic, S., Zerach, G., Shevlin, M., Zeligman, Z., Elklit, A., & Solomon, Z. (2016). Evidence of complex posttraumatic stress disorder (CPTSD) across populations with prolonged trauma of varying interpersonal intensity and ages of exposure. *Psychiatry Research*, 246, 692–699.
- Perkonig, A., Höfler, M., Cloitre, M., Wittchen, H. U., Trautmann, S., & Maercker, A. (2016). Evidence for two different ICD-11 posttraumatic stress disorders in a community sample of adolescents and young adults. *European Archives of Psychiatry and Clinical Neuroscience*, 266(4), 317–328.
- Raftery, A. E. (1995). Bayesian model selection in social research. *Sociological Methodology*, 111–163.
- Redican, E., Nolan, E., Hyland, P., Cloitre, M., McBride, O., Karatzias, T., & Shevlin, M. (2021). A Systematic Literature Review of Factor Analytic and Mixture Models of ICD-11 PTSD and CPTSD using the International Trauma Questionnaire. *Journal of Anxiety Disorders*, Article 102381.
- Reed, G. M., Sharan, P., Rebello, T. J., Keeley, J. W., Elena Medina-Mora, M., Gureje, O., & Pike, K. M. (2018). The ICD-11 developmental field study of reliability of diagnoses of high-burden mental disorders: results among adult patients in mental health settings of 13 countries. *World Psychiatry*, 17(2), 174–186.
- Rink, J., & Lipinska, G. (2020). Evidence of distinct profiles of ICD-11 post-traumatic stress disorder (PTSD) and complex PTSD in a South African sample. *European Journal of Psychotraumatology*, 11(1), 1818965.
- Roth, S., Newman, E., Pelcovitz, D., Van Der Kolk, B., & Mandel, F. S. (1997). Complex PTSD in victims exposed to sexual and physical abuse: Results from the DSM-IV field trial for posttraumatic stress disorder. *Journal of Traumatic Stress*, 10(4), 539–555.
- Sclove, S. L. (1987). Application of model-selection criteria to some problems in multivariate analysis. *Psychometrika*, 52(3), 333–343.
- Shevlin, M., Hyland, P., Roberts, N. P., Bisson, J. I., Brewin, C. R., & Cloitre, M. (2018). A psychometric assessment of Disturbances in Self-Organization symptom indicators for ICD-11 Complex PTSD using the International Trauma Questionnaire. *European Journal of Psychotraumatology*, 9(1), 1419749.
- Sölvä, K., Haselgruber, A., & Lueger-Schuster, B. (2020). Latent classes of childhood maltreatment in children and adolescents in foster care: associations with ICD-11 PTSD and complex PTSD. *European Journal of Psychotraumatology*, 11(1), 1832757.
- Spinhoven, P., Penninx, B. W., Van Hemert, A. M., De Rooij, M., & Elzinga, B. M. (2014). Comorbidity of PTSD in anxiety and depressive disorders: Prevalence and shared risk factors. *Child Abuse & Neglect*, 38(8), 1320–1330.
- Spitzer, R. L., Kroenke, K., Williams, J. B., & Löwe, B. (2006). A brief measure for assessing generalized anxiety disorder: the GAD-7. *Archives of Internal Medicine*, 166(10), 1092–1097.
- Steiger, J. H. (1990). Structural model evaluation and modification: An interval estimation approach. *Multivariate Behavioral Research*, 25(2), 173–180.
- Tian, Y., Wu, X., Wang, W., Zhang, D., Yu, Q., & Zhao, X. (2020). Complex posttraumatic stress disorder in Chinese young adults using the International Trauma Questionnaire (ITQ): A latent profile analysis. *Journal of Affective Disorders*, 267, 137–143.
- Tucker, L. R., & Lewis, C. (1973). A reliability coefficient for maximum likelihood factor analysis. *Psychometrika*, 38(1), 1–10.
- Vang, M. L., Dokkedahl, S. B., Løkkegaard, S. S., Jakobsen, A. V., Møller, L., Auning-Hansen, M. A., & Elklit, A. (2021). Validation of ICD-11 PTSD and DSO using the International Trauma Questionnaire in five clinical samples recruited in Denmark. *European Journal of Psychotraumatology*, 12(1), 1894806.
- Vermunt, J. K. (2010). Latent class modelling with covariates: Two improved three-step approaches. *Political analysis*, 450–469.
- Whalen, D. J. (2017). Using hybrid modeling to determine the latent structure of psychopathology. *Biological Psychiatry*, 81(6), e41–e42.
- Wolf, E. J., Miller, M. W., Kilpatrick, D., Resnick, H. S., Badour, C. L., Marx, B. P., & Friedman, M. J. (2015). ICD-11 complex PTSD in US national and veteran samples: Prevalence and structural associations with PTSD. *Clinical Psychological Science*, 3(2), 215–229.
- World Health Organization. (1992). The ICD-10 classification of mental and behavioural disorders: Clinical descriptions and diagnostic guidelines.
- World Health Organization: Regional Office for Europe (1998). Wellbeing measures in primary health care: The DepCare Project. Consensus meeting, Stockholm.
- World Health Organization. (1999). *Classification of mental and behavioural disorders, 10th revision*. Zagreb: Medicinska naklada.
- World Health Organization. (2018). International Statistical Classification of Diseases and Related Health Problems. Retrieved from (11th Revision). (<https://www.who.int/classifications/icd/en/>).
- Yuan, K. H., & Bentler, P. M. (2000). 5. Three likelihood-based methods for mean and covariance structure analysis with nonnormal missing data. *Sociological Methodology*, 30(1), 165–200.
- Zahn, R., Lythe, K. E., Gethin, J. A., Green, S., Deakin, J. F. W., Young, A. H., & Moll, J. (2015). The role of self-blame and worthlessness in the psychopathology of major depressive disorder. *Journal of Affective Disorders*, 186, 337–341.