

Role of the ancillary ligands on the stabilization of the imino-oxo tautomer of 1-methylcytosine in Pt^{II} complexes†

Bruno Longato,^{*a} Diego Montagner^a and Ennio Zangrando^b

Received 20th October 2008, Accepted 24th December 2008


First published as an Advance Article on the web 12th February 2009

DOI: 10.1039/b818514j

The mixed nucleobases complexes *cis*-[L₂Pt{1-MeTy(-H)}(1-MeCy,N³)]NO₃ (L = PPh₃, **1a**; PMePh₂, **1b**), containing the N(3)-deprotonated 1-methylthymine (1-MeTy(-H)) and the neutral 1-methylcytosine (1-MeCy) have been prepared and characterised. The compounds were obtained by reacting the hydroxo complexes *cis*-[L₂Pt(μ-OH)]₂(NO₃)₂ with 1-methylthymine (1-MeTy), followed by the addition of 1 equivalent of 1-MeCy. In solution of DMSO, DMF or chlorinated solvents, **1a** converts quantitatively into the isomer *cis*-[L₂Pt{1-MeTy(-H)}(1-MeCy,N⁴)]NO₃ (**2a**) containing the tautomeric form of the cytosine stabilized through the coordination at the N(4) atom, as shown by single-crystal X-ray analysis. The structural determination of **2a** shows the presence in the unit cell of two crystallographic independent complexes having similar conformation, with a different orientation of the two nucleobases (head–head and head–tail) according to the presence of both isomers in solution. Complex **1b**, having the less hindered PMePh₂ ligands, in DMSO solution, contains the tautomeric forms of the cytosine in equilibrium and the migration of the metal from the N(3) to N(4) site occurs only to a minor extent.

Introduction

The usual metal binding site of the model nucleobase 1-methylcytosine (1-MeCy) is the N(3) atom.¹ However, the stabilization of the tautomeric form of this molecule through the coordination at the exocyclic N(4), in particular at Pt^{II} centers, have been well documented.² In all these cases, the initial coordination of the metal occurs at the N(3) site, followed by the metal migration at the N(4) position. Such isomerisation implies the shift of one of the exocyclic NH₂ protons to the endocyclic N(3) atom of the cytosine ligand, as shown in Scheme 1.³



Scheme 1

^aDipartimento di Scienze Chimiche, Università di Padova, Via Marzolo 1, 35131, Padova, Italy. E-mail: bruno.longato@unipd.it; Tel: +39 049 8275197; Fax: +39 049 8275161

^bDipartimento di Scienze Chimiche, Università di Trieste, Via Giorgieri 1, 34127, Trieste, Italy

† Electronic supplementary information (ESI) available: ¹⁵N–¹H HMBC spectrum of **1b**. CCDC reference number 705857 for compound **2a**. For ESI and crystallographic data in CIF or other electronic format see DOI: 10.1039/b818514j

We have recently shown that the N(3)-bonded cytosine molecule in the mixed nucleobases complex *cis*-[(PPh₃)₂Pt{1-MeTy(-H)}(1-MeCy,N³)]⁺ (**1a**) slowly rearranges into its more stable tautomeric derivative *cis*-[(PPh₃)₂Pt{1-MeTy(-H)}(1-MeCy,N⁴)]⁺ (**2a**).⁴

In this paper we report the structural characterisation of this compound by single-crystal X-ray analysis showing that the binding mode of the cytosine, previously established in solution by multinuclear NMR techniques, is maintained in the solid state. Moreover, the role of the PPh₃ ligands in the stabilization of the cytosine imino-oxo tautomer has been further investigated by preparing the complex *cis*-[(PMePh₂)₂Pt{1-MeTy(-H)}(1-MeCy,N³)]⁺, **1b**, containing the less hindered PMePh₂ ligands. This simple change of the metal coordination sphere strongly effects the relative stability of the tautomeric forms of the cytosine ligand since the platination at the N(3) site appears largely predominant in a DMSO solution of **1b**.

Experimental

Synthesis and materials

cis-[(PMePh₂)₂Pt(μ-OH)]₂(NO₃)₂⁵ and 1-MeCy⁶ were prepared as previously reported. *cis*-[(PPh₃)₂Pt{1-MeTy(-H)}(1-MeCy,N⁴)]NO₃ was prepared as described in reference [4] and the sample for X-ray analysis was obtained by slow diffusion of Et₂O into a DMF solution of the compound, at room temperature. 1-MeTy and all the solvents (CH₂Cl₂, DMF, DMSO-*d*₆, CDCl₃, Et₂O) were Aldrich products.

***cis*-(PMePh₂)₂Pt{1-MeTy(-H)}(ONO₂)**. To a solution of *cis*-[(PMePh₂)₂Pt(μ-OH)]₂(NO₃)₂ (48.6 mg, 3.6·10⁻² mmol) in CH₂Cl₂ (3 cm³) 1-MeTy (11.0 mg, 7.8·10⁻² mmol) was added, and the suspension stirred at room temperature for ca. 12 h. Addition of pentane (25 cm³) to the resulting solution afforded a white

solid which was isolated and dried under vacuum. Purification of the solid from CHCl_3 , by vapour diffusion of Et_2O at room temperature, afforded crystals having the composition *cis*-(PMePh_2)₂Pt{1-MeTy(-H)}(ONO₂)·1H₂O·1/4CH₂Cl₂. (50.4 mg, 85%). (Found: C, 46.46; H, 4.02; N, 5.00. C_{32.25}H_{35.5}N₃Cl_{0.5}O₆P₂Pt requires C, 46.34; H, 4.29; N, 5.02). δ_{H} (300.13 MHz; CDCl₃; Me₄Si) 7.69–7.23 (20H, cm, PPh₂), 6.38 (1H, s, H6), 3.03 (3H, s, NCH₃), 1.89 (3H, d, ²J_{HP} 11.6, PMe), 1.80 (3H, d, ²J_{HP} 11.6, PMe), 1.65 (3H, s, CCH₃); δ_{H} (300.13 MHz; DMSO-*d*₆; Me₄Si) 7.64–7.25 (20H, cm, PPh₂), 6.84 (1H, s, H6), 2.96 (3H, s, NCH₃), 2.09 (3H, d, ²J_{HP} 12, PMe), 1.75 (3H, d, ²J_{HP} 12, PMe), 1.50 s (3H, s, CCH₃). δ_{P} (121.5 MHz; CDCl₃) AB multiplet –5.32 (1P, d, ¹J_{PPt} 3284), –13.74 (1P, d, ¹J_{PPt} 4062, ²J_{PP} 23.1). δ_{P} (121.5 MHz; DMSO-*d*₆) AB multiplet –2.58 (1P, d, ¹J_{PPt} 3241), –11.17 (1P, d, ¹J_{PPt} 4108, ²J_{PP} = 24.3).

***cis*-(PMePh_2)₂Pt{1-MeTy(-H)}(1-MeCy, N³)[NO₃] (1b).** To a solution of *cis*-(PMePh_2)₂Pt{1-MeTy(-H)}(ONO₂) (31 mg, 3.8·10⁻² mmol) in 3 cm³ of CH₂Cl₂ was added 1-MeCy (5.0 mg, 3.9·10⁻² mmol) which, under stirring at room temperature, dissolved in a few h. Addition of Et₂O afforded a white solid that after filtration, washing with Et₂O and dried under vacuum had the composition *cis*-(PMePh_2)₂Pt{1-MeTy(-H)}(1-MeCy, N³)[NO₃]·1H₂O·1/4CH₂Cl₂ (24.9 mg, 71%). (Found: C, 46.12; H, 4.08; N, 8.44. C_{37.25}H_{42.5}N₆O₇Cl_{0.5}P₂Pt requires C, 46.55; H, 4.47; N, 8.74). δ_{H} (300.13 MHz; CDCl₃; Me₄Si) 7.86–7.21 (20H, cm, PPh₂), 1.34 (3H, d, ²J_{HP} 11, PCH₃), 1.32 (3H, d, ²J_{HP} 11, PCH₃). Main conformer (1-MeTy(-H) resonances): 6.31 (1H, s, H6), 2.99 (3H, s, NCH₃), 1.56 (3H, s, CCH₃); (1-MeCy resonances): 8.69 (1H, br s, NH), 8.60 (1H, br s, NH), 6.63 (1H, d, J 7.1, H6), 6.06 (1H, d, J 7.1, H5), 2.98 (3H, s, NCH₃). Minor conformer: (1-MeTy(-H) resonances): 6.27 (1H, s, H6), 2.85 (3H, s, NCH₃), 1.58 (3H, s, CCH₃), (1-MeCy resonances): 8.75 (1H, br s, NH), 8.63 (1H, br s, NH), 6.61 (1H, d, J 7.2, H6), 6.10 (1H, d, J 7.2, H5), 3.02 (3H, s, NCH₃). δ_{H} (300.13 MHz; DMSO-*d*₆; Me₄Si) 8.01–7.39 (20H, cm, PPh₂), 1.64 (3H, d, ²J_{HP} 10.3, PCH₃), 1.63 (3H, d, ²J_{HP} 10.3, PCH₃). Main conformer (1-MeTy(-H) resonances): 6.72 (1H, s, H6), 2.89 (3H, s, NCH₃), 1.41 (3H, s, CCH₃); (1-MeCy resonances): 8.75 (1H, br s, NH), 8.37 (1H, br s, NH), 7.21 (1H, d, J 7.1, H6), 5.42 (1H, d, J 7.1, H5), 2.93 (3H, s, NCH₃). Minor conformer: (1-MeTy(-H) resonances): 6.68 (1H, s, H6), 2.89 (3H, s, NCH₃), 1.38 (3H, s, CCH₃), (1-MeCy resonances): 8.68 (1H, br s, NH), 8.37 (1H, br s, NH), 7.21 (1H, d, J 7.1, H6), 5.42 (1H, d, J 7.1, H5), 2.93 (3H, s, NCH₃). δ_{P} (121.5 MHz; CDCl₃), minor conformer: AB multiplet –11.48 (1P, d, ¹J_{PPt} 3249), –13.17 (1P, d, ¹J_{PPt} 3489, ²J_{PP} 24.4); main conformer: AB multiplet –11.75 (1P, d, ¹J_{PPt} 3249), –13.17 (1P, d, ¹J_{PPt} 3485, ²J_{PP} 24.4). δ_{P} (121.5 MHz; DMSO-*d*₆), minor conformer: AB multiplet –9.83 (1P, d, ¹J_{PPt} 3224), –11.82 (1P, d, ¹J_{PPt} 3495, ²J_{PP} 24.7); main conformer: AB multiplet –10.11 (1P, d, ¹J_{PPt} 3224), –11.88 (1P, d, ¹J_{PPt} 3495, ²J_{PP} 24.3).

NMR measurements

NMR spectra were obtained in solution of various solvents at 298 K, with a Bruker AVANCE 300 MHz for ¹H and ³¹P (operating at 300.13 and 121.5 MHz, respectively) and a Bruker 400 AMX-WB spectrometer for ¹⁵N (operating at 40.6 MHz). δ are in ppm and J in Hz. The ¹H chemical shifts were referenced to the residual impurity of the solvent and to Me₄Si. The external refer-

ences were H₃PO₄ (85 w/w in D₂O) for ³¹P, and CH₃NO₂ (in CDCl₃ at 50% w/w) for ¹⁵N. Inverse detected spectra were obtained through heteronuclear multiple bond correlation (HMBC) experiments, using parameters similar to those previously reported.⁷

X-Ray structure determination

Diffraction data for compound **2a** were collected at room temperature on a Nonius DIP-1030H system with Mo-K α radiation ($\lambda = 0.71073 \text{ \AA}$). Cell refinement, indexing and scaling of the data set were carried out using programs Denzo⁸ and Scalepack.⁸ The structure was solved by direct method and subsequent Fourier analyses⁹ and refined by the full-matrix least-squares method based on F^2 with all observed reflections.⁹ A residual in the ΔF map was interpreted as a lattice water oxygen (hydrogen atoms not located). All the calculations were performed using the WinGX System, Ver 1.70.01.¹⁰

Crystal data of **2a**·0.5(H₂O): C₄₇H₄₅N₆O_{6.50}P₂Pt, M = 1054.92, triclinic, space group $P\bar{1}$, $a = 14.868(4)$, $b = 17.155(4)$, $c = 22.120(5) \text{ \AA}$, $\alpha = 105.94(2)$, $\beta = 96.01(3)$, $\gamma = 112.59(3)^\circ$, $V = 4866(2) \text{ \AA}^3$, $Z = 4$, $D_c = 1.440 \text{ g/cm}^3$, $\mu(\text{Mo-K}\alpha) = 3.002 \text{ mm}^{-1}$, $F(000) = 2116$, θ range = $1.53\text{--}25.35^\circ$. Final $R1 = 0.0521$, $wR2 = 0.1262$, $S = 0.768$ for 1086 parameters and 58142 reflections, 15539 unique [R(int) = 0.0674], of which 6194 with $I > 2\sigma(I)$, max positive and negative peaks in ΔF map 1.101, $-0.776 \text{ e}\cdot\text{\AA}^{-3}$.

Results and discussion

Crystal and molecular structure of

cis-(PPh_3)₂Pt{1-MeTy(-H)}(1-MeCy, N⁴)[NO₃] (2a)

We have recently shown that *cis*-(PPh_3)₂Pt(μ -OH)₂(NO₃)₂ reacts with 1-MeTy, in CH₂Cl₂, DMF or CH₃CN, affording the neutral complex *cis*-(PPh_3)₂Pt{1-MeTy(-H)}(ONO₂) in which the thyminato ligand is N(3) platinated and the nitrate group acts as monodentate ligand.⁴ Addition of one equivalent of 1-MeCy leads to the mixed complex *cis*-(PPh_3)₂Pt{1-MeTy(-H)}(1-MeCy, N³)[NO₃] (**1a**), resulting in the immediate replacement of the nitrate ligand. The deprotonated 1-MeTy and the neutral 1-MeCy, are both platinated at the N(3) atom. In a few days at room temperature in chlorinated solvents, DMSO or DMF, **1a** converts into the isomer *cis*-(PPh_3)₂Pt{1-MeTy(-H)}(1-MeCy, N⁴)[NO₃], **2a**, in which the cytosine is coordinated to the metal through the exocyclic N(4) atom, as shown by multinuclear NMR studies in solution and now confirmed in the solid state.

The X-ray structural determination of **2a** shows the presence in the unit cell of two crystallographic independent complexes (A and B), disordered nitrate anions and a lattice water molecule. The metal ion has a square planar coordination geometry achieved through the phosphorous atoms and the nitrogen donors of the nucleobases. The thyminato is bound through the endocyclic N(3) atom, the cytosine through the deprotonated amino group N(4) (Fig. 1 and 2 and Table 1).

The atoms of the coordination N₂P₂ plane are almost coplanar with max deviations of $\pm 0.02 \text{ \AA}$. The two complexes are conformationally very similar, differing slightly in the orientation of the nucleobase and phenyl rings. The position of the thymine N(1) nitrogen atom was not definitely assigned, having a pseudo two-fold axis that does not allow differentiation on the Δ Fourier




Fig. 1 ORTEP drawing of complex cation A (head–tail orientation of the bases) with indication of the intramolecular H-bond and π – π stacking interactions.





Fig. 2 ORTEP drawing of complex cation B (head–head orientation, the cytosine and thymine bases are labelled as “d” and “u”, respectively).

map of the endocyclic N(1) and C(5) atoms. Based on the thermal parameters values and bond distances, we tentatively assigned the N(1) thymine atom in the two complexes corresponding to the two possible head-to-head and head-to-tail conformational isomers present in solution. In the head-to-head (*hh*) and head-to-tail (*ht*) conformations the methyl group on cytosine and thymine N1 nitrogen atoms lie on the same and on the opposite side, respectively, of the P_2Pt plane (see Scheme 2). On the other hand, in both molecules the hydrogen at the cytosine N(3) atom (close to the metal) is indicative of a *syn* isomer.⁴



Scheme 2

The bond lengths and angles in the two independent complexes (not highly accurate) fall in a wide range, comparable within 2 – 3σ (Table 1). These data indicate that the Pt–N(3)(thyminato) bond distances (2.095(8) and 2.045(9) Å) appear shorter with respect to the Pt–N(4)(cytosine) ones (2.110(9) and 2.089(8) Å). The coordination bond angle C(4c)–N(4c)–Pt(1) at cytosine is also very similar in the two complexes with a mean value of $127.1(9)^\circ$. The N(3t)–Pt–N(4c) angle, $87.2(3)$ and $86.5(3)^\circ$ in complexes A and B, respectively, is narrower in contrast to the P(1)–Pt–P(2) one that average to 96.3° , likely induced by steric requirements. The thymine base is oriented almost normal to the coordination mean plane forming a dihedral angle of 82.88° (average value in the two complexes), while the cytosine plane is more bent, and its

Table 1 Coordination bond lengths and angles for the two independent complexes

	Complex A	Complex B
Pt(1)–N(3t)	2.095(8)	2.045(9)
Pt(1)–N(4c)	2.110(9)	2.089(8)
Pt(1)–P(1)	2.274(3)	2.281(3)
Pt(1)–P(2)	2.286(3)	2.272(3)
N(3t)–Pt(1)–N(4c)	87.2(3)	86.5(3)
N(3t)–Pt(1)–P(1)	88.9(2)	89.4(3)
N(3t)–Pt(1)–P(2)	174.8(3)	174.0(2)
N(4c)–Pt(1)–P(1)	175.9(2)	175.9(2)
N(4c)–Pt(1)–P(2)	87.9(2)	87.6(2)
P(1)–Pt(1)–P(2)	96.13(11)	96.48(12)
C(4c)–N(4c)–Pt(1)	127.2(9)	127.0(8)

ring forms a dihedral angle averaging to $78.6(2)^\circ$. This favours the formation of an intramolecular H bond between the N(3)–H and the thymine oxygen (see Fig. 1, mean values of N \cdots O distances and N–H \cdots O angles of *ca.* 2.86 Å and 161° , respectively).

The species are stabilized by two intramolecular π – π interactions given that two phosphine phenyl rings are oriented to stack with the model nucleobases. The centroid-to-centroid distance is shorter for the phenyl–thymine coupling, 3.394(9) Å with a dihedral angle of 14.9° (3.413(9) Å and 14.77° in complex B), in comparison to the values measured for the phenyl–cytosine pair, of 3.657(10) Å and 26.46° (3.887(9) Å, 30.39° in complex B). An additional intramolecular π – π interaction (not indicated in Fig. 1 and 2) is realized between two, almost parallel, phenyl groups.

The crystal packing evidences both the complexes, located near a center of symmetry, forming pairs of molecules connected by H-bonds occurring between the N(4)–H cytosine donor with the thymine oxygen O(2) of the symmetry related species, the N \cdots O distance being 2.992 Å (Fig. 3 and Table 2). A similar arrangement is also detected for the other independent complex B involving N(4)–H with O(4) with a more labile interaction of 3.102 Å.




Fig. 3 Crystal packing showing the pairing of complexes about an inversion center with an indication of H-bonds (phosphine phenyl groups not shown for clarity).

Most of the structural features in complexes A and B are similar to those previously described in the complex $cis\text{-}[(\text{PPh}_3)_2\text{Pt}\{1\text{-MeCy}(-\text{H}), \text{N}^4\}]^+$,¹¹ although in this species the coordination distances showed a reverse trend, being the Pt–N(3)

bond length is significantly longer (2.100(3) Å) than the Pt–N(4a) one (2.061(4) Å). Similar to the structure reported here, a N(4)–H \cdots N(3) hydrogen bond occurs between the bases along with intramolecular π – π stacking, confirming that the PPh₃ derivatives appear stabilized by these interactions.

On the other hand the different crystal packing of $cis\text{-}[(\text{PPh}_3)_2\text{Pt}\{1\text{-MeCy}(-\text{H}), \text{N}^4\}]^+$ (a polymer built by a H-bonding scheme) and of **2a** (pair of molecules, see above) could explain the differences observed in the coordination distance values.

Characterisation of $cis\text{-}[(\text{PMePh}_2)_2\text{Pt}\{1\text{-MeTy}(-\text{H})\text{-}(1\text{-MeCy}, \text{N}^3)]\text{NO}_3$ (**1b**)

With a procedure analogous to that used for the PPh₃ derivative,⁴ the thymine complex $cis\text{-}(\text{PMePh}_2)_2\text{Pt}\{1\text{-MeTy}(-\text{H})\}(\text{ONO}_2)$ was prepared. The ¹H and ³¹P NMR data of the new compound, in CDCl₃ and DMSO, strongly support a structure in which the nitrate group acts as monodentate ligand, as found in the PPh₃ analogue. Addition of one equivalent of 1-MeCy affords the mixed complex $cis\text{-}[(\text{PMePh}_2)_2\text{Pt}\{1\text{-MeTy}(-\text{H})\}(1\text{-MeCy}, \text{N}^3)]\text{NO}_3$ (**1b**). The spectroscopic analysis of the isolated product is consistent with the presence of the deprotonated 1-MeTy and the neutral 1-MeCy, both platinated at the N(3) atom. Due to the different orientations of the nucleobases with respect to the metal coordination plane (Scheme 3), two conformers are expected.




Table 2 Intra- and inter-molecular hydrogen bonds

D–H	d(D–H)	d(H \cdots A)	<DHA	d(D \cdots A)	A	Symmetry
Complex A						
N(3c)–H3c	0.860	2.039	163.63	2.875	O(4t)	
N(4c)–H4c	0.860	2.188	155.45	2.992	O(2t)	$[-x, -y, -z + 1]$
Complex B						
N(3d)–H3d	0.860	2.020	158.68	2.838	O(2u)	
N(4d)–H4d	0.860	2.310	153.18	3.102	O(4u)	$[-x + 1, -y + 1, -z + 2]$

Note: In molecule B the cytosine and thymine bases are labelled as “d” and “u”, respectively.




Fig. 4 $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum of **1b** (central part) in CDCl_3 at 25°C .

In this Scheme, the methyl group, bound to the N(1) atom in 1-MeCy and 1-MeTy, is oriented in opposite directions in the *ht* conformer. Accordingly, the ^{31}P NMR spectrum of **1b** in CDCl_3 is characterised by two partially overlapped AB multiplets, of relative intensities 1.5 : 1 (Fig. 4). Heterocorrelate ^{31}P and ^{15}N experiments indicate that the two phosphorous resonances at lower field, having $^1J_{\text{PPt}} = 3249$ Hz, are attributable to the phosphine in a *trans* position to the thymine ligand.

In the ^1H NMR spectrum each nucleobase shows two sets of resonances having relative intensities of 1.5 : 1, whose attribution (see Experimental) was obtained through a COSY experiment. The assignments of the cytosine NH_2 protons were obtained through inverse detected $^1\text{H},^{15}\text{N}$ heteronuclear multiple bond coherence experiments (HMBC). In CDCl_3 , the NH_2 resonances are observed at δ 8.69 and 8.60 ppm for the major conformer, and at δ 8.75 and 8.63 ppm for the other. As shown in Figure A in the ESI,[†] the couple of proton signals at lower field correlate with the same N(4) nucleus (^{15}N at $\delta = -271$ ppm), whereas the protons at δ 8.69 and 8.60 ppm correlate with the ^{15}N signal at $\delta = -270$ ppm ($^1J_{\text{NH}}$ in the range 80–90 Hz).

The NMR spectra of **1b** in CDCl_3 , exhibit other very weak resonances, whose attribution remains uncertain. Similar results were obtained in $\text{DMSO}-d_6$ in which, however, complex **1b** appears in equilibrium with the tautomeric species *cis*-[(PMePh₂)₂Pt{1-MeTy(-H)}(1-MeCy, N⁴)]⁺ (**2b**) as clearly indicated by the presence of a weak singlet at δ 10.50 ppm, attributable to the N(3)H proton of the N(4)-coordinated cytosine, and by a weak AX multiplet ($\delta_{\text{P}} = -7.05$ and -8.42 with $^2J_{\text{PP}} = 23.4$ Hz) in the corresponding ^{31}P NMR spectrum. The relative intensities of the signals indicate that the iminoxo tautomer is *ca.* 7% of the isomeric mixture and its concentration does not change after several weeks at room temperature.

Since the formal migration of the metal from N(3) to the N(4) site of the cytosine occurs quantitatively in **1a**, but only to a minor extent in **1b**, the nature of the ancillary ligands plays an important role. The stabilisation of the cytosine ligand in its unusual iminoxo tautomeric form is clearly favoured by the bulkier PPh₃ molecules, probably for steric reasons. The platination of the cytosine at the N(4) position, in fact, permits less crowding around the metal being one of the two pyrimidinic rings relatively further away from the coordination center.

In this context, it is interesting to note that the diphosphine analogue, *cis*-[(dppf)Pt{1-MeTy(-H)}(1-MeCy, N³)]BF₄ (dppf = 1,1'-bis(diphenylphosphino)ferrocene),¹² undergoes a similar rearrangement of the cytosine in DMF solution, with a large predominance of the *cis*-[(dppf)Pt{1-MeTy(-H)}(1-MeCy, N⁴)]⁺ species present at the equilibrium. In that case, however, the solid isolated from the mixture turned out to be the starting complex.

Conclusion

The X-ray structure of the compound *cis*-[(PPh₃)₂Pt{1-MeTy(-H)}(1-MeCy, N⁴)]NO₃ here reported represents the first example of a phosphino complex in which the neutral 1-MeCy exhibits N(4)-coordination to a metal centre. The two crystallographic independent molecules have been modelled taking into account the conformational isomers previously characterised in solution.⁴ These complexes appear stabilized by strong intramolecular π - π interactions between the pyrimidinic rings and the phosphine phenyl substituents, allowing the formation of intra- and intermolecular hydrogen bonds.

The peculiar properties of the PPh₃ ligands in the stabilisation of the cytosine molecule in the iminoxo form stem from the

following observations: (i) the PMe_3 complex, $\text{cis}-[(\text{PMe}_3)_2\text{Pt}\{1\text{-MeTy}(-\text{H})\}(1\text{-MeCy}, N^3)]^+$, in solution slowly rearranges into the polynuclear cytosinate species $\text{cis}-[(\text{PMe}_3)_2\text{Pt}\{1\text{-MeCy}(-\text{H})\}]_n^{n+}$ ($n = 2, 3$) and free 1-MeTy;⁴ (ii) the tautomeric equilibrium is largely shifted toward the usual N(3)-coordination for PMePh_2 derivatives, in particular in chlorinated solvents; (iii) the opposite holds for the dppf analogue of **1a**. However, in spite of its thermodynamic stability, we were unable to isolate the iminoxo species, $\text{cis}-[(\text{dppf})\text{Pt}\{1\text{-MeTy}(-\text{H})\}(1\text{-MeCy}, N^4)]^+$.¹²

References

- 1 B. Lippert, *Coord. Chem. Rev.*, 2000, **200–202**, 487; B. Lippert, *Prog. Inorg. Chem.*, 2005, **54**, 385.
- 2 B. Lippert, H. Schollhorn and U. Thewald, *J. Am. Chem. Soc.*, 1986, **108**, 6616; F. Pichierri, D. Holthenrich, E. Zangrando and L. Randaccio, *J. Biol. Inorg. Chem.*, 1996, **1**, 439; J. Müller, E. Zangrando, N. Pahlke, E. Freisinger, L. Randaccio and B. Lippert, *Chem. Eur. J.*, 1998, **4**, 397; P. J. Sanz Miguel, P. Lax, M. Willermann and B. Lippert, *Inorg. Chim. Acta*, 2004, **357**, 4552.
- 3 B. Lippert, *Prog. Inorg. Chem.*, 1989, **1**, 371.
- 4 D. Montagner, E. Zangrando and B. Longato, *Inorg. Chim. Acta*, 2009, **362**, 725.
- 5 B. Longato, G. Bandoli and A. Dolmella, *Eur. J. Inorg. Chem.*, 2004, 1092.
- 6 T. J. Kistenmacher, M. Rossi, J. P. Caradonna and L. G. Marzilli, *Adv. Mol. Relax. Interact. Processes*, 1979, **15**, 119.
- 7 L. Schenetti, A. Mucci and B. Longato, *J. Chem. Soc., Dalton Trans.*, 1996, 299.
- 8 Z. Otwinowski, and W. Minor, "Processing of X-ray Diffraction Data Collected in Oscillation Mode". in *Methods in Enzymology*, ed. C. W. Carter, Jr. and R. M. Sweet, Volume 276, New York, Academic Press, 1997, 307.
- 9 *SHELX97 Programs for Crystal Structure Analysis (Release 97–2)*, G. M. Sheldrick, University of Göttingen, Germany, 1998.
- 10 L. J. Farrugia, *J. Appl. Crystallogr.*, 1999, **32**, 837.
- 11 B. Longato, D. Montagner and E. Zangrando, *Inorg. Chem.*, 2006, **45**, 8179.
- 12 G. Bandoli, G. Trovò, A. Dolmella and B. Longato, *Inorg. Chem.*, 1992, **31**, 3169.